

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number
WO 01/46256 A2

(51) International Patent Classification⁷: **C07K 14/47**

(21) International Application Number: **PCT/US00/34919**

(22) International Filing Date:
21 December 2000 (21.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/172,968 21 December 1999 (21.12.1999) US
60/172,066 23 December 1999 (23.12.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:
US 60/172,968 (CIP)
Filed on 21 December 1999 (21.12.1999)
US 60/172,066 (CIP)
Filed on 23 December 1999 (23.12.1999)

(71) Applicant (for all designated States except US): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **TANG, Y., Tom** [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **BANDMAN, Olga** [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). **HILLMAN, Jennifer, L.** [US/US]; 230 Monrow Drive #17, Mountain View, CA 94040 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **LU,**

Dyung, Aina, M. [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **AZIMZAI, Yalda** [US/US]; 5518 Boulder Canyon Drive, Castro Valley, CA 94552 (US). **YANG, Junming** [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US). **BURFORD, Neil** [US/US]; 105 Wildwood Circle, Durham, CT 06522 (US). **AU-YOUNG, Janice** [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). **REDDY, Roopa** [IN/US]; 1233 W. McKinley Avenue #3, Sunnyvale, CA 94086 (US).

(74) Agents: **HAMLET-COX, Diana** et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **VESICLE TRAFFICKING PROTEINS**

(57) Abstract: The invention provides human vesicle trafficking proteins (VETRP) and polynucleotides which identify and encode VETRP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing disorders associated with expression of VETRP.



WO 01/46256 A2

VESICLE TRAFFICKING PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of vesicle trafficking proteins and to the use of these sequences in the diagnosis, treatment, and prevention of vesicle trafficking disorders, autoimmune/inflammatory disorders, and cancer, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of vesicle trafficking proteins.

BACKGROUND OF THE INVENTION

Eukaryotic cells are bound by a lipid bilayer membrane and subdivided into functionally distinct, membrane-bound compartments. The membranes maintain the essential differences between the cytosol, the extracellular environment, and the luminal space of each intracellular organelle. As lipid membranes are highly impermeable to most polar molecules, transport of essential nutrients, metabolic waste products, cell signaling molecules, macromolecules, and proteins across lipid membranes and between organelles must be mediated by a variety of transport-associated molecules.

Integral membrane proteins, secreted proteins, and proteins destined for the lumen of organelles are synthesized within the endoplasmic reticulum (ER), delivered to the Golgi complex for post-translational processing and sorting, and then transported to specific intracellular and extracellular destinations. Material is internalized from the extracellular environment by endocytosis, a process essential for transmission of neuronal, metabolic, and proliferative signals; uptake of many essential nutrients; and defense against invading organisms. This intracellular and extracellular movement of protein molecules is termed vesicle trafficking. Trafficking is accomplished by the packaging of protein molecules into specialized vesicles which bud from the donor organelle membrane and fuse to the target membrane (Rothman, J.E and F.T. Wieland (1996) Science 272:227-234).

The transport of proteins across the ER membrane involves a process that is similar in bacteria, yeast, and mammals (Gorlich, D. et al. (1992) Cell 71: 489-503). In mammalian systems, transport is initiated by the action of a cytoplasmic signal recognition particle (SRP) which recognizes a signal sequence on a growing, nascent polypeptide and binds the polypeptide and its ribosome complex to the ER membrane through an SRP receptor located on the ER membrane. The signal peptide is cleaved and the ribosome complex, together with the attached polypeptide, becomes membrane bound. The polypeptide is subsequently translocated across the ER membrane and into a vesicle (Blobel, G. and B. Dobberstein (1975) J. Cell Biol. 67:852-862).

Proteins implicated in the translocation of polypeptides across the ER membrane in yeast

include SEC61p, SEC62p, and SEC63p. Mutations in the genes encoding these proteins lead to defects in the translocation process. SEC61 may be of particular importance since certain mutations in the gene for this protein inhibit the translocation of many proteins (Gorlich, supra).

Mammalian homologs of yeast SEC61 (mSEC61) have been identified in dog and rat (Gorlich, supra). Mammalian SEC61 is also structurally similar to SECYp, the bacterial cytoplasmic membrane translocation protein. mSEC61 is found in tight association with membrane-bound ribosomes. This association is induced by membrane-targeting of nascent polypeptide chains and is weakened by dissociation of the ribosomes into their constituent subunits. mSEC61 is postulated to be a component of a putative protein-conducting channel, located in the ER membrane, to which nascent polypeptides are transferred following the completion of translation by ribosomes (Gorlich, supra).

Several steps in the transit of material along the secretory and endocytic pathways requires the formation of transport vesicles. Specifically, vesicles form at the transitional endoplasmic reticulum (tER), the rim of Golgi cisternae, the face of the Trans-Golgi Network (TGN), the plasma membrane (PM), and tubular extensions of the endosomes. Vesicle formation occurs when a region of membrane buds off from the donor organelle. The membrane-bound vesicle contains proteins to be transported and is surrounded by a proteinaceous coat, the components of which are recruited from the cytosol. Vesicle formation begins with the budding of a vesicle out of a donor organelle. The initial budding and coating processes are controlled by a cytosolic ras-like GTP-binding protein, ADP-ribosylating factor (Arf), and adapter proteins (AP). Different isoforms of both Arf and AP are involved at different sites of budding. For example, Arfs 1, 3, and 5 are required for Golgi budding, Arf4 for endosomal budding, and Arf6 for plasma membrane budding. Two different classes of coat protein have also been identified. Clathrin coats form on vesicles derived from the TGN and PM, whereas coatamer (COP) coats form on vesicles derived from the ER and Golgi (Mellman, I. (1996) *Annu. Rev. Cell Dev. Biol.* 12:575-625).

Vesicle formation begins when an adapter protein (AP) interacts with cargo proteins within the donor membrane and recruits clathrin to the bud site. APs are heterotetrameric complexes composed of two large chains (a, g, d, or e, and b), a medium chain (m), and a small chain (s). Clathrin binds to APs via the carboxy-terminal appendage domain of the b-adaptin subunit (Le Bourgne, R. and B. Hoflack (1998) *Curr. Opin. Cell. Biol.* 10:499-503). AP-1 functions in protein sorting from the TGN and endosomes to compartments of the endosomal/lysosomal system. AP-2 functions in clathrin-mediated endocytosis at the plasma membrane, while AP-3 is associated with endosomes and/or the TGN and recruits integral membrane proteins for transport to lysosomes and lysosome-related organelles. The recently isolated AP-4 complex localizes to the TGN or a neighboring compartment and may play a role in sorting events thought to take place in post-Golgi

compartments (Dell'Angelica, E.C. et al. (1999) *J. Biol. Chem.* 274:7278-7285). Cytosolic GTP-bound Arf is also incorporated into the vesicle as it forms. Another GTP-binding protein, dynamin, forms a ring complex around the neck of the forming vesicle and provides the mechanochemical force required to release the vesicle from the donor membrane. The coated vesicle complex is then
5 transported through the cytosol. During the transport process, Arf-bound GTP is hydrolyzed to GDP and the coat dissociates from the transport vesicle (West, M.A. et al. (1997) *J. Cell Biol.* 138:1239-1254).

Coat protein (COP) coats form on the ER and Golgi. COP coats can further be distinguished as COPI, involved in retrograde traffic through the Golgi to the ER, and COPII, involved in
10 anterograde traffic from the ER to the Golgi. The COP coat consists of two major components, a GTP-binding protein (Arf or Sar) and coat protomer (coatomer). Coatomer is an equimolar complex of seven proteins, termed alpha-, beta-, beta'-, gamma-, delta-, epsilon- and zeta-COP. The coatomer complex binds to dilysine motifs contained on the cytoplasmic tails of integral membrane proteins. These include the dilysine-containing retrieval motif of membrane proteins of the ER and
15 dibasic/diphenylamine motifs of members of the p24 family. The p24 family of type I membrane proteins represent the major membrane proteins of COPI vesicles (Harter, C. and F.T. Wieland (1998) *Proc. Natl. Acad. Sci. USA* 95:11649-11654).

Vesicles can undergo homotypic or heterotypic fusion. Molecules required for appropriate targeting and fusion of vesicles include proteins in the vesicle membrane, the target membrane, and
20 proteins recruited from the cytosol. During budding of the vesicle from the donor compartment, an integral membrane protein, VAMP (vesicle-associated membrane protein) is incorporated into the vesicle. Soon after the vesicle uncoats, a cytosolic prenylated GTP-binding protein, Rab, is inserted into the vesicle membrane. The amino acid sequence of Rab proteins reveals conserved GTP-binding domains characteristic of Ras superfamily members. In the vesicle membrane, GTP-bound Rab
25 interacts with VAMP. Once the vesicle reaches the target membrane, a GTPase activating protein (GAP) in the target membrane converts the Rab protein to the GDP-bound form. A cytosolic protein, guanine-nucleotide dissociation inhibitor (GDI) then removes GDP-bound Rab from the vesicle membrane. Several Rab isoforms have been identified and appear to associate with specific compartments within the cell. For example, Rabs 4, 5, and 11 are associated with the early
30 endosome, whereas Rabs 7 and 9 associate with the late endosome. These differences may provide selectivity in the association between vesicles and their target membranes (Novick, P. and M. Zerial (1997) *Cur. Opin. Cell Biol.* 9:496-504).

Docking of the transport vesicle with the target membrane involves the formation of a complex between the vesicle SNAP receptor (v-SNARE), target membrane (t-) SNAREs, and certain
35 other membrane and cytosolic proteins. Many of these other proteins have been identified although

their exact functions in the docking complex remain uncertain (Tellam, J.T. et al. (1995) J. Biol. Chem. 270:5857-5863; Hata, Y. and T.C. Sudhof (1995) J. Biol. Chem. 270:13022-13028).

N-ethylmaleimide sensitive factor (NSF) and soluble NSF-attachment protein (α -SNAP and β -SNAP) are two such proteins that are conserved from yeast to man and function in most intracellular

5 membrane fusion reactions. Sec1 represents a family of yeast proteins that function at many different stages in the secretory pathway including membrane fusion. Recently, mammalian homologs of Sec1, called Munc-18 proteins, have been identified (Katagiri, H. et al. (1995) J. Biol. Chem. 270:4963-4966; Hata et al. supra).

The SNARE complex involves three SNARE molecules, one in the vesicular membrane and
10 two in the target membrane. Together they form a rod-shaped complex of four α -helical coiled-coils.

The membrane anchoring domains of all three SNAREs project from one end of the rod. This complex is similar to the rod-like structures formed by fusion proteins characteristic of the enveloped viruses, such as myxovirus, influenza, filovirus (Ebola), and the HIV and SIV retroviruses. (Skehel,

J.J. and D.C. Wiley (1998) Cell 95:871-874). It has been proposed that the SNARE complex is

15 sufficient for membrane fusion, suggesting that the proteins which associate with the complex provide regulation over the fusion event (Weber, T. et al. (1998) Cell 92:759-772). For example, in neurons, which exhibit regulated exocytosis, docked vesicles do not fuse with the presynaptic membrane until depolarization, which leads to an influx of calcium (Bennett, M.K. and R.H. Scheller (1994) Annu. Rev. Biochem. 63:63-100). Synaptotagmin, an integral membrane protein in the

20 synaptic vesicle, associates with the t-SNARE syntaxin in the docking complex. Synaptotagmin binds calcium in a complex with negatively charged phospholipids, which allows the cytosolic SNAP protein to displace synaptotagmin from syntaxin and fusion to occur. Thus, synaptotagmin is a negative regulator of fusion in the neuron (Littleton, J.T. et al. (1993) Cell 74:1125-1134). The most abundant membrane protein of synaptic vesicles appears to be the glycoprotein synaptophysin,, a 38
25 kDa protein with four transmembrane domains. Although the function of synaptophysin is not known, its calcium-binding ability, tyrosine phosphorylation, and widespread distribution in neural tissues suggest a potential role in neurosecretion (Bennett, supra).

The transport of proteins into and out of vesicles relies on interactions between cell membranes and a supporting membrane cytoskeleton consisting of spectrin and other proteins. A
30 large family of related proteins called ankyrins participate in the transport process by binding to the membrane skeleton protein spectrin and to a protein in the cell membrane called band 3, a component of an anion channel in the cell membrane. Ankyrins therefore function as a critical link between the cytoskeleton and the cell membrane.

Originally found in association with erythroid cells, ankyrins are also found in other tissues
35 as well (Birkenmeier, C.S. et al. (1993) J. Biol. Chem. 268:9533-9540). Ankyrins are large proteins

(~1800 amino acids) containing an N-terminal, 89 kDa domain that binds the cell membrane proteins band 3 and tubulin, a central 62 kDa domain that binds the cytoskeletal proteins spectrin and vimentin, and a C-terminal, 55 kDa regulatory domain that functions as a modifier of the binding activities of the other two domains. Individual genes for ankyrin are able to produce multiple ankyrin isoforms by various insertions and deletions. These isoforms are of nearly identical size but may have different functions. In addition, smaller transcripts are produced which are missing large regions of the coding sequences from the N-terminal (band 3 binding), and central (spectrin binding) domains. The existence of such a large family of ankyrin proteins and the observation that more than one type of ankyrin may be expressed in the same cell type suggests that ankyrins may have more specialized functions than simply binding the membrane skeleton to the plasma membrane (Birkenmeier, *supra*).

In humans, two isoforms of ankyrin are expressed, alternatively, in developing erythroids and mature erythroids, respectively (Lambert, S. et al. (1990) Proc. Natl. Acad. Sci. USA 87:1730-1734). A deficiency in erythroid spectrin and ankyrin has been associated with the hemolytic anemia, hereditary spherocytosis (Coetzer, T.L. et al. (1988) New Engl. J. Med. 318:230-234).

Correct trafficking of proteins is of particular importance for the proper function of epithelial cells, which are polarized into distinct apical and basolateral domains containing different cell membrane components such as lipids and membrane-associated proteins. Certain proteins are flexible and may be sorted to the basolateral or apical side depending upon cell type or growth conditions. For example, the kidney anion exchanger (kAE1) can be retargeted from the apical to the basolateral domain if cells are plated at higher density. The protein kanadaplin was isolated as a protein which binds to the cytoplasmic domain of kAE1. It also colocalizes with kAE1 in vesicles, but not in the membrane, suggesting that kanadaplin's function is to guide kAE1-containing vesicles to the basolateral target membrane (Chen, J. et al. (1998) J. Biol. Chem. 273:1038-1043).

Vesicle trafficking is crucial in the process of neurotransmission. Synaptic vesicles carry neurotransmitter molecules from the cytoplasm of a neuron to the synapse. Rab3's are a family of GTP-binding proteins located on synaptic vesicles. The RIM family of proteins are thought to be effectors for Rab3's (Wang, Y. et al. (2000) J. Biol. Chem. 275:20033-20044). Rabphilin-3 is a synaptic vesicle protein. Granophilins are proteins with homology to rabphilins, and may have a unique role in exocytosis (Wang, J. et al. (1999) J. Biol. Chem. 274:28542-28548).

The etiology of numerous human diseases and disorders can be attributed to defects in the trafficking of proteins to organelles or the cell surface. Defects in the trafficking of membrane-bound receptors and ion channels are associated with cystic fibrosis (cystic fibrosis transmembrane conductance regulator; CFTR), glucose-galactose malabsorption syndrome (Na⁺/glucose cotransporter), hypercholesterolemia (low-density lipoprotein (LDL) receptor), and forms of diabetes

mellitus (insulin receptor). Abnormal hormonal secretion is linked to disorders including diabetes insipidus (vasopressin), hyper- and hypoglycemia (insulin, glucagon), Grave's disease and goiter (thyroid hormone), and Cushing's and Addison's diseases (adrenocorticotrophic hormone; ACTH).

Cancer cells secrete excessive amounts of hormones or other biologically active peptides.

- 5 Disorders related to excessive secretion of biologically active peptides by tumor cells include: fasting hypoglycemia due to increased insulin secretion from insulinoma-islet cell tumors; hypertension due to increased epinephrine and norepinephrine secreted from pheochromocytomas of the adrenal medulla and sympathetic paraganglia; and carcinoid syndrome, which includes abdominal cramps, diarrhea, and valvular heart disease, caused by excessive amounts of vasoactive substances
- 10 (serotonin, bradykinin, histamine, prostaglandins, and polypeptide hormones) secreted from intestinal tumors. Ectopic synthesis and secretion of biologically active peptides (peptides not expected from a tumor) includes ACTH and vasopressin in lung and pancreatic cancers; parathyroid hormone in lung and bladder cancers; calcitonin in lung and breast cancers; and thyroid-stimulating hormone in medullary thyroid carcinoma.

- 15 Various human pathogens alter host cell protein trafficking pathways to their own advantage. For example, the HIV protein Nef downregulates cell-surface expression of CD4 molecules by accelerating their endocytosis through clathrin coated pits. This function of Nef is important for the spread of HIV from the infected cell (Harris, M. (1999) Curr. Biol. 9:R449-R461). A recently identified human protein, Nef-associated factor 1 (Naf1), a protein with four extended coiled-coil
- 20 domains, has been found to associate with Nef. Overexpression of Naf1 increased cell surface expression of CD4, an effect which could be suppressed by Nef (Fukushi, M. et al. (1999) FEBS Lett. 442:83-88).

- The discovery of new vesicle trafficking proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis,
- 25 prevention, and treatment of vesicle trafficking disorders, autoimmune/inflammatory disorders, and cancer, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of vesicle trafficking proteins.

SUMMARY OF THE INVENTION

- 30 The invention features purified polypeptides, vesicle trafficking proteins, referred to collectively as "VETRP" and individually as "VETRP-1," "VETRP-2," "VETRP-3," "VETRP-4," "VETRP-5," "VETRP-6," "VETRP-7," "VETRP-8," "VETRP-9," "VETRP-10," "VETRP-11," "VETRP-12," "VETRP-13," "VETRP-14," "VETRP-15," "VETRP-16," "VETRP-17," "VETRP-18," "VETRP-19," "VETRP-20," "VETRP-21," "VETRP-22," and "VETRP-23." In one aspect, the
- 35 invention provides an isolated polypeptide comprising an amino acid sequence selected from the

group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-23.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-23. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:24-46.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the

polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said

target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional VETRP, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional VETRP, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. The

method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional VETRP, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:24-46, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding VETRP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of VETRP.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding VETRP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and

polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

5 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

10 It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

15 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing
20 the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"VETRP" refers to the amino acid sequences of substantially purified VETRP obtained from
25 any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of VETRP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of VETRP either by directly interacting with
30 VETRP or by acting on components of the biological pathway in which VETRP participates.

An "allelic variant" is an alternative form of the gene encoding VETRP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to
35 allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides.

Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding VETRP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as VETRP or a polypeptide with at least one functional characteristic of VETRP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding VETRP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding VETRP. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent VETRP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of VETRP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to a sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

“Amplification” relates to the production of additional copies of a nucleic acid sequence.

Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of VETRP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of VETRP either by directly interacting with VETRP or by acting on components of the biological pathway in which VETRP participates.

The term “antibody” refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind VETRP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or

oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize
5 the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures
10 on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as
15 phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring
20 nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic"
25 refers to the capability of the natural, recombinant, or synthetic VETRP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement,
30 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding VETRP or fragments of VETRP may
35 be employed as hybridization probes. The probes may be stored in freeze-dried form and may be

associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

Original Residue	Conservative Substitution
Ala	Gly, Ser
Arg	His, Lys
Asn	Asp, Gln, His
Asp	Asn, Glu
Cys	Ala, Ser
Gln	Asn, Glu, His
Glu	Asp, Gln, His
Gly	Ala
His	Asn, Arg, Gln, Glu
Ile	Leu, Val
Leu	Ile, Val
Lys	Arg, Gln, Glu
Met	Leu, Ile
Phe	His, Met, Leu, Trp, Tyr
Ser	Cys, Thr
Thr	Ser, Val
Trp	Phe, Tyr
Tyr	His, Phe, Trp
Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the

absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of VETRP or the polynucleotide encoding VETRP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:24-46 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:24-46, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:24-46 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:24-46 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:24-46 and the region of SEQ ID NO:24-46 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-23 is encoded by a fragment of SEQ ID NO:24-46. A fragment of SEQ ID NO:1-23 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-23. For example, a fragment of SEQ ID NO:1-23 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-23. The precise length of a fragment of SEQ ID NO:1-23 and the region of SEQ ID NO:1-23 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended

purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

5 "Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps
10 in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of
15 molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent
20 similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at
25 <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>.
30 The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

35 *Reward for match: 1*

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

5 *Word Size: 11*

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous
10 nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode
15 similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a
20 standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default
25 parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by
30 CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

35 *Matrix: BLOSUM62*

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and

conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of VETRP which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of VETRP which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of VETRP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of VETRP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an VETRP may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of VETRP.

"Probe" refers to nucleic acid sequences encoding VETRP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule.

Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have

been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding VETRP, or fragments thereof, or VETRP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides

by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook, J. et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example,

an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human vesicle trafficking proteins (VETRP), the polynucleotides encoding VETRP, and the use of these compositions for the diagnosis, treatment, or prevention of vesicle trafficking disorders, autoimmune/inflammatory disorders, and cancer.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding VETRP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each VETRP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each VETRP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6

shows homologous sequences as identified by BLAST analysis along with relevant citations, all of which are expressly incorporated by reference herein in their entirety; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding VETRP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:24-46 and to distinguish between SEQ ID NO:24-46 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express VETRP as a fraction of total tissues expressing VETRP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing VETRP as a fraction of total tissues expressing VETRP. Column 5 lists the vectors used to subclone each cDNA library. Of particular note is the expression of SEQ ID NO:25 in nervous tissue. SEQ ID NO:41 is noted for its expression in both cancer and reproductive tissue, and SEQ ID NO:43 is expressed in cancer and nervous tissue.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding VETRP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses VETRP variants. A preferred VETRP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the VETRP amino acid sequence, and which contains at least one functional or structural characteristic of VETRP.

SEQ ID NO:31 maps to chromosome 12 within the interval from 70.60 to 76.50 centiMorgans, and to chromosome 1 within the interval from 159.60 to 164.10 centiMorgans. SEQ ID NO:36 maps to chromosome 3 within the interval from 129.00 to 131.80 centiMorgans, and to chromosome 4 within the interval from 86.00 to 91.90 centiMorgans. SEQ ID NO:38 maps to chromosome 6 within the interval from the p-terminus to 27.10 centiMorgans. SEQ ID NO:42 maps to chromosome 2 within the interval from 233.10 to 236.10 centiMorgans. SEQ ID NO:44 maps to chromosome 5 within the interval from 61.10 to 69.60 centiMorgans, to chromosome 11 within the interval from 117.90 to 123.50 centiMorgans, and to chromosome 17 within the interval from 99.30 to 103.70 centiMorgans.

The invention also encompasses polynucleotides which encode VETRP. In a particular

embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:24-46, which encodes VETRP. The polynucleotide sequences of SEQ ID NO:24-46, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding VETRP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding VETRP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:24-46 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:24-46. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of VETRP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding VETRP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring VETRP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode VETRP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring VETRP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding VETRP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding VETRP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode VETRP and VETRP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell

systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding VETRP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:24-46 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) *Short Protocols in Molecular Biology*, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) *Molecular Biology and Biotechnology*, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding VETRP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region

of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode VETRP may be cloned in recombinant DNA molecules that direct expression of VETRP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express VETRP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter VETRP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or
5 improve the biological properties of VETRP, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of
10 DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby
15 maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding VETRP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively,
20 VETRP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid
25 sequence of VETRP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.)
30 The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active VETRP, the nucleotide sequences encoding VETRP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding
35 sequence in a suitable host. These elements include regulatory sequences, such as enhancers,

constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding VETRP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding VETRP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding VETRP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding VETRP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding VETRP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for

delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.)

5 The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding VETRP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding VETRP can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1
10 plasmid (Life Technologies). Ligation of sequences encoding VETRP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.)
15 When large quantities of VETRP are needed, e.g. for the production of antibodies, vectors which direct high level expression of VETRP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of VETRP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH
20 promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, et al. 1995, *supra*; Bitter, *supra*; and Scorer, *supra*.)

Plant systems may also be used for expression of VETRP. Transcription of sequences
25 encoding VETRP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, *supra*; Broglie, *supra*; and Winter, *supra*.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated
30 transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding VETRP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader
35 sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain

infective virus which expresses VETRP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of VETRP in cell lines is preferred. For example, sequences encoding VETRP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk⁻* and *apr⁻* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the

sequence encoding VETRP is inserted within a marker gene sequence, transformed cells containing sequences encoding VETRP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding VETRP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding VETRP and that express VETRP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of VETRP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on VETRP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding VETRP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding VETRP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding VETRP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence.

and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode VETRP may be designed to contain signal sequences which direct secretion of VETRP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding VETRP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric VETRP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of VETRP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the VETRP encoding sequence and the heterologous protein sequence, so that VETRP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled VETRP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

VETRP of the present invention or fragments thereof may be used to screen for compounds that specifically bind to VETRP. At least one and up to a plurality of test compounds may be screened for specific binding to VETRP. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

5 In one embodiment, the compound thus identified is closely related to the natural ligand of VETRP, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which VETRP binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the
10 compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express VETRP, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing VETRP or cell membrane fractions which contain VETRP are then contacted with a test compound and binding, stimulation, or inhibition of activity of either VETRP or
15 the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with VETRP, either in solution or affixed to a solid support, and detecting the binding of VETRP to the compound.
20 Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

VETRP of the present invention or fragments thereof may be used to screen for compounds
25 that modulate the activity of VETRP. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for VETRP activity, wherein VETRP is combined with at least one test compound, and the activity of VETRP in the presence of a test compound is compared with the activity of VETRP in the absence of the test compound. A change in the activity of VETRP in the presence of the test compound is
30 indicative of a compound that modulates the activity of VETRP. Alternatively, a test compound is combined with an in vitro or cell-free system comprising VETRP under conditions suitable for VETRP activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of VETRP may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding VETRP or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding VETRP may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding VETRP can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding VETRP is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress VETRP, e.g., by secreting VETRP in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of VETRP and vesicle trafficking proteins. In addition, the expression of VETRP is closely associated with reproductive tissue, nervous tissue, cancer and inflammation/trauma.

Therefore, VETRP appears to play a role in vesicle trafficking disorders, autoimmune/inflammatory disorders, and cancer. In the treatment of disorders associated with increased VETRP expression or activity, it is desirable to decrease the expression or activity of VETRP. In the treatment of disorders associated with decreased VETRP expression or activity, it is desirable to increase the expression or activity of VETRP.

Therefore, in one embodiment, VETRP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of VETRP. Examples of such disorders include, but are not limited to, a vesicle trafficking disorder, such as cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, diabetes mellitus, diabetes insipidus, hyper- and hypoglycemia, Grave's disease, goiter, Cushing's disease, and Addison's disease; gastrointestinal disorders including ulcerative colitis, gastric and duodenal ulcers; other conditions associated with abnormal vesicle trafficking, including acquired immunodeficiency syndrome (AIDS); allergies including hay fever, asthma, and urticaria (hives); autoimmune hemolytic anemia; proliferative glomerulonephritis; inflammatory bowel disease; multiple sclerosis; myasthenia gravis; rheumatoid and osteoarthritis; scleroderma; Chediak-Higashi and Sjogren's syndromes; systemic lupus erythematosus; toxic shock syndrome; traumatic tissue damage; and viral, bacterial, fungal, helminthic, and protozoal infections; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cancer, such as, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and

uterus.

In another embodiment, a vector capable of expressing VETRP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of VETRP including, but not limited to, those described above.

5 In a further embodiment, a composition comprising a substantially purified VETRP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of VETRP including, but not limited to, those provided above.

10 In still another embodiment, an agonist which modulates the activity of VETRP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of VETRP including, but not limited to, those listed above.

In a further embodiment, an antagonist of VETRP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of VETRP. Examples of such disorders include, but are not limited to, those vesicle trafficking disorders,
15 autoimmune/inflammatory disorders, and cancer described above. In one aspect, an antibody which specifically binds VETRP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express VETRP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding VETRP may be administered to a subject to treat or prevent a disorder associated with
20 increased expression or activity of VETRP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The
25 combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of VETRP may be produced using methods which are generally known in the art. In particular, purified VETRP may be used to produce antibodies or to screen libraries of
30 pharmaceutical agents to identify those which specifically bind VETRP. Antibodies to VETRP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

35 For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans,

and others may be immunized by injection with VETRP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to VETRP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of VETRP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to VETRP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce VETRP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for VETRP may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and

easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between VETRP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering VETRP epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for VETRP. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of VETRP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple VETRP epitopes, represents the average affinity, or avidity, of the antibodies for VETRP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular VETRP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the VETRP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of VETRP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of VETRP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

In another embodiment of the invention, the polynucleotides encoding VETRP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene

encoding VETRP. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding VETRP. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

5 In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 10 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et 15 al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding VETRP may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X- 20 linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475); cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial 25 hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. 30 (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in VETRP expression or regulation causes disease, the expression of VETRP from an appropriate population of transduced cells may alleviate the clinical manifestations 35 caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in VETRP are treated by constructing mammalian expression vectors encoding VETRP and introducing these vectors by mechanical means into VETRP-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) 5 ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of VETRP include, but are not 10 limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). VETRP may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the 15 tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. 20 and H.M. Blau, supra), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding VETRP from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver 25 polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with 30 respect to VETRP expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding VETRP under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are 35 commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc.*

Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding VETRP to cells which have one or more genetic abnormalities with respect to the expression of VETRP. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544; and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding VETRP to target cells which have one or more genetic abnormalities with respect to the expression of VETRP. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing VETRP to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a

cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned
5 herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to
10 deliver polynucleotides encoding VETRP to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA,
15 resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for VETRP into the alphavirus genome in place of the capsid-coding region results in the production of a large number of VETRP-coding RNAs and the synthesis of high levels of VETRP in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a
20 persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of VETRP into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of
25 manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly,
30 inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-
35 177.) A complementary sequence or antisense molecule may also be designed to block translation of

mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding VETRP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding VETRP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding VETRP. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide

sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased VETRP expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding VETRP may be therapeutically useful, and in the treatment of disorders associated with decreased VETRP expression or activity, a compound which specifically promotes expression of the polynucleotide encoding VETRP may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding VETRP is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding VETRP are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding VETRP. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient.

Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of VETRP, antibodies to VETRP, and mimetics, agonists, antagonists, or inhibitors of VETRP.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising VETRP or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, VETRP or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example VETRP or fragments thereof, antibodies of VETRP, and agonists, antagonists or inhibitors of VETRP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about $0.1 \mu\text{g}$ to $100,000 \mu\text{g}$, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind VETRP may be used for the diagnosis of disorders characterized by expression of VETRP, or in assays to monitor patients being treated with VETRP or agonists, antagonists, or inhibitors of VETRP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics.

Diagnostic assays for VETRP include methods which utilize the antibody and a label to detect VETRP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring VETRP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of VETRP expression. Normal or standard values for VETRP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to VETRP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of VETRP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding VETRP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of VETRP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of VETRP, and to monitor regulation of VETRP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding VETRP or closely related molecules may be used to identify nucleic acid sequences which encode VETRP. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding VETRP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the VETRP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:24-46 or from genomic sequences including promoters, enhancers, and introns of the VETRP gene.

Means for producing specific hybridization probes for DNAs encoding VETRP include the cloning of polynucleotide sequences encoding VETRP or VETRP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA

polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding VETRP may be used for the diagnosis of disorders associated with expression of VETRP. Examples of such disorders include, but are not limited to, a vesicle trafficking disorder, such as cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, diabetes mellitus, diabetes insipidus, hyper- and hypoglycemia, Grave's disease, goiter, Cushing's disease, and Addison's disease; gastrointestinal disorders including ulcerative colitis, gastric and duodenal ulcers; other conditions associated with abnormal vesicle trafficking, including acquired immunodeficiency syndrome (AIDS); allergies including hay fever, asthma, and urticaria (hives); autoimmune hemolytic anemia; proliferative glomerulonephritis; inflammatory bowel disease; multiple sclerosis; myasthenia gravis; rheumatoid and osteoarthritis; scleroderma; Chediak-Higashi and Sjogren's syndromes; systemic lupus erythematosus; toxic shock syndrome; traumatic tissue damage; and viral, bacterial, fungal, helminthic, and protozoal infections; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cancer, such as, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding VETRP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect

altered VETRP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding VETRP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding VETRP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding VETRP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of VETRP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding VETRP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding VETRP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding VETRP, or a fragment of a polynucleotide complementary to the polynucleotide encoding

VETRP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding VETRP may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding VETRP are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of VETRP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to

monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, antibodies specific for VETRP, or VETRP or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes

are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, supra). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be

obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for VETRP to quantify the levels of VETRP expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) Anal. Biochem. 270:103-111; Mendoz, L.G. et al. (1999) Biotechniques 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) Electrophoresis 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al.

(1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

5 In another embodiment of the invention, nucleic acid sequences encoding VETRP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during
10 chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J.
15 (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, e.g., Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic
20 map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding VETRP on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

25 In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery
30 techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to
35 translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, VETRP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes
5 between VETRP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with VETRP, or fragments thereof,
10 and washed. Bound VETRP is then detected by methods well known in the art. Purified VETRP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing
15 antibodies capable of binding VETRP specifically compete with a test compound for binding VETRP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with VETRP.

In additional embodiments, the nucleotide sequences which encode VETRP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on
20 properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific
25 embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/172,968 and U.S. Ser. No. 60/172,066 are hereby expressly incorporated by reference.

30 EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a
35 monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged

over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence

scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire

annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:24-46. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel, 1995, *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum \{length(Seq. 1), length(Seq. 2)\}}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the

entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

5 The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding VETRP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma,
10 cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Chromosomal Mapping of ABBR Encoding Polynucleotides

15 The cDNA sequences which were used to assemble SEQ ID NO:24-46 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:24-46 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available
20 from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

 The genetic map locations of SEQ ID NO:31, SEQ ID NO:36, SEQ ID NO:38, SEQ ID
25 NO:42, and SEQ ID NO:44 are described in The Invention as ranges, or intervals, of human chromosomes. More than one map location is reported for SEQ ID NO:31, SEQ ID NO:36, and SEQ ID NO:44, indicating that previously mapped sequences having similarity, but not complete identity, to SEQ ID NO:31, SEQ ID NO:36, and SEQ ID NO:44 were assembled into their respective clusters. The map position of an interval, in centiMorgans, is measured relative to the terminus of the
30 chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome
35 maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web

site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

VI. Extension of VETRP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:24-46 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviII cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%)

agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, the polynucleotide sequences of SEQ ID NO:24-46 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

VII. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:24-46 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 µCi of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and

compared.

VIII. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the
5 aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding
10 procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Skena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may
15 comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a
20 fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

25 Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40
30 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37 °C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and
35 incubated for 20 minutes at 85 °C to stop the reaction and degrade the RNA. Samples are purified

using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and

5 resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are
10 amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water
15 washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US
20 Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water.
25 Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60 °C followed by washes in 0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and
30 Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65 °C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is

incubated for about 6.5 hours at 60 °C. The arrays are washed for 10 min at 45 °C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45 °C in a second wash buffer (0.1X SSC), and dried.

Detection

5 Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each

spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

IX. Complementary Polynucleotides

Sequences complementary to the VETRP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring VETRP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of VETRP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the VETRP-encoding transcript.

X. Expression of VETRP

Expression and purification of VETRP is achieved using bacterial or virus-based expression systems. For expression of VETRP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express VETRP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of VETRP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding VETRP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, VETRP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on

immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from VETRP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10 and 16). Purified VETRP obtained by these methods can be used directly in the assays shown in Examples XI and XV.

XI. Demonstration of VETRP Activity

VETRP activity is measured by its inclusion in coated vesicles. VETRP can be expressed by transforming a mammalian cell line such as COS7, HeLa, or CHO with an eukaryotic expression vector encoding VETRP. Eukaryotic expression vectors are commercially available, and the techniques to introduce them into cells are well known to those skilled in the art. A small amount of a second plasmid, which expresses any one of a number of marker genes, such as β -galactosidase, is co-transformed into the cells in order to allow rapid identification of those cells which have taken up and expressed the foreign DNA. The cells are incubated for 48-72 hours after transformation under conditions appropriate for the cell line to allow expression and accumulation of VETRP and β -galactosidase.

Transformed cells are collected and cell lysates are assayed for vesicle formation. A non-hydrolyzable form of GTP, GTP γ S, and an ATP regenerating system are added to the lysate and the mixture is incubated at 37 °C for 10 minutes. Under these conditions, over 90% of the vesicles remain coated (Orsi, L. et al (1989) Cell 56:357-368). Transport vesicles are salt-released from the Golgi membranes, loaded under a sucrose gradient, centrifuged, and fractions are collected and analyzed by SDS-PAGE. Co-localization of VETRP with clathrin or COP coatamer is indicative of VETRP activity in vesicle formation. The contribution of VETRP in vesicle formation can be confirmed by incubating lysates with antibodies specific for VETRP prior to GTP γ S addition. The antibody will bind to VETRP and interfere with its activity, thus preventing vesicle formation.

In the alternative, VETRP activity is measured by its ability to alter vesicle trafficking pathways. Vesicle trafficking in cells transformed with VETRP is examined using fluorescence microscopy. Antibodies specific for vesicle coat proteins or typical vesicle trafficking substrates such as transferrin or the mannose-6-phosphate receptor are commercially available. Various cellular components such as ER, Golgi bodies, peroxisomes, endosomes, lysosomes, and the plasmalemma are examined. Alterations in the numbers and locations of vesicles in cells transformed with VETRP as compared to control cells are characteristic of VETRP activity.

XII. Functional Assays

VETRP function is assessed by expressing the sequences encoding VETRP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of VETRP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding VETRP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding VETRP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIII. Production of VETRP Specific Antibodies

VETRP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495); or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the VETRP amino acid sequence is analyzed using LASERGENE software

(DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

5 Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for
10 antipeptide and anti-VETRP activity by, for example, binding the peptide or VETRP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIV. Purification of Naturally Occurring VETRP Using Specific Antibodies

Naturally occurring or recombinant VETRP is substantially purified by immunoaffinity
15 chromatography using antibodies specific for VETRP. An immunoaffinity column is constructed by covalently coupling anti-VETRP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing VETRP are passed over the immunoaffinity column, and the column is
20 washed under conditions that allow the preferential absorbance of VETRP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/VETRP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and VETRP is collected.

XV. Identification of Molecules Which Interact with VETRP

25 VETRP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled VETRP, washed, and any wells with labeled VETRP complex are assayed. Data obtained using different concentrations of VETRP are used to calculate values for the number, affinity, and association of
30 VETRP with the candidate molecules.

Alternatively, molecules interacting with VETRP are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, Nature 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

VETRP may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT)
35 which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions

between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

Various modifications and variations of the described methods and systems of the invention
5 will be apparent to those skilled in the art without departing from the scope and spirit of the
invention. Although the invention has been described in connection with certain embodiments, it
should be understood that the invention as claimed should not be unduly limited to such specific
embodiments. Indeed, various modifications of the described modes for carrying out the invention
which are obvious to those skilled in molecular biology or related fields are intended to be within the
10 scope of the following claims.

Table 1

Polyptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	24	381039	HYPONOB01	381039H1 (HYPONOB01), 1558030F6 (BLADTUT04), 1653756H1 (PROSTUT08), 2327916X32C1 (COLNNOT11), 2327916X57C1 (COLNNOT11), 2503308H1 (CONUTUT01), 2649725F6 (KIDNFET01), 3105456F6 (HEAONOT05), 3278946H1 (STOMFET02)
2	25	383249	HYPONOB01	383249H1 (HYPONOB01), 4310090T6 (BRAUNOT01), SXBC01838V1, SXBC00070V1, SCSA04368V1, SXBC00956V1
3	26	618769	PGANNOT01	618769H1 (PGANNOT01), 618769R6 (PGANNOT01), 897423R1 (BRSTNOT05), 897423T1 (BRSTNOT05), 2225425T6 (SEMVNOT01), 3026516F6 (HEARFET02), 3031626H2 (TLYMNOT05)
4	27	1234837	LUNGFET03	763436R1 (BRAITUT02), 1234837H1 (LUNGFET03), 1237053F6 (LUNGFET03), 1721329F6 (BLADNOT06), 2751788R7 (THP1AZS08), g777187
5	28	1607223	LUNGNOT15	1250975F6 (LUNGFET03), 1270831F1 (TESTTUT02), 1399306F1 (BRAITUT08), 1607223H1 (LUNGNOT15), 1684804F6 (PROSNOT15), 1803848F6 (SINTNOT13), 2814466T6 (OVARNOT10), 3274147F6 (PROSBPT06), 3289542F6 (BONRFET01), SCFA04890V1
6	29	1621554	BRAITUT13	795816F1 (OVARNOT03), 795816R1 (OVARNOT03), 855789R1 (NGANNOT01), 1357564T6 (LUNGNOT09), 1621554H1 (BRAITUT13), 2081463H1 (UTRSNOT08), 2474413F6 (SMCANOT01), 2812794F6 (OVARNOT10), 3296847H1 (TLYJINT01), 4642794H1 (PROSTMT03)
7	30	1751553	LIVRTUT01	816693R6 (OVARNOT01), 816693X311D1 (OVARNOT01), 816693X313D1 (OVARNOT01), 1678292F6 (STOMFET01), 1678292T6 (STOMFET01), 1751553H1 (LIVRTUT01), 1981902R6 (LUNGNOT03), 1981902T6 (LUNGNOT03), 3050018H1 (LUNGNOT25), 4419520T6 (LIVRDIT02), 5208961H1 (BRAFNOT02), g2823700
8	31	1832403	BRAINON01	025882F1 (SPLNFET01), 1255756F2 (MENITUT03), 1832403H1 (BRAINON01), 2305321R6 (NGANNOT01)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
9	32	1971747	UCMCL5T01	285815F1 (EOSIHET02), 936208R1 (CERVNOT01), 1510143H1 (LUNGNOT14), 1673683F6 (BLADNOT05), 1912830F6 (LEUKNOT02), 1971747F6 (UCMCL5T01), 1971747H1 (UCMCL5T01), 2264823R6 (UTRSNOT02), 2858746F6 (SININOT03), 3108785F6 (BRSTTUT15), 3391023H1 (LUNGNOT17), 3736985F6 (SMCCNOS01), SBVA02967V1, SBVA04682V1, SBVA04527V1, g3882214
10	33	2285348	BRAINON01	228256F1 (PANCNOT01), 453676H1 (TLYMNOT02), 857831R1 (NGANNOT01), 1350071F1 (LATRTUT02), 1558538F1 (SPLNNOT04), 2170558F6 (ENDCNOT03), 2285348H1 (BRAINON01), 2291739T6 (BRAINON01), 2418450F6 (HNT3AZT01), 2717843F6 (THYRNOT09), 3404595F6 (ESOGNOT03), 3427632F6 (BRSTNOR01)
11	34	2374186	ISLTNOT01	1309077F1 (COLNFET02), 1526217F1 (UCMCL5T01), 2374186H1 (ISLTNOT01), 3581348F6 (293TF3T01), 3581348T6 (293TF3T01), SCGA06229V1, SCGA12945V1, SCGA02017V1, SCGA13028V1
12	35	2476232	SMCANOT01	1394147T1 (THYRNOT03), 1709170F6 (PROSNOT16), 2476232F6 (SMCANOT01), 2476232H1 (SMCANOT01), 2733026T6 (OVRTUT04), 3589738H1 (293TF5T01), 4638417T6 (MYEPTXT01), 5027954H1 (COLCDIT01)
13	36	2503986	CONUTUT01	449043X14 (TLYMNOT02), 632170R6 (KIDNNOT05), 1450259F1 (PENITUT01), 1798335F6 (COLNNOT27), 2503986H1 (CONUTUT01), 2872984H1 (THYRNOT10), 300424IT6 (TLYMNOT06), 3027284T6 (HEARFET02), SBFA03755F1, SBFA04352F1, SBFA04697F1, SBFA00555F1, SBFA01887F1
14	37	2596566	OVRTUT02	349546R6 (LVENNOT01), 840332R1 (PROSTUT05), 1333240F6 (COLNNOT13), 1628086F6 (COLNPOT01), 3293723F6 (TLYJINT01), 4008413H1 (ENDCNOT04), 4818893H1 (PROSTUT17), 2596566H2 (OVRTUT02)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
15	38	2685253	LUNGNOT23	680657R6 (UTRSNOT02), 736092R1 (TONSNOT01), 1486268T6 (CORPNOT02), 2445454T6 (THPINOT03), 2685253H1 (LUNGNOT23), 3731545H1 (SMCCNON03), 4020957H1 (BRAXNOT02), 4741584H1 (THYMNOR02), SZAT00201V1, SZAT01557V1, SZAT01529V1, SZAT00077V1, SZAT01181V1, SZAT01587V1, SZAT00683V1, SZAT00014V1, SZAT00017V1
16	39	2762252	BRSTNOT12	2685254H1 (LUNGNOT23), 2762252H1 (BRSTNOT12), 2766358T6 (BRSTNOT12), 2766358T6 (BRSTNOT12), 4364345H1 (SKIRNOT01), g3560562
17	40	3452009	UTRSNON03	3149588T6 (ADRENON04), 3452009H1 (UTRSNON03), SBHA03400F1, SBHA02544F1, SBHA03511F1
18	41	4644780	PROSTMT03	1494403R6 (PROSNON01), 1818377F6 (PROSNOT20), 3362284H1 (PROSBPT02), 3364760T6 (PROSBPT02), 4643280H1 (PROSTMT03), 4644736H1 (PROSTMT03), 4815885H1 (PROSTUS11), 5423826H1 (PROSTMT07), 5424290H1 (PROSTMT07)
19	42	4946103	SINTNOT25	1482075H1 (CORPNOT02), 1729337F6 (BRSTTUT08), 1729337X14C1 (BRSTTUT08), 1729337X16C1 (BRSTTUT08), 1730230X11C1 (BRSTTUT08), 1899544F6 (BLADTUT06), 2290603R6 (BRAINON01), 3042988H1 (HEAANOT01), 4946103H1 (SINTNOT25), SBAA00190F1, SBAA03996F1
20	43	5562355	BRSTDIT01	1909929F6 (CONNTUT01), 4027671H1 (BRAINOT23), 5091054F6 (UTRSTMR01), 5512655H1 (BRADDIR01), 5562355H1 (BRSTDIT01), g1625519
21	44	5678824	BRAENOT02	320465F1 (EOSIHET02), 506242X23R1 (TMLR3DT02), 1255854F2 (MENITUT03), 1351588F6 (LATRTUT02), 2051796F6 (LIVRFET02), 3421889H1 (UCMCNOT04), 5678824H1 (BRAENOT02), SCAA06386V1, SCAA02158V1, SBYA05287U1, SCAA04949V1

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
22	45	5870962	COLTDIT04	1415307F6 (BRAINOT12), 1504159F6 (BRAITUT07), 1692553T6 (COLNNOT23), 1868556T6 (SKINBIT01), 2911311H1 (KIDNTUT15), 2912463F6 (KIDNTUT15), 2926162T6 (TLYMNOT04), 5121545H1 (SMCBUNT01), 5812063H1 (KIDCTMT02), 5870962H1 (COLTDIT04), 6094909H1 (THP1TXT03)
23	46	2818605		2818605F6 (BRSTNOT14), 4904212T6 (TLYMNOT08), 1495645T6 (PROSNON01)

Table 2

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
1	481	S222 T56 T90 T92 S168 S313 S436 S217 S442	N433	Putative GTPase activating protein for ADP ribosylation factor (ArfGAP): C30-G153 HIV REV interacting protein: N44-P80, V83-N104, N43-W144 GATA-type zinc finger domain: N44-E93	g3135319 nucleoporin (Homo sapiens) Mikoshihara, K. et al. (1999) Chem. Phys. Lipids 98:59-67; Salcini, A.E. et al. (1997) Genes Dev. 11:2239-2249; Glockner, G. et al. (1998) Genome Res. 8: 1060-1073	Motifs BLAST-GENBANK BLIMPS_PRINTS BLAST_DOMO HMMER_PFBAM BLIMPS_PRODUM
2	195	S40 S110 S144		Podocalyxin-like protein: M1-T195 P24 protein: M1-T195	g1890141 Neural organelle transport protein P24 (Mus musculus) Kadota, Y. et al. (1997) Brain Res. Mol. Brain Res. 46:265-273	Motifs BLAST-GENBANK BLAST_PRODUM

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
3	313	T66 T76 S150 T212 S251 T256 S277 T80 S130 T155	N8	Protein kinase C C2 domain: L26-I115, L185-T273 Protein kinase C C2 domain: T13-Q69 C2 domain motif: L54-E79 C2 domain: G11-T128	g5926736 granuphilin-a (Mus musculus) Wang, J. et al. (1999) J. Biol. Chem. 274: 28542- 28548	Motifs BLAST-GENBANK HMMER_PFAM PROFILES SCAN BLIMPS_PFAM BLAST_DOMO
4	201	S11 T14 S82 T94 S119 T149		SNF7 Nuclear transcription regulation protein motif: N3-E166	g3873551 coiled- coil protein (Schizosaccharomyc es pombe)	Motifs BLAST-GENBANK BLAST_PRODOME
5	566	S112 S171 S195 S250 S282 S345 T381 T443 S503 T542 T547 S26 S35 S139 T377 S433 Y47	N213 N387	Gamma-adaptin Clathrin assembly protein complex: S407-W563 Gamma adaptin motif: M331-F541	g961444 related to mouse gamma adaptin (Homo sapiens) Nagase, T. et al. (1995) DNA Res. 2: 167-174	Motifs BLAST-GENBANK BLAST_PRODOME BLAST_DOMO
6	270	S192 S34 T91 S131 T146 T266 T18 T50 S187 S246			g4689260 sorting nexin 10 (Homo sapiens) Kurten, R. et al. (1996) Science 272:1008-1010	Motifs BLAST-GENBANK

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
7	490	S205 T320 T407 S94 S106 S156 S174 S218 S235 S255 S279 T320 S332 S397 S430 T18 S50 S102 S141 T162 S182 S200 S201 S208 S263 T304 T315 S327 S445	N22 N38 N180 N209 N234	C2 domain: I368-W456	g2724126 synaptotagmin VII (Homo sapiens) Cooper, P.R. et al. (1998) Genomics 49: 419-429	Motifs BLAST-GENBANK HMMER_PFM
8	136	T20 S50 T20 T117 Y81 Y129	N110		g4206090 snapin SNARE-associated synaptic transmission protein (Mus musculus) Ilardi, J.M. et al. (1999) Nat. Neurosci. 2:119- 124	Motifs BLAST-GENBANK

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
9	1104	T269 T816 T26 S27 T32 S4 S12 S123 S132 T182 S288 S342 S427 S453 S470 S506 T566 S626 T637 T752 T773 T826 S918 S933 S950 T959 S963 S101 T116 S123 S469 S686 S758 T769 T789 S831 S997 T1022 S1027 S1058	N212 N273 N1062	Transmembrane motif: Y69-R92 ATP-GTP binding site: G667-S674 C2 profile: L649-K735, I331- K417, L988-Q1077, L800- S877, L480-T558 C2 domain: V670-L682, E694- I707, L1014-E1039, L716- D724 G protein-coupled receptor: S881-L897 C2 domain: G973-D1097	g4193489 GLUT4 vesicle protein (Rattus norvegicus) Hashiramoto, M. (1998) Adv. Exp. Med. Biol. 441:47-61; Morris, N.J. et al. (1999) Biochim. Biophys. Acta 1431:525-530	Motifs BLAST-GENBANK HMMER HMMER_PFBM BLIMPS_PRINTS BLAST_DOMO BLIMPS_PFBM
10	411	S15 S19 S29 S49 T84 S135 T209 T322 S387 T406 S19 S75 T117 T128 T322	N27 N126 N237		g3560143 putative vacuolar protein sorting- associated protein (Schizosaccharomyc es pombe)	Motifs BLAST-GENBANK
11	201	S28 S141 S183 T44 Y171			g4689262 sorting nexin 11 (Homo sapiens)	Motifs BLAST-GENBANK

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
12	476	S61 S6 T44 S74 S156 S207 T245 T306 T330 T432 S6 T162 T245 S323	N43 N192 N305 N345 N385 N418		g3483017 TOM1-like protein (Homo sapiens) Seroussi, E. et al. (1999) Genomics 57:380-388	Motifs BLAST-GENBANK
13	1220	T1108 S81 T194 S233 S269 T434 S440 S465 T467 S527 S584 T674 S716 T1121 T1122 S1133 S66 S190 S211 T613 S799 S986 T1104 T1108 T1139 S1147 Y495 Y154 Y182	N40 N143 N562 N1181 N1197	WEI1 protein transport protein: M1-M229	g4104321 vesicle associated protein (Rattus norvegicus)	Motifs BLAST-GENBANK BLIMPS_PRODOR
14	222	T58 S93 T103 Y154 Y182	N197	Pleckstrin homology domain: A2-T109	g6013425 evectin-2 (Mus musculus) Krappa, R. et al. (1999) Proc. Natl. Acad. Sci. U.S.A. 96:4633-4638	Motifs BLAST-GENBANK HMMER_PFAM

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
15	924	S431 S14 S145 T178 T211 T330 T349 S387 S435 S525 T563 T574 T594 T651 T675 S691 S698 S751 S889 T25 T77 S89 T118 S128 T231 T239 S269 S426 T440 T554 S631 S674 S876 Y534	N164 N174 N177 N411 N475	ATP-GTP binding site: A309-T316 Exocyst complex component: N494-S918	g2827158 rsec5 exocyst complex subunit (Rattus norvegicus) Kee, Y. et al. (1997) Proc. Natl. Acad. Sci. U.S.A. 94:14438-14443	Motifs BLAST-GENBANK BLAST_PRODOM
16	435	T48 T63 S239 S241 T256 S261 S308 S313 S370 S390 T409 T426 T44 S86 T137 T164 S317 S325 T398 Y230	N254 N404 N407 N418		g3560561 PAM COOH-terminal interactor protein (Rattus norvegicus) Chen, L. et al. (1998) J. Biol. Chem. 273:33524-33532	Motifs BLAST-GENBANK

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
17	321	T72 S130 S168 S226 S253 S280 S104 T264	N41 N262	Vacuolar sorting- associated protein: L17-P152	g3319953 TOM1 (Homo sapiens) Seroussi, E. et al. (1999) Genomics 57:380-388	Motifs BLAST-GENBANK BLAST_PRODOM
18	499	S89 S140 T207 S209 S223 T259 S267 S456 S463 S38 S72 T264 S351 T481	N75 N461	Signal peptide: M1-T25 Lectin precursor: S38-P235 Lumenal domain: P9-D215	g606828 er-Golgi mannose- specific lectin (Homo sapiens) Arar, C. et al. (1995) J. Biol. Chem. 270:3551-3553	Motifs BLAST-GENBANK BLAST_PRODOM BLAST_DOMO HMMER
19	879	T14 S104 T150 S183 S236 S237 S244 S271 T436 S476 S484 S486 T494 S505 T533 S559 T708 S731 S834 T864 T44 S58 S210 S297 S382 S440 S449 T627 S649 S705 Y403 Y687 Y785 Y814	N131 N146 N482 N703		g2078441 similar to S. cerevisiae intracellular protein transport protein (Caenorhabditis elegans) Wilson, R. et al. (1994) Nature 368:32-38	Motifs BLAST-GENBANK

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
20	298	T36 S77 S84 T126 S201 S270	N57 N158	NSF attachment protein signature: R37-K56, A100-F117, E129-E146, N162- Y185, M193-C212, P233- E253, K264-K284 NSF attachment protein: K19-D294 Soluble attachment protein SNAP: M1-E292	g3929617 alpha SNAP (platelet SNARE- associated protein) (Homo sapiens) Lemons, P. et al. (1997) Blood 90:1490-1500	Motifs BLAST-GENBANK BLIMPS_PRINTS BLAST_PRODUM BLAST_DOMO
21	941	S293 S478 T941 T50 S101 T147 S217 T299 T325 T359 S468 S478 S579 T625 S707 S718 S827 S849 S851 S853 S923 S54 S217 S306 S467 S656 T800 S813 T918 S937 Y532 Y737	N435 N763 N892	ATP/GTP binding site: A193-T200 Zinc finger C3HC4: C822-C860, K815-C860 PHD finger: H835-S849 ATP-binding vacuolar biogenesis protein: L146-D466	g666102 vacuolar biogenesis protein END1 (PEP5) protein (Saccharomyces cerevisiae) Woolford, C. (1990) Genetics 125:739- 752	Motifs BLAST-GENBANK HMMER_PFBM BLIMPS_PFBM BLAST_PRODUM BLAST_DOMO

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphoryla- tion Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs and Domains	Homologous Sequence	Analytical Methods and Databases
22	336	T87 S130 T161 S43 T192 T267 T294 S301 T302 T318 T113 T125 T226 S227 T288	N117 N246	C2 domain: L47-E136, V196-R283, S34-V90 C2 domain signature: K63-L75, K92-K105, L137-D145 C2 Domain: L28-L137	g5926736 granuphilin-a (Mus musculus) Wang, J. et al. (1999) J. Biol. Chem. 274:28542- 28548	Motifs BLAST-GENBANK HMMER_PFAM BLIMPS_PRINTS BLAST_DOMO PROFILESCAN
23	163	T11 T54 S145 T155 S86	N125	Ankyrin repeats: K64-R96, E97-V129, F130-K162	g6624055 similar to ankyrin motif (Homo sapiens) Sulston, J.E. and Waterston, R. (1998) Genome Res. 8:1097-1108	MOTIFS HMMER_PFAM

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
24	1-292 686-1285 1715-1983	Gastrointestinal (0.250) Nervous (0.167) Cardiovascular (0.125) Reproductive (0.125)	Cancer (0.417) Inflammation (0.208) Cell proliferation (0.167)	PBLUESCRIPT
25	1302-1559	Nervous (1.000)	Cancer (0.250) Inflammation (0.250) Trauma (0.167)	PBLUESCRIPT
26	1-386 1034-1179	Reproductive (0.538) Musculoskeletal (0.154)	Cancer (0.615) Inflammation (0.154) Cell proliferation (0.077)	PSPORT1
27	1-1124	Reproductive (0.255) Cardiovascular (0.149) Nervous (0.128)	Cancer (0.404) Cell proliferation (0.298) Inflammation (0.234)	pINCY
28	1-839 1697-2291	Reproductive (0.276) Gastrointestinal (0.161) Nervous (0.149)	Cancer (0.540) Inflammation (0.207) Cell proliferation (0.138)	pINCY
29	1-1509	Reproductive (0.213) Nervous (0.170) Endocrine (0.128) Gastrointestinal (0.128)	Cancer (0.447) Cell proliferation (0.191) Inflammation (0.191)	pINCY
30	659-1352 1976-2599	Reproductive (0.485) Cardiovascular (0.121) Gastrointestinal (0.121)	Cancer (0.576) Inflammation (0.212) Trauma (0.121)	pINCY
31	1-415	Reproductive (0.273) Nervous (0.227) Gastrointestinal (0.159)	Cancer (0.500) Cell proliferation (0.182) Inflammation (0.182)	PSPORT1
32	1-578 1020-3418	Reproductive (0.254) Hematopoietic/Immune (0.176) Nervous (0.162)	Cancer (0.401) Inflammation (0.310) Cell proliferation (0.176)	PBLUESCRIPT

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
33	696-916 1980-2046 2794-3343	Reproductive (0.255) Cardiovascular (0.157) Nervous (0.137)	Cancer (0.510) Inflammation (0.196) Trauma (0.157)	PSPORT1
34	473-730 1404-1496	Nervous (0.278) Hematopoietic/Immune (0.208) Gastrointestinal (0.125)	Cancer (0.417) Inflammation (0.347) Cell proliferation (0.153)	pINCY
35	1-42 1247-1482	Reproductive (0.341) Gastrointestinal (0.205) Urologic (0.114)	Cancer (0.636) Cell proliferation (0.114) Inflammation (0.114)	pINCY
36	1-948 1687-1972 2448-2534 2800-2960 3035-3074	Reproductive (0.231) Nervous (0.166) Hematopoietic/Immune (0.126)	Cancer (0.467) Inflammation (0.226) Cell proliferation (0.196)	pINCY
37	1-38	Hematopoietic/Immune (0.222) Nervous (0.204) Reproductive (0.204)	Inflammation (0.370) Cancer (0.352) Cell proliferation (0.259)	pINCY
38	1-71 565-625 1142-2946 3445-3899	Reproductive (0.250) Nervous (0.235) Gastrointestinal (0.132)	Cancer (0.368) Inflammation (0.265) Cell proliferation (0.162)	pINCY
39	1-33 601-723 993-1319 1467-1551	Cardiovascular (0.250) Dermatologic (0.250) Reproductive (0.250) Urologic (0.250)	Cancer (0.250) Inflammation (0.250) Trauma (0.250)	pINCY
40	1-72 651-1014 1066-1088	Reproductive (0.267) Gastrointestinal (0.200) Hematopoietic/Immune (0.133) Nervous (0.133) Urologic (0.133)	Cancer (0.333) Inflammation (0.267) Trauma (0.200)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
41	1-1367	Reproductive (0.75)	Cancer (0.75) Trauma (0.167)	pINCY
42	1-569 1069-2802	Nervous (0.193) Reproductive (0.193) Gastrointestinal (0.169)	Cancer (0.422) Cell proliferation (0.253) Inflammation (0.229)	pINCY
43		Nervous (0.600) Musculoskeletal (0.300) Reproductive (0.100)	Cancer (0.600) Neurological (0.200) Trauma (0.100)	pINCY
44	1-600	Hematopoietic/Immune (0.192) Reproductive (0.192) Nervous (0.179)	Cancer (0.410) Inflammation (0.295) Cell proliferation (0.167)	pINCY
45	1-1363	Reproductive (0.298) Gastrointestinal (0.234) Nervous (0.149)	Cancer (0.532) Inflammation (0.234) Cell proliferation (0.149)	pINCY
46	1-448 388-985 529-1034	Breast T-lymphocyte Prostate	Adenocarcinoma Ductal Type	

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
24	HYPONOB01	This library was constructed using RNA (Clontech, #6579-2) isolated from the hypothalamus tissues of 51 male and female Caucasian donors, 16 to 75 years old.
25	HYPONOB01	This library was constructed using RNA (Clontech, #6579-2) isolated from the hypothalamus tissues of 51 male and female Caucasian donors, 16 to 75 years old.
26	PGANNOT01	This library was constructed using RNA isolated from paraganglionic tumor tissue removed from the intra-abdominal region of a 46-year-old Caucasian male. Pathology indicated a benign paraganglioma and was associated with renal cell carcinoma, clear cell type, which did not penetrate the capsule.
27	LUNGFEET03	This library was constructed using RNA isolated from lung tissue removed from a Caucasian female fetus, who died at 20 weeks' gestation.
28	LUNGNOT15	This library was constructed using RNA isolated from lung tissue removed from a 69-year-old Caucasian male. Pathology for the associated tumor tissue indicated residual invasive squamous cell carcinoma. Patient history included acute myocardial infarction, prostatic hyperplasia, and malignant skin neoplasm. Family history included cerebrovascular disease, type I diabetes, acute myocardial infarction, and arteriosclerotic coronary disease.
29	BRAITUT13	This library was constructed using RNA isolated from brain tumor tissue from the frontal lobe of a 68-year-old Caucasian male. Pathology indicated a meningioma in the frontal lobe.
30	LIVRTUT01	This library was constructed using RNA isolated from liver tumor tissue removed from a 51-year-old Caucasian female. Pathology indicated metastatic adenocarcinoma consistent with colon cancer. Family history included malignant neoplasm of the liver.
31	BRAINON01	This library was constructed and normalized from 4.88 million independent clones from a brain library. RNA was made from brain tissue from a 26-year-old Caucasian male. Pathology for the associated tumor tissue indicated oligoastrocytoma in the right fronto-parietal part of the brain.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
32	UCMCL5T01	This library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of 12 individuals. The cells were cultured for 12 days with IL-5 before RNA was obtained from the pooled lysates.
33	BRAINON01	This library was constructed and normalized from 4.88 million independent clones from a brain library. RNA was made from brain tissue from a 26-year-old Caucasian male. Pathology for the associated tumor tissue indicated oligoastrocytoma in the right fronto-parietal part of the brain.
34	ISLTNOT01	This library was constructed using RNA isolated from a pooled collection of pancreatic islet cells.
35	SMCANOT01	This library was constructed using RNA isolated from an aortic smooth muscle cell line derived from the explanted heart of a male during a heart transplant.
36	CONUTUT01	This library was constructed using RNA isolated from sigmoid mesentery tumor tissue obtained from a 61-year-old female during a total abdominal hysterectomy and bilateral salpingo-oophorectomy with regional lymph node excision. Pathology indicated a metastatic malignant mixed müllerian tumor present in the sigmoid mesentery at two sites.
37	OVARTUT02	This library was constructed using RNA isolated from ovarian tumor tissue removed from a 51-year-old Caucasian female during an exploratory laparotomy, total abdominal hysterectomy, salpingo-oophorectomy. Pathology indicated mucinous cystadenoma. Family history included atherosclerotic coronary artery disease, benign hypertension, breast cancer, and uterine cancer.
38	LUNGNOT23	This library was constructed using RNA isolated from lung tissue from a 58-year-old Caucasian male. Pathology for the associated tumor tissue indicated metastatic grade 3 (of 4) osteosarcoma. Patient history included soft tissue cancer, secondary cancer of the lung, prostate cancer, and an acute duodenal ulcer with hemorrhage. Family history included prostate cancer, breast cancer, and acute leukemia.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
39	BRSTNOT12	This library was constructed using RNA isolated from diseased breast tissue removed from a 32-year-old Caucasian female during a bilateral reduction mammoplasty. Pathology indicated nonproliferative fibrocystic disease. Family history included benign hypertension and atherosclerotic coronary artery disease.
40	UTRSNON03	This normalized library was constructed from 6.4 million independent clones from a uterus library. RNA was isolated from uterine myometrial tissue removed from a 41-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. Pathology for the associated tumor tissue indicated uterine leiomyoma. Patient history included ventral hernia and a benign ovarian neoplasm. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9228).
41	PROSTMT03	This library was constructed using RNA isolated from prostate tissue removed from a 68-year-old Caucasian male during a radical prostatectomy and regional lymph node excision. Pathology for the associated tumor indicated adenocarcinoma. The patient presented with elevated prostate specific antigen (PSA) and induration. Patient history included pure hypercholesterolemia, kidney calculus, an unspecified allergy, and atopic dermatitis. Family history included colon cancer.
42	SINTNOT25	This library was constructed using RNA isolated from small intestine tissue removed from a 13-year-old Caucasian male, who died from a gunshot wound to the head. Family history included diabetes.
43	BRSTDIT01	This library was constructed using RNA isolated from diseased breast tissue from a 48-year-old Caucasian female. Pathology for the associated tumor tissue indicated intraductal cancer. The patient presented with a malignant neoplasm of the breast and unspecified breast symptoms. Patient history included mitral valve disorder and an unspecified disease of the shoulder region. Family history included malignant neoplasm of the breast and hyperlipidemia, malignant neoplasm of the colon and cardiac dysrhythmias, and malignant neoplasm of the colon.
44	BRAENOT02	This library was constructed using RNA isolated from posterior parietal cortex tissue removed from the brain of a 35-year-old Caucasian male.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
45	COLTDIT04	This library was constructed from diseased transverse colon tissue removed from a 16-year-old Caucasian male during partial colectomy, temporary ileostomy, and colonoscopy. Pathology indicated innumerable (greater than 100) adenomatous polyps with low-grade dysplasia involving the entire colonic mucosa in the setting of familial polyposis coli. The anal mucosa showed 10 adenomatous polyps with low-grade dysplasia in the setting of familial polyposis coli. The patient presented with abdominal pain and flatulence. Family history included benign colon neoplasm in the father; benign colon neoplasm in the sibling(s); and benign hypertension, cerebrovascular disease, breast cancer, uterine cancer, and type II diabetes in the grandparent(s).

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score > GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-23,
 b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-23,
 c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23, and
10 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-23.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-23.

15 3. An isolated polynucleotide encoding a polypeptide of claim 1.

 4. An isolated polynucleotide encoding a polypeptide of claim 2.

20 5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:24-46.

 6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

25 7. A cell transformed with a recombinant polynucleotide of claim 6.

 8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

30 9. A method for producing a polypeptide of claim 1, the method comprising:
 a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

5 11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46,

b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:24-46,

10 c) a polynucleotide sequence complementary to a),

d) a polynucleotide sequence complementary to b), and

e) an RNA equivalent of a)-d).

15 12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

20 a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

25 14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

30 a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide comprises an amino acid sequence
5 selected from the group consisting of SEQ ID NO:1-23.

18. A method for treating a disease or condition associated with decreased expression of functional VETRP, comprising administering to a patient in need of such treatment the composition of claim 16.

10 19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

15 20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

20 21. A method for treating a disease or condition associated with decreased expression of functional VETRP, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- 25
- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
 - b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

30 24. A method for treating a disease or condition associated with overexpression of functional VETRP, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim

1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
 - b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying
- 5 a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under conditions
- 10 permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
 - c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change
- 15 in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method

20 comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
 - b) detecting altered expression of the target polynucleotide, and
 - c) comparing the expression of the target polynucleotide in the presence of varying amounts
- 25 of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
 - b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at
- 30 least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- c) quantifying the amount of hybridization complex; and

d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

<110> INCYTE GENOMICS, INC.

TANG, Y. Tom
YUE, Henry
BANDMAN, Olga
HILLMAN, Jennifer
BAUGHN, Mariah R.
LU, Dyung Aina M.
AZIMZAI, Yalda

<120> VESICLE TRAFFICKING PROTEINS

<130> PF-0756 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/172,968; 60/172,066

<151> 1999-12-21; 1999-12-23

<160> 46

<170> PERL Program

<210> 1

<211> 481

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 381039CD1

<400> 1

Met	Val	Met	Ala	Ala	Lys	Lys	Gly	Pro	Gly	Pro	Gly	Gly	Gly	Val
1				5					10					15
Ser	Gly	Gly	Lys	Ala	Glu	Ala	Glu	Ala	Ala	Ser	Glu	Val	Trp	Cys
				20					25					30
Arg	Arg	Val	Arg	Glu	Leu	Gly	Gly	Cys	Ser	Gln	Ala	Gly	Asn	Arg
				35					40					45
His	Cys	Phe	Glu	Cys	Ala	Gln	Arg	Gly	Val	Thr	Tyr	Val	Asp	Ile
				50					55					60
Thr	Val	Gly	Ser	Phe	Val	Cys	Thr	Thr	Cys	Ser	Gly	Leu	Leu	Arg
				65					70					75
Gly	Leu	Asn	Pro	Pro	His	Arg	Val	Lys	Ser	Ile	Ser	Met	Thr	Thr
				80					85					90
Phe	Thr	Glu	Pro	Glu	Val	Val	Phe	Leu	Gln	Ser	Arg	Gly	Asn	Glu
				95					100					105
Val	Cys	Arg	Lys	Ile	Trp	Leu	Gly	Leu	Phe	Asp	Ala	Arg	Thr	Ser
				110					115					120
Leu	Val	Pro	Asp	Ser	Arg	Asp	Pro	Gln	Lys	Val	Lys	Glu	Phe	Leu
				125					130					135
Gln	Glu	Lys	Tyr	Glu	Lys	Lys	Arg	Trp	Tyr	Val	Pro	Pro	Asp	Gln
				140					145					150
Val	Lys	Gly	Pro	Thr	Tyr	Thr	Lys	Gly	Ser	Ala	Ser	Thr	Pro	Val
				155					160					165
Gln	Gly	Ser	Ile	Pro	Glu	Gly	Lys	Pro	Leu	Arg	Thr	Leu	Leu	Gly
				170					175					180
Asp	Pro	Ala	Pro	Ser	Leu	Ser	Val	Ala	Ala	Ser	Thr	Ser	Ser	Gln
				185					190					195
Pro	Val	Ser	Gln	Ser	His	Ala	Arg	Thr	Ser	Gln	Ala	Arg	Ser	Thr
				200					205					210
Gln	Pro	Pro	Pro	His	Ser	Ser	Val	Lys	Lys	Ala	Ser	Thr	Asp	Leu
				215					220					225
Leu	Ala	Asp	Ile	Gly	Gly	Asp	Pro	Phe	Ala	Ala	Pro	Gln	Met	Ala
				230					235					240
Pro	Ala	Phe	Ala	Ala	Phe	Pro	Ala	Phe	Gly	Gly	Gln	Thr	Pro	Ser
				245					250					255
Gln	Gly	Gly	Phe	Ala	Asn	Phe	Asp	Ala	Phe	Ser	Ser	Gly	Pro	Ser

	260		265		270
Ser Ser Val Phe	Gly	Ser Leu Pro Pro	Ala Gly Gln Ala Ser	Phe	
	275		280		285
Gln Ala Gln Pro	Thr	Pro Ala Gly Ser	Ser Gln Gly Thr Pro	Phe	
	290		295		300
Gly Ala Thr Pro	Leu	Ala Pro Ala Ser	Gln Pro Asn Ser Leu	Ala	
	305		310		315
Asp Val Gly Ser	Phe	Leu Gly Pro Gly	Val Pro Ala Ala Gly	Val	
	320		325		330
Pro Ser Ser Leu	Phe	Gly Met Ala Gly	Gln Val Pro Pro Leu	Gln	
	335		340		345
Ser Val Thr Met	Gly	Gly Gly Gly Gly	Ser Ser Thr Gly Leu	Ala	
	350		355		360
Phe Gly Ala Phe	Thr	Asn Pro Phe Thr	Ala Pro Ala Ala Gln	Ser	
	365		370		375
Pro Leu Pro Ser	Thr	Asn Pro Phe Gln	Pro Asn Gly Leu Ala	Pro	
	380		385		390
Gly Pro Gly Phe	Gly	Met Ser Ser Ala	Gly Pro Gly Phe Pro	Gln	
	395		400		405
Ala Val Pro Pro	Thr	Gly Ala Phe Ala	Ser Ser Phe Pro Ala	Pro	
	410		415		420
Leu Phe Pro Pro	Gln	Thr Pro Leu Val	Gln Gln Gln Asn Gly	Ser	
	425		430		435
Ser Phe Gly Asp	Leu	Gly Ser Ala Lys	Leu Gly Gln Arg Pro	Leu	
	440		445		450
Ser Gln Pro Ala	Gly	Ile Ser Thr Asn	Pro Phe Met Thr Gly	Pro	
	455		460		465
Ser Ser Ser Pro	Phe	Ala Ser Lys Pro	Pro Thr Thr Asn Pro	Phe	
	470		475		480

Leu

<210> 2
 <211> 195
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 383249CD1

<400> 2

Met Ser Ser Cys Ser	Asn Val Cys Gly Ser	Arg Gln Ala Gln Ala	
1	5	10	15
Ala Ala Glu Gly Gly	Tyr Gln Arg Tyr Gly	Val Arg Ser Tyr Leu	
	20	25	30
His Gln Phe Tyr Glu	Asp Cys Thr Ala Ser	Ile Trp Glu Tyr Glu	
	35	40	45
Asp Asp Phe Gln Ile	Gln Arg Ser Pro Asn	Arg Trp Ser Ser Val	
	50	55	60
Phe Trp Lys Val Gly	Leu Ile Ser Gly Thr	Val Phe Val Ile Leu	
	65	70	75
Gly Leu Thr Val Leu	Ala Val Gly Phe Leu	Val Pro Pro Lys Ile	
	80	85	90
Glu Ala Phe Gly Glu	Ala Asp Phe Val Val	Val Asp Thr His Ala	
	95	100	105
Val Gln Phe Asn Ser	Ala Leu Asp Met Tyr	Lys Leu Ala Gly Ala	
	110	115	120
Val Leu Phe Cys Ile	Gly Gly Thr Ser Met	Ala Gly Cys Leu Leu	
	125	130	135
Met Ser Val Phe Val	Lys Ser Tyr Ser Lys	Glu Glu Lys Phe Leu	
	140	145	150
Gln Gln Lys Phe Lys	Glu Arg Ile Ala Asp	Ile Lys Ala His Thr	
	155	160	165
Gln Pro Val Thr Lys	Ala Pro Gly Pro Gly	Glu Thr Lys Ile Pro	
	170	175	180
Val Thr Leu Ser Arg	Val Gln Asn Val Gln	Pro Leu Leu Ala Thr	

185

190

195

<210> 3
 <211> 313
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 618769CD1

<400> 3
 Met Gly Asn Phe Asp Asn Ala Asn Val Thr Gly Glu Ile Glu Phe
 1 5 10 15
 Ala Ile His Tyr Cys Phe Lys Thr His Ser Leu Glu Ile Cys Ile
 20 25 30
 Lys Ala Cys Lys Asn Leu Ala Tyr Gly Glu Glu Lys Lys Lys Lys
 35 40 45
 Cys Asn Pro Tyr Val Lys Thr Tyr Leu Leu Pro Asp Arg Ser Ser
 50 55 60
 Gln Gly Lys Arg Lys Thr Gly Val Gln Arg Asn Thr Val Asp Pro
 65 70 75
 Thr Phe Gln Glu Thr Leu Lys Tyr Gln Val Ala Pro Ala Gln Leu
 80 85 90
 Val Thr Arg Gln Leu Gln Val Ser Val Trp His Leu Gly Thr Leu
 95 100 105
 Ala Arg Arg Val Phe Leu Gly Glu Val Ile Ile Pro Leu Ala Thr
 110 115 120
 Trp Asp Phe Glu Asp Ser Thr Thr Gln Ser Phe Arg Trp His Pro
 125 130 135
 Leu Arg Ala Lys Ala Glu Lys Tyr Glu Asp Ser Val Pro Gln Ser
 140 145 150
 Asn Gly Glu Leu Thr Val Arg Ala Lys Leu Val Leu Pro Ser Arg
 155 160 165
 Pro Arg Lys Leu Gln Glu Ala Gln Glu Gly Thr Asp Gln Pro Ser
 170 175 180
 Leu His Gly Gln Leu Cys Leu Val Val Leu Gly Ala Lys Asn Leu
 185 190 195
 Pro Val Arg Pro Asp Gly Thr Leu Asn Ser Phe Val Lys Gly Cys
 200 205 210
 Leu Thr Leu Pro Asp Gln Gln Lys Leu Arg Leu Lys Ser Pro Val
 215 220 225
 Leu Arg Lys Gln Ala Cys Pro Gln Trp Lys His Ser Phe Val Phe
 230 235 240
 Ser Gly Val Thr Pro Ala Gln Leu Arg Gln Ser Ser Leu Glu Leu
 245 250 255
 Thr Val Trp Asp Gln Ala Leu Phe Gly Met Asn Asp Arg Leu Leu
 260 265 270
 Gly Gly Thr Arg Leu Gly Ser Lys Gly Asp Thr Ala Val Gly Gly
 275 280 285
 Asp Ala Cys Ser Leu Ser Lys Leu Gln Trp Gln Lys Val Leu Ser
 290 295 300
 Ser Pro Asn Leu Trp Thr Asp Met Thr Leu Val Leu His
 305 310

<210> 4
 <211> 201
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1234837CD1

<400> 4
 Met Gly Asn Leu Phe Gly Arg Lys Lys Gln Ser Arg Val Thr Glu

1	5	10	15
Gln Asp Lys Ala Ile	Leu Gln Leu Lys	Gln Gln Arg Asp Lys	Leu
20	25	30	
Arg Gln Tyr Gln Lys	Arg Ile Ala Gln	Gln Leu Glu Arg Glu	Arg
35	40	45	
Ala Leu Ala Arg Gln	Leu Leu Arg Asp	Gly Arg Lys Glu Arg	Ala
50	55	60	
Lys Leu Leu Leu Lys	Lys Lys Arg Tyr	Gln Glu Gln Leu Leu	Asp
65	70	75	
Arg Thr Glu Asn Gln	Ile Ser Ser Leu	Glu Ala Met Val Gln	Ser
80	85	90	
Ile Glu Phe Thr Gln	Ile Glu Met Lys	Val Met Glu Gly Leu	Gln
95	100	105	
Phe Gly Asn Glu Cys	Leu Asn Lys Met	His Gln Val Met Ser	Ile
110	115	120	
Glu Glu Val Glu Arg	Ile Leu Asp Glu	Thr Gln Glu Ala Val	Glu
125	130	135	
Tyr Gln Arg Gln Ile	Asp Glu Leu Leu	Ala Gly Ser Phe Thr	Gln
140	145	150	
Glu Asp Glu Asp Ala	Ile Leu Glu Glu	Leu Ser Ala Ile Thr	Gln
155	160	165	
Glu Gln Ile Glu Leu	Pro Glu Val Pro	Ser Glu Pro Leu Pro	Glu
170	175	180	
Lys Ile Pro Glu Asn	Val Pro Val Lys	Ala Arg Pro Arg Gln	Ala
185	190	195	
Glu Leu Val Ala Ala	Ser		
200			

<210> 5

<211> 566

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1607223CD1

<400> 5

Met Lys Ser Cys Gly	Lys Arg Phe His	Asp Glu Val Gly Lys	Phe
1	5	10	15
Arg Phe Leu Asn Glu	Leu Ile Lys Val	Val Ser Pro Lys Tyr	Leu
20	25	30	
Gly Ser Arg Thr Ser	Glu Lys Val Lys	Asn Lys Ile Leu Glu	Leu
35	40	45	
Leu Tyr Ser Trp Thr	Val Gly Leu Pro	Glu Glu Val Lys Ile	Ala
50	55	60	
Glu Ala Tyr Gln Met	Leu Lys Lys Gln	Gly Ile Val Lys Ser	Asp
65	70	75	
Pro Lys Leu Pro Asp	Asp Thr Thr Phe	Pro Leu Pro Pro Pro	Arg
80	85	90	
Pro Lys Asn Val Ile	Phe Glu Asp Glu	Glu Lys Ser Lys Met	Leu
95	100	105	
Ala Arg Leu Leu Lys	Ser Ser His Pro	Glu Asp Leu Arg Ala	Ala
110	115	120	
Asn Lys Leu Ile Lys	Glu Met Val Gln	Glu Asp Gln Lys Arg	Met
125	130	135	
Glu Lys Ile Ser Lys	Arg Val Asn Ala	Ile Glu Glu Val Asn	Asn
140	145	150	
Asn Val Lys Leu Leu	Thr Glu Met Val	Met Ser His Ser Gln	Gly
155	160	165	
Gly Ala Ala Ala Gly	Ser Ser Glu Asp	Leu Met Lys Glu Leu	Tyr
170	175	180	
Gln Arg Cys Glu Arg	Met Arg Pro Thr	Leu Phe Arg Leu Ala	Ser
185	190	195	
Asp Thr Glu Asp Asn	Asp Glu Ala Leu	Ala Glu Ile Leu Gln	Ala
200	205	210	
Asn Asp Asn Leu Thr	Gln Val Ile Asn	Leu Tyr Lys Gln Leu	Val

Arg Gly Glu Glu	215	Val Asn Gly Asp Ala	220	Thr Ala Gly Ser Ile	225
	230		235		240
Gly Ser Thr Ser	245	Ala Leu Leu Asp Leu	250	Ser Gly Leu Asp Leu	255
	260		265		270
Pro Ala Gly Thr	275	Thr Tyr Pro Ala Met	280	Pro Thr Arg Pro Gly	285
	290		295		300
Gln Ala Ser Pro	305	Glu Gln Pro Ser Ala	310	Ser Val Ser Leu Leu	315
	320		325		330
Asp Glu Leu Met	335	Ser Leu Gly Leu Ser	340	Asp Pro Thr Pro Pro	345
	350		355		360
Gly Pro Ser Leu	365	Asp Gly Thr Gly Trp	370	Ser Phe Gln Ser Ser	375
	380		385		390
Asp Ala Thr Glu	395	Pro Pro Ala Pro Ala	400	Leu Ala Gln Ala Pro	405
	410		415		420
Met Glu Ser Arg	425	Pro Pro Ala Gln Thr	430	Ser Leu Pro Ala Ser	435
	440		445		450
Gly Leu Asp Asp	455	Leu Asp Leu Leu Gly	460	Lys Thr Leu Leu Gln	465
	470		475		480
Ser Leu Pro Pro	485	Glu Ser Gln Gln Val	490	Arg Trp Glu Lys Gln	495
	500		505		510
Pro Thr Pro Arg	515	Leu Thr Leu Arg Asp	520	Gln Asn Lys Ser Ser	525
	530		535		540
Ser Cys Ser Ser	545	Pro Ser Ser Ser Ala	550	Thr Ser Leu Leu His	555
	560		565		
Val Ser Pro Glu		Pro Pro Arg Pro Pro		Gln Gln Pro Val Pro	
Glu Leu Ser Leu		Ala Ser Ile Thr Val		Pro Leu Glu Ser Ile	
Pro Ser Asn Ile		Leu Pro Val Thr Val		Tyr Asp Gln His Gly	
Arg Ile Leu Phe		His Phe Ala Arg Asp		Pro Leu Pro Gly Arg	
Asp Val Leu Val		Val Val Val Ser Met		Leu Ser Thr Ala Pro	
Pro Ile Arg Asn		Ile Val Phe Gln Ser		Ala Val Pro Lys Val	
Lys Val Lys Leu		Gln Pro Pro Ser Gly		Thr Glu Leu Pro Ala	
Asn Pro Ile Val		His Pro Ser Ala Ile		Thr Gln Val Leu Leu	
Ala Asn Pro Gln		Lys Glu Lys Val Arg		Leu Arg Tyr Lys Leu	
Phe Thr Met Gly		Asp Gln Thr Tyr Asn		Glu Met Gly Asp Val	
Gln Phe Pro Pro		Glu Thr Trp Gly		Ser Leu	

<210> 6

<211> 270

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1621554CD1

<400> 6

Met Gly Phe Trp Cys	Arg Met Ser Glu Asn	Gln Glu Gln Glu Glu
1	5	10
Val Ile Thr Val Arg	Val Gln Asp Pro Arg	Val Gln Asn Glu Gly
	20	25
Ser Trp Asn Ser Tyr	Val Asp Tyr Lys Ile	Phe Leu His Thr Asn
	35	40
Ser Lys Ala Phe Thr	Ala Lys Thr Ser Cys	Val Arg Arg Arg Tyr
	50	55
Arg Glu Phe Val Trp	Leu Arg Lys Gln Leu	Gln Arg Asn Ala Gly

65	70	75
Leu Val Pro Val Pro	Glu Leu Pro Gly Lys	Ser Thr Phe Phe Gly
80	85	90
Thr Ser Asp Glu Phe	Ile Glu Lys Arg Arg	Gln Gly Leu Gln His
95	100	105
Phe Leu Glu Lys Val	Leu Gln Ser Val Val	Leu Leu Ser Asp Ser
110	115	120
Gln Leu His Leu Phe	Leu Gln Ser Gln Leu	Ser Val Pro Glu Ile
125	130	135
Glu Ala Cys Val Gln	Gly Arg Ser Thr Met	Thr Val Ser Asp Ala
140	145	150
Ile Leu Arg Tyr Ala	Met Ser Asn Cys Gly	Trp Ala Gln Glu Glu
155	160	165
Arg Gln Ser Ser Ser	His Leu Ala Lys Gly	Asp Gln Pro Lys Ser
170	175	180
Cys Cys Phe Leu Pro	Arg Ser Gly Arg Arg	Ser Ser Pro Ser Pro
185	190	195
Pro Pro Ser Glu Glu	Lys Asp His Leu Glu	Val Trp Ala Pro Val
200	205	210
Val Asp Ser Glu Val	Pro Ser Leu Glu Ser	Pro Thr Leu Pro Pro
215	220	225
Leu Ser Ser Pro Leu	Cys Cys Asp Phe Gly	Arg Pro Lys Glu Gly
230	235	240
Thr Ser Thr Leu Gln	Ser Val Arg Arg Ala	Val Gly Gly Asp His
245	250	255
Ala Val Pro Leu Asp	Pro Gly Gln Leu Glu	Thr Val Leu Glu Lys
260	265	270

<210> 7
 <211> 490
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1751553CD1

<400> 7
Met Ala Thr Glu Phe Ile Lys Ser Cys Cys Gly Gly Cys Phe Tyr
1 5 10 15
Gly Glu Thr Glu Lys His Asn Phe Ser Val Glu Arg Asp Phe Lys
20 25 30
Ala Ala Val Pro Asn Ser Gln Asn Ala Thr Ile Ser Val Pro Pro
35 40 45
Leu Thr Ser Val Ser Val Lys Pro Gln Leu Gly Cys Thr Glu Asp
50 55 60
Tyr Leu Leu Ser Lys Leu Pro Ser Asp Gly Lys Glu Val Pro Phe
65 70 75
Val Val Pro Lys Phe Lys Leu Ser Tyr Ile Gln Pro Arg Thr Gln
80 85 90
Glu Thr Pro Ser His Leu Glu Glu Leu Gly Ser Ala Arg Ala
95 100 105
Ser Phe Gly Asp Arg Lys Val Glu Leu Ser Ser Ser Ser Gln His
110 115 120
Gly Pro Ser Tyr Asp Val Tyr Asn Pro Phe Tyr Met Tyr Gln His
125 130 135
Ile Ser Pro Asp Leu Ser Arg Arg Phe Pro Pro Arg Ser Glu Val
140 145 150
Thr Arg Leu Tyr Gly Ser Val Cys Asp Leu Arg Thr Asn Lys Leu
155 160 165
Pro Gly Ser Pro Gly Leu Ser Lys Ser Met Phe Asp Leu Thr Asn
170 175 180
Ser Ser Gln Arg Phe Ile Gln Arg His Asp Ser Leu Ser Ser Val
185 190 195
Pro Ser Ser Ser Ser Ser Arg Lys Asn Ser Gln Gly Ser Asn Arg
200 205 210

Ser	Leu	Asp	Thr	Ile	Thr	Leu	Ser	Gly	Asp	Glu	Arg	Asp	Phe	Gly	215	220	225
Arg	Leu	Asn	Val	Lys	Leu	Phe	Tyr	Asn	Ser	Ser	Val	Glu	Gln	Ile	230	235	240
Trp	Ile	Thr	Val	Leu	Gln	Cys	Arg	Asp	Leu	Ser	Trp	Pro	Ser	Ser	245	250	255
Tyr	Gly	Asp	Thr	Pro	Thr	Val	Ser	Ile	Lys	Gly	Ile	Leu	Thr	Leu	260	265	270
Pro	Lys	Pro	Val	His	Phe	Lys	Ser	Ser	Ala	Lys	Glu	Gly	Ser	Asn	275	280	285
Ala	Ile	Glu	Phe	Met	Glu	Thr	Phe	Val	Phe	Ala	Ile	Lys	Leu	Gln	290	295	300
Asn	Leu	Gln	Thr	Val	Arg	Leu	Val	Phe	Lys	Ile	Gln	Thr	Gln	Thr	305	310	315
Pro	Arg	Lys	Lys	Thr	Ile	Gly	Glu	Cys	Ser	Met	Ser	Leu	Arg	Thr	320	325	330
Leu	Ser	Thr	Gln	Glu	Met	Asp	Tyr	Ser	Leu	Asp	Ile	Thr	Pro	Pro	335	340	345
Ser	Lys	Ile	Ser	Val	Cys	His	Ala	Glu	Leu	Glu	Leu	Gly	Thr	Cys	350	355	360
Phe	Gln	Ala	Val	Asn	Ser	Arg	Ile	Gln	Leu	Gln	Ile	Leu	Glu	Ala	365	370	375
Arg	Tyr	Leu	Pro	Ser	Ser	Ser	Thr	Pro	Leu	Thr	Leu	Ser	Phe	Phe	380	385	390
Val	Lys	Val	Gly	Met	Phe	Ser	Ser	Gly	Glu	Leu	Ile	Tyr	Lys	Lys	395	400	405
Lys	Thr	Arg	Leu	Leu	Lys	Ala	Ser	Asn	Gly	Arg	Val	Lys	Trp	Gly	410	415	420
Glu	Thr	Met	Ile	Phe	Pro	Leu	Ile	Gln	Ser	Glu	Lys	Glu	Ile	Val	425	430	435
Phe	Leu	Ile	Lys	Leu	Tyr	Ser	Arg	Ser	Ser	Val	Arg	Arg	Lys	His	440	445	450
Phe	Val	Gly	Gln	Ile	Trp	Ile	Ser	Glu	Asp	Ser	Asn	Asn	Ile	Glu	455	460	465
Ala	Val	Asn	Gln	Trp	Lys	Glu	Thr	Val	Ile	Asn	Pro	Glu	Lys	Val	470	475	480
Val	Ile	Arg	Trp	His	Lys	Leu	Asn	Pro	Ser						485	490	

<210> 8

<211> 136

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1832403CD1

<400> 8

Met	Ala	Gly	Ala	Gly	Ser	Ala	Ala	Val	Ser	Gly	Ala	Gly	Thr	Pro	1	5	10	15
Val	Ala	Gly	Pro	Thr	Gly	Arg	Asp	Leu	Phe	Ala	Glu	Gly	Leu	Leu	20	25	30	35
Glu	Phe	Leu	Arg	Pro	Ala	Val	Gln	Gln	Leu	Asp	Ser	His	Val	His	40	45	50	55
Ala	Val	Arg	Glu	Ser	Gln	Val	Glu	Leu	Arg	Glu	Gln	Ile	Asp	Asn	60	65	70	75
Leu	Ala	Thr	Glu	Leu	Cys	Arg	Ile	Asn	Glu	Asp	Gln	Lys	Val	Ala	80	85	90	95
Leu	Asp	Leu	Asp	Pro	Tyr	Val	Lys	Lys	Leu	Leu	Asn	Ala	Arg	Arg	100	105	110	115
Arg	Val	Val	Leu	Val	Asn	Asn	Ile	Leu	Gln	Asn	Ala	Gln	Glu	Arg	120	125	130	135
Leu	Arg	Arg	Leu	Asn	His	Ser	Val	Ala	Lys	Glu	Thr	Ala	Arg	Arg				
Arg	Ala	Met	Leu	Asp	Ser	Gly	Ile	Tyr	Pro	Pro	Gly	Ser	Pro	Gly				

Lys

<210> 9

<211> 1104

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1971747CD1

<400> 9

Met	Glu	Arg	Ser	Pro	Gly	Glu	Gly	Pro	Ser	Pro	Ser	Pro	Met	Asp
1				5					10					15
Gln	Pro	Ser	Ala	Pro	Ser	Asp	Pro	Thr	Asp	Gln	Pro	Pro	Ala	Ala
				20					25					30
His	Ala	Lys	Pro	Asp	Pro	Gly	Ser	Gly	Gly	Gln	Pro	Ala	Gly	Pro
				35					40					45
Gly	Ala	Ala	Gly	Glu	Ala	Leu	Ala	Val	Leu	Thr	Ser	Phe	Gly	Arg
				50					55					60
Arg	Leu	Leu	Val	Leu	Ile	Pro	Val	Tyr	Leu	Ala	Gly	Ala	Val	Gly
				65					70					75
Leu	Ser	Val	Gly	Phe	Val	Leu	Phe	Gly	Leu	Ala	Leu	Tyr	Leu	Gly
				80					85					90
Trp	Arg	Arg	Val	Arg	Asp	Glu	Lys	Glu	Arg	Ser	Leu	Arg	Ala	Ala
				95					100					105
Arg	Gln	Leu	Leu	Asp	Asp	Glu	Glu	Gln	Leu	Thr	Ala	Lys	Thr	Leu
				110					115					120
Tyr	Met	Ser	His	Arg	Glu	Leu	Pro	Ala	Trp	Val	Ser	Phe	Pro	Asp
				125					130					135
Val	Glu	Lys	Ala	Glu	Trp	Leu	Asn	Lys	Ile	Val	Ala	Gln	Val	Trp
				140					145					150
Pro	Phe	Leu	Gly	Gln	Tyr	Met	Glu	Lys	Leu	Leu	Ala	Glu	Thr	Val
				155					160					165
Ala	Pro	Ala	Val	Arg	Gly	Ser	Asn	Pro	His	Leu	Gln	Thr	Phe	Thr
				170					175					180
Phe	Thr	Arg	Val	Glu	Leu	Gly	Glu	Lys	Pro	Leu	Arg	Ile	Ile	Gly
				185					190					195
Val	Lys	Val	His	Pro	Gly	Gln	Arg	Lys	Glu	Gln	Ile	Leu	Leu	Asp
				200					205					210
Leu	Asn	Ile	Ser	Tyr	Val	Gly	Asp	Val	Gln	Ile	Asp	Val	Glu	Val
				215					220					225
Lys	Lys	Tyr	Phe	Cys	Lys	Ala	Gly	Val	Lys	Gly	Met	Gln	Leu	His
				230					235					240
Gly	Val	Leu	Arg	Val	Ile	Leu	Glu	Pro	Leu	Ile	Gly	Asp	Leu	Pro
				245					250					255
Phe	Val	Gly	Ala	Val	Ser	Met	Phe	Phe	Ile	Arg	Arg	Pro	Thr	Leu
				260					265					270
Asp	Ile	Asn	Trp	Thr	Gly	Met	Thr	Asn	Leu	Leu	Asp	Ile	Pro	Gly
				275					280					285
Leu	Ser	Ser	Leu	Ser	Asp	Thr	Met	Ile	Met	Asp	Ser	Ile	Ala	Ala
				290					295					300
Phe	Leu	Val	Leu	Pro	Asn	Arg	Leu	Leu	Val	Pro	Leu	Val	Pro	Asp
				305					310					315
Leu	Gln	Asp	Val	Ala	Gln	Leu	Arg	Ser	Pro	Leu	Pro	Arg	Gly	Ile
				320					325					330
Ile	Arg	Ile	His	Leu	Leu	Ala	Ala	Arg	Gly	Leu	Ser	Ser	Lys	Asp
				335					340					345
Lys	Tyr	Val	Lys	Gly	Leu	Ile	Glu	Gly	Lys	Ser	Asp	Pro	Tyr	Ala
				350					355					360
Leu	Val	Arg	Leu	Gly	Thr	Gln	Thr	Phe	Cys	Ser	Arg	Val	Ile	Asp
				365					370					375
Glu	Glu	Leu	Asn	Pro	Gln	Trp	Gly	Glu	Thr	Tyr	Glu	Val	Met	Val
				380					385					390
His	Glu	Val	Pro	Gly	Gln	Glu	Ile	Glu	Val	Glu	Val	Phe	Asp	Lys
				395					400					405

Asp	Pro	Asp	Lys	Asp	Asp	Phe	Leu	Gly	Arg	Met	Lys	Leu	Asp	Val
				410					415					420
Gly	Lys	Val	Leu	Gln	Ala	Ser	Val	Leu	Asp	Asp	Trp	Phe	Pro	Leu
				425					430					435
Gln	Gly	Gly	Gln	Gly	Gln	Val	His	Leu	Arg	Leu	Glu	Trp	Leu	Ser
				440					445					450
Leu	Leu	Ser	Asp	Ala	Glu	Lys	Leu	Glu	Gln	Val	Leu	Gln	Trp	Asn
				455					460					465
Trp	Gly	Val	Ser	Ser	Arg	Pro	Asp	Pro	Pro	Ser	Ala	Ala	Ile	Leu
				470					475					480
Val	Val	Tyr	Leu	Asp	Arg	Ala	Gln	Asp	Leu	Pro	Leu	Lys	Lys	Gly
				485					490					495
Asn	Lys	Glu	Pro	Asn	Pro	Met	Val	Gln	Leu	Ser	Ile	Gln	Asp	Val
				500					505					510
Thr	Gln	Glu	Ser	Lys	Ala	Val	Tyr	Ser	Thr	Asn	Cys	Pro	Val	Trp
				515					520					525
Glu	Glu	Ala	Phe	Arg	Phe	Phe	Leu	Gln	Asp	Pro	Gln	Ser	Gln	Glu
				530					535					540
Leu	Asp	Val	Gln	Val	Lys	Asp	Asp	Ser	Arg	Ala	Leu	Thr	Leu	Gly
				545					550					555
Ala	Leu	Thr	Leu	Pro	Leu	Ala	Arg	Leu	Leu	Thr	Ala	Pro	Glu	Leu
				560					565					570
Ile	Leu	Asp	Gln	Trp	Phe	Gln	Leu	Ser	Ser	Ser	Gly	Pro	Asn	Ser
				575					580					585
Arg	Leu	Tyr	Met	Lys	Leu	Val	Met	Arg	Ile	Leu	Tyr	Leu	Asp	Ser
				590					595					600
Ser	Glu	Ile	Cys	Phe	Pro	Thr	Val	Pro	Gly	Cys	Pro	Gly	Ala	Trp
				605					610					615
Asp	Val	Asp	Ser	Glu	Asn	Pro	Gln	Arg	Gly	Ser	Ser	Val	Asp	Ala
				620					625					630
Pro	Pro	Arg	Pro	Cys	His	Thr	Thr	Pro	Asp	Ser	Gln	Phe	Gly	Thr
				635					640					645
Glu	His	Val	Leu	Arg	Ile	His	Val	Leu	Glu	Ala	Gln	Asp	Leu	Ile
				650					655					660
Ala	Lys	Asp	Arg	Phe	Leu	Gly	Gly	Leu	Val	Lys	Gly	Lys	Ser	Asp
				665					670					675
Pro	Tyr	Val	Lys	Leu	Lys	Leu	Ala	Gly	Arg	Ser	Phe	Arg	Ser	His
				680					685					690
Val	Val	Arg	Glu	Asp	Leu	Asn	Pro	Arg	Trp	Asn	Glu	Val	Phe	Glu
				695					700					705
Val	Ile	Val	Thr	Ser	Val	Pro	Gly	Gln	Glu	Leu	Glu	Val	Glu	Val
				710					715					720
Phe	Asp	Lys	Asp	Leu	Asp	Lys	Asp	Asp	Phe	Leu	Gly	Arg	Cys	Lys
				725					730					735
Val	Arg	Leu	Thr	Thr	Val	Leu	Asn	Ser	Gly	Phe	Leu	Asp	Glu	Trp
				740					745					750
Leu	Thr	Leu	Glu	Asp	Val	Pro	Ser	Gly	Arg	Leu	His	Leu	Arg	Leu
				755					760					765
Glu	Arg	Leu	Thr	Pro	Arg	Pro	Thr	Ala	Ala	Glu	Leu	Glu	Glu	Val
				770					775					780
Leu	Gln	Val	Asn	Ser	Leu	Ile	Gln	Thr	Gln	Lys	Ser	Ala	Glu	Leu
				785					790					795
Ala	Ala	Ala	Leu	Leu	Ser	Ile	Tyr	Met	Glu	Arg	Ala	Glu	Asp	Leu
				800					805					810
Pro	Leu	Arg	Lys	Gly	Thr	Lys	His	Leu	Ser	Pro	Tyr	Ala	Thr	Leu
				815					820					825
Thr	Val	Gly	Asp	Ser	Ser	His	Lys	Thr	Lys	Thr	Ile	Ser	Gln	Thr
				830					835					840
Ser	Ala	Pro	Val	Trp	Asp	Glu	Ser	Ala	Ser	Phe	Leu	Ile	Arg	Lys
				845					850					855
Pro	His	Thr	Glu	Ser	Leu	Glu	Leu	Gln	Val	Arg	Gly	Glu	Gly	Thr
				860					865					870
Gly	Val	Leu	Gly	Ser	Leu	Ser	Leu	Pro	Leu	Ser	Glu	Leu	Leu	Val
				875					880					885
Ala	Asp	Gln	Leu	Cys	Leu	Asp	Arg	Trp	Phe	Thr	Leu	Ser	Ser	Gly
				890					895					900
Gln	Gly	Gln	Val	Leu	Leu	Arg	Ala	Gln	Leu	Gly	Ile	Leu	Val	Ser

Gln	His	Ser	Gly	905	Val	Glu	Ala	His	Ser	910	His	Ser	Tyr	Ser	His	915
Ser	Ser	Ser	Leu	920	Ser	Glu	Glu	Pro	Glu	925	Leu	Ser	Gly	Gly	Pro	930
His	Ile	Thr	Ser	935	Ser	Ala	Pro	Glu	Leu	940	Arg	Gln	Arg	Leu	Thr	945
Val	Asp	Ser	Pro	950	Leu	Glu	Ala	Pro	Ala	955	Gly	Pro	Leu	Gly	Gln	960
Lys	Leu	Thr	Leu	965	Trp	Tyr	Tyr	Ser	Glu	970	Glu	Arg	Lys	Leu	Val	975
Ile	Val	His	Gly	980	Cys	Arg	Ser	Leu	Arg	985	Gln	Asn	Gly	Arg	Asp	990
Pro	Asp	Pro	Tyr	995	Val	Ser	Leu	Leu	Leu	1000	Leu	Pro	Asp	Lys	Asn	1005
Gly	Thr	Lys	Arg	1010	Arg	Thr	Ser	Gln	Lys	1015	Lys	Arg	Thr	Leu	Ser	1020
Glu	Phe	Asn	Glu	1025	Arg	Phe	Glu	Trp	Glu	1030	Leu	Pro	Leu	Asp	Glu	1035
Gln	Arg	Arg	Lys	1040	Leu	Asp	Val	Ser	Val	1045	Lys	Ser	Asn	Ser	Ser	1050
Met	Ser	Arg	Glu	1055	Arg	Glu	Leu	Leu	Gly	1060	Lys	Val	Gln	Leu	Asp	1065
Ala	Glu	Thr	Asp	1070	Leu	Ser	Gln	Gly	Val	1075	Ala	Arg	Trp	Tyr	Asp	1080
Met	Asp	Asn	Lys	1085	Asp	Lys	Gly	Ser	Ser	1090						1095
				1100												

<210> 10

<211> 411

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2285348CD1

<400> 10

Met	Ala	Asn	Tyr	Glu	Ser	Thr	Glu	Val	Met	Gly	Asp	Gly	Glu	Ser	
1				5					10					15	
Ala	His	Asp	Ser	Pro	Arg	Asp	Glu	Ala	Leu	Gln	Asn	Ile	Ser	Ala	
				20					25					30	
Asp	Asp	Leu	Pro	Asp	Ser	Ala	Ser	Gln	Ala	Ala	His	Pro	Gln	Asp	
				35					40					45	
Ser	Ala	Phe	Ser	Tyr	Arg	Asp	Ala	Lys	Lys	Lys	Leu	Arg	Leu	Ala	
				50					55					60	
Leu	Cys	Ser	Ala	Asp	Ser	Val	Ala	Phe	Pro	Val	Leu	Thr	His	Ser	
				65					70					75	
Thr	Arg	Asn	Gly	Leu	Pro	Asp	His	Thr	Asp	Pro	Glu	Asp	Asn	Glu	
				80					85					90	
Ile	Val	Cys	Phe	Leu	Lys	Val	Gln	Ile	Ala	Glu	Ala	Ile	Asn	Leu	
				95					100					105	
Gln	Asp	Lys	Asn	Leu	Met	Ala	Gln	Leu	Gln	Glu	Thr	Met	Arg	Cys	
				110					115					120	
Val	Cys	Arg	Phe	Asp	Asn	Arg	Thr	Cys	Arg	Lys	Leu	Leu	Ala	Ser	
				125					130					135	
Ile	Ala	Glu	Asp	Tyr	Arg	Lys	Arg	Ala	Pro	Tyr	Ile	Ala	Tyr	Leu	
				140					145					150	
Thr	Arg	Cys	Arg	Gln	Gly	Leu	Gln	Thr	Thr	Gln	Ala	His	Leu	Glu	
				155					160					165	
Arg	Leu	Leu	Gln	Arg	Val	Leu	Arg	Asp	Lys	Glu	Val	Ala	Asn	Arg	
				170					175					180	
Tyr	Phe	Thr	Thr	Val	Cys	Val	Arg	Leu	Leu	Leu	Glu	Ser	Lys	Glu	
				185					190					195	
Lys	Lys	Ile	Arg	Glu	Phe	Ile	Gln	Asp	Phe	Gln	Lys	Leu	Thr	Ala	
				200					205					210	
Ala	Asp	Asp	Lys	Thr	Ala	Gln	Val	Glu	Asp	Phe	Leu	Gln	Phe	Leu	

	215		220		225
Tyr Gly Ala Met	Ala Gln Asp Val Ile	Trp Gln Asn Ala Ser	Glu		
	230		235		240
Glu Gln Leu Gln	Asp Ala Gln Leu Ala	Ile Glu Arg Ser Val	Met		
	245		250		255
Asn Arg Ile Phe	Lys Leu Ala Phe Tyr	Pro Asn Gln Asp Gly	Asp		
	260		265		270
Ile Leu Arg Asp	Gln Val Leu His Glu	His Ile Gln Arg Leu	Ser		
	275		280		285
Lys Val Val Thr	Ala Asn His Arg Ala	Leu Gln Ile Pro Glu	Val		
	290		295		300
Tyr Leu Arg Glu	Ala Pro Trp Pro Ser	Ala Gln Ser Glu Ile	Arg		
	305		310		315
Thr Ile Ser Ala	Tyr Lys Thr Pro Arg	Asp Lys Val Gln Cys	Ile		
	320		325		330
Leu Arg Met Cys	Ser Thr Ile Met Asn	Leu Leu Ser Leu Ala	Asn		
	335		340		345
Glu Asp Ser Val	Pro Gly Ala Asp Asp	Phe Val Pro Val Leu	Val		
	350		355		360
Phe Val Leu Ile	Lys Ala Asn Pro Pro	Cys Leu Leu Ser Thr	Val		
	365		370		375
Gln Tyr Ile Ser	Ser Phe Tyr Ala Ser	Cys Leu Ser Gly Glu	Glu		
	380		385		390
Ser Tyr Trp Trp	Met Gln Phe Thr Ala	Ala Val Glu Phe Ile	Lys		
	395		400		405
Thr Ile Asp Asp	Arg Lys				
	410				

<210> 11

<211> 201

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2374186CD1

<400> 11..

Met. Phe Pro Glu Gln	Gln Lys Glu Glu	Phe Val Ser Val	Trp Val
1	5	10	15
Arg Asp Pro Arg Ile	Gln Lys Glu Asp	Phe Trp His Ser Tyr	Ile
	20	25	30
Asp Tyr Glu Ile Cys	Ile His Thr Asn	Ser Met Cys Phe Thr	Met
	35	40	45
Lys Thr Ser Cys Val	Arg Arg Arg Tyr	Arg Glu Phe Val	Trp Leu
	50	55	60
Arg Gln Arg Leu Gln	Ser Asn Ala Leu	Leu Val Gln Leu Pro	Glu
	65	70	75
Leu Pro Ser Lys Asn	Leu Phe Phe Asn	Met Asn Asn Arg Gln	His
	80	85	90
Val Asp Gln Arg Arg	Gln Gly Leu Glu	Asp Phe Leu Arg Lys	Val
	95	100	105
Leu Gln Asn Ala Leu	Leu Leu Leu Ser	Asp Ser Ser Leu His	Leu Phe
	110	115	120
Leu Gln Ser His Leu	Asn Ser Glu Asp	Ile Glu Ala Cys Val	Ser
	125	130	135
Gly Gln Thr Lys Tyr	Ser Val Glu Glu	Ala Ile His Lys Phe	Ala
	140	145	150
Leu Met Asn Arg Arg	Phe Pro Glu Glu	Asp Glu Glu Gly Lys	Lys
	155	160	165
Glu Asn Asp Ile Asp	Tyr Asp Ser Glu	Ser Ser Ser Ser Gly	Leu
	170	175	180
Gly His Ser Ser Asp	Asp Ser Ser Ser	His Gly Cys Lys Val	Asn
	185	190	195
Thr Ala Pro Gln Glu	Ser		
	200		

<210> 12
 <211> 476
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2476232CD1

<400> 12

Met	Ala	Phe	Gly	Lys	Ser	His	Arg	Asp	Pro	Tyr	Ala	Thr	Ser	Val
1				5					10					15
Gly	His	Leu	Ile	Glu	Lys	Ala	Thr	Phe	Ala	Gly	Val	Gln	Thr	Glu
				20					25					30
Asp	Trp	Gly	Gln	Phe	Met	His	Ile	Cys	Asp	Ile	Ile	Asn	Thr	Thr
				35					40					45
Gln	Asp	Gly	Pro	Lys	Asp	Ala	Val	Lys	Ala	Leu	Lys	Lys	Arg	Ile
				50					55					60
Ser	Lys	Asn	Tyr	Asn	His	Lys	Glu	Ile	Gln	Leu	Thr	Leu	Ser	Leu
				65					70					75
Ile	Asp	Met	Cys	Val	Gln	Asn	Cys	Gly	Pro	Ser	Phe	Gln	Ser	Leu
				80					85					90
Ile	Val	Lys	Lys	Glu	Phe	Val	Lys	Glu	Asn	Leu	Val	Lys	Leu	Leu
				95					100					105
Asn	Pro	Arg	Tyr	Asn	Leu	Pro	Leu	Asp	Ile	Gln	Asn	Arg	Ile	Leu
				110					115					120
Asn	Phe	Ile	Lys	Thr	Trp	Ser	Gln	Gly	Phe	Pro	Gly	Gly	Val	Asp
				125					130					135
Val	Ser	Glu	Val	Lys	Glu	Val	Tyr	Leu	Asp	Leu	Val	Lys	Lys	Gly
				140					145					150
Val	Gln	Phe	Pro	Pro	Ser	Glu	Ala	Glu	Ala	Glu	Thr	Ala	Arg	Gln
				155					160					165
Glu	Thr	Ala	Gln	Ile	Ser	Ser	Asn	Pro	Pro	Thr	Ser	Val	Pro	Thr
				170					175					180
Ala	Pro	Ala	Leu	Ser	Ser	Val	Ile	Ala	Pro	Lys	Asn	Ser	Thr	Val
				185					190					195
Thr	Leu	Val	Pro	Glu	Gln	Ile	Gly	Lys	Leu	His	Ser	Glu	Leu	Asp
				200					205					210
Met	Val	Lys	Met	Asn	Val	Arg	Val	Met	Ser	Ala	Ile	Leu	Met	Glu
				215					220					225
Asn	Thr	Pro	Gly	Ser	Glu	Asn	His	Glu	Asp	Ile	Glu	Leu	Leu	Gln
				230					235					240
Lys	Leu	Tyr	Lys	Thr	Gly	Arg	Glu	Met	Gln	Glu	Arg	Ile	Met	Asp
				245					250					255
Leu	Leu	Val	Val	Val	Glu	Asn	Glu	Asp	Val	Thr	Val	Glu	Leu	Ile
				260					265					270
Gln	Val	Asn	Glu	Asp	Leu	Asn	Asn	Ala	Ile	Leu	Gly	Tyr	Glu	Arg
				275					280					285
Phe	Thr	Arg	Asn	Gln	Gln	Arg	Ile	Leu	Glu	Gln	Asn	Lys	Asn	Gln
				290					295					300
Lys	Glu	Ala	Thr	Asn	Thr	Thr	Ser	Glu	Pro	Ser	Ala	Pro	Ser	Gln
				305					310					315
Asp	Leu	Leu	Asp	Leu	Ser	Pro	Ser	Pro	Arg	Met	Pro	Arg	Ala	Thr
				320					325					330
Leu	Gly	Glu	Leu	Asn	Thr	Met	Asn	Asn	Gln	Leu	Ser	Gly	Leu	Asn
				335					340					345
Phe	Ser	Leu	Pro	Ser	Ser	Asp	Val	Thr	Asn	Asn	Leu	Lys	Pro	Ser
				350					355					360
Leu	His	Pro	Gln	Met	Asn	Leu	Leu	Ala	Leu	Glu	Asn	Thr	Glu	Ile
				365					370					375
Pro	Pro	Phe	Ala	Gln	Arg	Thr	Ser	Gln	Asn	Leu	Thr	Ser	Ser	His
				380					385					390
Ala	Tyr	Asp	Asn	Phe	Leu	Glu	His	Ser	Asn	Ser	Val	Phe	Leu	Gln
				395					400					405
Pro	Val	Ser	Leu	Gln	Thr	Ile	Ala	Ala	Ala	Pro	Ser	Asn	Gln	Ser
				410					415					420
Leu	Pro	Pro	Leu	Pro	Ser	Asn	His	Pro	Ala	Met	Thr	Lys	Ser	Asp

Leu	Gln	Pro	Pro	Asn	Tyr	Tyr	Glu	Val	Met	Glu	Phe	Asp	Pro	Leu
				425					430					435
Ala	Pro	Ala	Val	Thr	Thr	Glu	Ala	Ile	Tyr	Glu	Glu	Ile	Asp	Ala
				440					445					450
His	Gln	His	Lys	Gly	Ala	Gln	Asn	Asp	Gly	Asp				
				455					460					465
				470					475					

<210> 13
 <211> 1220
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2503986CD1

<400> 13

Met	Lys	Leu	Lys	Glu	Val	Asp	Arg	Thr	Ala	Met	Gln	Ala	Trp	Ser
1				5					10					15
Pro	Ala	Gln	Asn	His	Pro	Ile	Tyr	Leu	Ala	Thr	Gly	Thr	Ser	Ala
				20					25					30
Gln	Gln	Leu	Asp	Ala	Thr	Phe	Ser	Thr	Asn	Ala	Ser	Leu	Glu	Ile
				35					40					45
Phe	Glu	Leu	Asp	Leu	Ser	Asp	Pro	Ser	Leu	Asp	Met	Lys	Ser	Cys
				50					55					60
Ala	Thr	Phe	Ser	Ser	Ser	His	Arg	Tyr	His	Lys	Leu	Ile	Trp	Gly
				65					70					75
Pro	Tyr	Lys	Met	Asp	Ser	Lys	Gly	Asp	Val	Ser	Gly	Val	Leu	Ile
				80					85					90
Ala	Gly	Gly	Glu	Asn	Gly	Asn	Ile	Ile	Leu	Tyr	Asp	Pro	Ser	Lys
				95					100					105
Ile	Ile	Ala	Gly	Asp	Lys	Glu	Val	Val	Ile	Ala	Gln	Asn	Asp	Lys
				110					115					120
His	Thr	Gly	Pro	Val	Arg	Ala	Leu	Asp	Val	Asn	Ile	Phe	Gln	Thr
				125					130					135
Asn	Leu	Val	Ala	Ser	Gly	Ala	Asn	Glu	Ser	Glu	Ile	Tyr	Ile	Trp
				140					145					150
Asp	Leu	Asn	Asn	Phe	Ala	Thr	Pro	Met	Thr	Pro	Gly	Ala	Lys	Thr
				155					160					165
Gln	Pro	Pro	Glu	Asp	Ile	Ser	Cys	Ile	Ala	Trp	Asn	Arg	Gln	Val
				170					175					180
Gln	His	Ile	Leu	Ala	Ser	Ala	Ser	Pro	Ser	Gly	Arg	Ala	Thr	Val
				185					190					195
Trp	Asp	Leu	Arg	Lys	Asn	Glu	Pro	Ile	Ile	Lys	Val	Ser	Asp	His
				200					205					210
Ser	Asn	Arg	Met	His	Cys	Ser	Gly	Leu	Ala	Trp	His	Pro	Asp	Val
				215					220					225
Ala	Thr	Gln	Met	Val	Leu	Ala	Ser	Glu	Asp	Asp	Arg	Leu	Pro	Val
				230					235					240
Ile	Gln	Met	Trp	Asp	Leu	Arg	Phe	Ala	Ser	Ser	Pro	Leu	Arg	Val
				245					250					255
Leu	Glu	Asn	His	Ala	Arg	Gly	Ile	Leu	Ala	Ile	Ala	Trp	Ser	Met
				260					265					270
Ala	Asp	Pro	Glu	Leu	Leu	Leu	Ser	Cys	Gly	Lys	Asp	Ala	Lys	Ile
				275					280					285
Leu	Cys	Ser	Asn	Pro	Asn	Thr	Gly	Glu	Val	Leu	Tyr	Glu	Leu	Pro
				290					295					300
Thr	Asn	Thr	Gln	Trp	Cys	Phe	Asp	Ile	Gln	Trp	Cys	Pro	Arg	Asn
				305					310					315
Pro	Ala	Val	Leu	Ser	Ala	Ala	Ser	Phe	Asp	Gly	Arg	Ile	Ser	Val
				320					325					330
Tyr	Ser	Ile	Met	Gly	Gly	Ser	Thr	Asp	Gly	Leu	Arg	Gln	Lys	Gln
				335					340					345
Val	Asp	Lys	Leu	Ser	Ser	Ser	Phe	Gly	Asn	Leu	Asp	Pro	Phe	Gly
				350					355					360
Thr	Gly	Gln	Pro	Leu	Pro	Pro	Leu	Gln	Ile	Pro	Gln	Gln	Thr	Ala

				365					370					375
Gln	His	Ser	Ile	Val	Leu	Pro	Leu	Lys	Lys	Pro	Pro	Lys	Trp	Ile
				380					385					390
Arg	Arg	Pro	Val	Gly	Ala	Ser	Phe	Ser	Phe	Gly	Gly	Lys	Leu	Val
				395					400					405
Thr	Phe	Glu	Asn	Val	Arg	Met	Pro	Ser	His	Gln	Gly	Ala	Glu	Gln
				410					415					420
Gln	Gln	Gln	Gln	His	His	Val	Phe	Ile	Ser	Gln	Val	Val	Thr	Glu
				425					430					435
Lys	Glu	Phe	Leu	Ser	Arg	Ser	Asp	Gln	Leu	Gln	Gln	Ala	Val	Gln
				440					445					450
Ser	Gln	Gly	Phe	Ile	Asn	Tyr	Cys	Gln	Lys	Lys	Ile	Asp	Ala	Ser
				455					460					465
Gln	Thr	Glu	Phe	Glu	Lys	Asn	Val	Trp	Ser	Phe	Leu	Lys	Val	Asn
				470					475					480
Phe	Glu	Asp	Asp	Ser	Arg	Gly	Lys	Tyr	Leu	Glu	Leu	Leu	Gly	Tyr
				485					490					495
Arg	Lys	Glu	Asp	Leu	Gly	Lys	Lys	Ile	Ala	Leu	Ala	Leu	Asn	Lys
				500					505					510
Val	Asp	Gly	Ala	Asn	Val	Ala	Leu	Lys	Asp	Ser	Asp	Gln	Val	Ala
				515					520					525
Gln	Ser	Asp	Gly	Glu	Glu	Ser	Pro	Ala	Ala	Glu	Glu	Gln	Leu	Leu
				530					535					540
Gly	Glu	His	Ile	Lys	Glu	Glu	Lys	Glu	Glu	Ser	Glu	Phe	Leu	Pro
				545					550					555
Ser	Ser	Gly	Gly	Thr	Phe	Asn	Ile	Ser	Val	Ser	Gly	Asp	Ile	Asp
				560					565					570
Gly	Leu	Ile	Thr	Gln	Ala	Leu	Leu	Thr	Gly	Asn	Phe	Glu	Ser	Ala
				575					580					585
Val	Asp	Leu	Cys	Leu	His	Asp	Asn	Arg	Met	Ala	Asp	Ala	Ile	Ile
				590					595					600
Leu	Ala	Ile	Ala	Gly	Gly	Gln	Glu	Leu	Leu	Ala	Arg	Thr	Gln	Lys
				605					610					615
Lys	Tyr	Phe	Ala	Lys	Ser	Gln	Ser	Lys	Ile	Thr	Arg	Leu	Ile	Thr
				620					625					630
Ala	Val	Val	Met	Lys	Asn	Trp	Lys	Glu	Ile	Val	Glu	Ser	Cys	Asp
				635					640					645
Leu	Lys	Asn	Trp	Arg	Glu	Ala	Leu	Ala	Ala	Val	Leu	Thr	Tyr	Ala
				650					655					660
Lys	Pro	Asp	Glu	Phe	Ser	Ala	Leu	Cys	Asp	Leu	Leu	Gly	Thr	Arg
				665					670					675
Leu	Glu	Asn	Glu	Gly	Asp	Ser	Leu	Leu	Gln	Thr	Gln	Ala	Cys	Leu
				680					685					690
Cys	Tyr	Ile	Cys	Ala	Gly	Asn	Val	Glu	Lys	Leu	Val	Ala	Cys	Trp
				695					700					705
Thr	Lys	Ala	Gln	Asp	Gly	Ser	His	Pro	Leu	Ser	Leu	Gln	Asp	Leu
				710					715					720
Ile	Glu	Lys	Val	Val	Ile	Leu	Arg	Lys	Ala	Val	Gln	Leu	Thr	Gln
				725					730					735
Ala	Met	Asp	Thr	Ser	Thr	Val	Gly	Val	Leu	Leu	Ala	Ala	Lys	Met
				740					745					750
Ser	Gln	Tyr	Ala	Asn	Leu	Leu	Ala	Ala	Gln	Gly	Ser	Ile	Ala	Ala
				755					760					765
Ala	Leu	Ala	Phe	Leu	Pro	Asp	Asn	Thr	Asn	Gln	Pro	Asn	Ile	Met
				770					775					780
Gln	Leu	Arg	Asp	Arg	Leu	Cys	Arg	Ala	Gln	Gly	Glu	Pro	Val	Ala
				785					790					795
Gly	His	Glu	Ser	Pro	Lys	Ile	Pro	Tyr	Glu	Lys	Gln	Gln	Leu	Pro
				800					805					810
Lys	Gly	Arg	Pro	Gly	Pro	Val	Ala	Gly	His	His	Gln	Met	Pro	Arg
				815					820					825
Val	Gln	Thr	Gln	Gln	Tyr	Tyr	Pro	His	Gly	Glu	Asn	Pro	Pro	Pro
				830					835					840
Pro	Gly	Phe	Ile	Met	His	Gly	Asn	Val	Asn	Pro	Asn	Ala	Ala	Gly
				845					850					855
Gln	Leu	Pro	Thr	Ser	Pro	Gly	His	Met	His	Thr	Gln	Val	Pro	Pro
				860					865					870

```

Tyr Pro Gln Pro Gln Pro Tyr Gln Pro Ala Gln Pro Tyr Pro Phe
875 880 885
Gly Thr Gly Gly Ser Ala Met Tyr Arg Pro Gln Gln Pro Val Ala
890 895 900
Pro Pro Thr Ser Asn Ala Tyr Pro Asn Thr Pro Tyr Ile Ser Ser
905 910 915
Ala Ser Ser Tyr Thr Gly Gln Ser Gln Leu Tyr Ala Ala Gln His
920 925 930
Gln Ala Ser Ser Pro Thr Ser Ser Pro Ala Thr Ser Phe Pro Pro
935 940 945
Pro Pro Ser Ser Gly Ala Ser Phe Gln His Gly Gly Pro Gly Ala
950 955 960
Pro Pro Ser Ser Ser Ala Tyr Ala Leu Pro Pro Gly Thr Thr Gly
965 970 975
Thr Leu Pro Ala Ala Ser Glu Leu Pro Ala Ser Gln Arg Thr Gly
980 985 990
Pro Gln Asn Gly Trp Asn Asp Pro Pro Ala Leu Asn Arg Val Pro
995 1000 1005
Lys Lys Lys Lys Met Pro Glu Asn Phe Met Pro Pro Val Pro Ile
1010 1015 1020
Thr Ser Pro Ile Met Asn Pro Leu Gly Asp Pro Gln Ser Gln Met
1025 1030 1035
Leu Gln Gln Gln Pro Ser Ala Pro Val Pro Leu Ser Ser Gln Ser
1040 1045 1050
Ser Phe Pro Gln Pro His Leu Pro Gly Gly Gln Pro Phe His Gly
1055 1060 1065
Val Gln Gln Pro Leu Gly Gln Thr Gly Met Pro Pro Ser Phe Ser
1070 1075 1080
Lys Pro Asn Ile Glu Gly Ala Pro Gly Ala Pro Ile Gly Asn Thr
1085 1090 1095
Phe Gln His Val Gln Ser Leu Pro Thr Lys Lys Ile Thr Lys Lys
1100 1105 1110
Pro Ile Pro Asp Glu His Leu Ile Leu Lys Thr Thr Phe Glu Asp
1115 1120 1125
Leu Ile Gln Arg Cys Leu Ser Ser Ala Thr Asp Pro Gln Thr Lys
1130 1135 1140
Arg Lys Leu Asp Asp Ala Ser Lys Arg Leu Glu Phe Leu Tyr Asp
1145 1150 1155
Lys Leu Arg Glu Gln Thr Leu Ser Pro Thr Ile Thr Ser Gly Leu
1160 1165 1170
His Asn Ile Ala Arg Ser Ile Glu Thr Arg Asn Tyr Ser Glu Gly
1175 1180 1185
Leu Thr Met His Thr His Ile Val Ser Thr Ser Asn Phe Ser Glu
1190 1195 1200
Thr Ser Ala Phe Met Pro Val Leu Lys Val Val Leu Thr Gln Ala
1205 1210 1215
Asn Lys Leu Gly Val
1220

```

<210> 14
 <211> 222
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID: No: 2596566CD1

```

<400> 14
Met Ala Phe Val Lys Ser Gly Trp Leu Leu Arg Gln Ser Thr Ile
1 5 10 15
Leu Lys Arg Trp Lys Lys Asn Trp Phe Asp Leu Trp Ser Asp Gly
20 25 30
His Leu Ile Tyr Tyr Asp Asp Gln Thr Arg Gln Asn Ile Glu Asp
35 40 45
Lys Val His Met Pro Met Asp Cys Ile Asn Ile Arg Thr Gly Gln
50 55 60

```

Glu	Cys	Arg	Asp	Thr	Gln	Pro	Pro	Asp	Gly	Lys	Ser	Lys	Asp	Cys
				65					70					75
Met	Leu	Gln	Ile	Val	Cys	Arg	Asp	Gly	Lys	Thr	Ile	Ser	Leu	Cys
				80					85					90
Ala	Glu	Ser	Thr	Asp	Asp	Cys	Leu	Ala	Trp	Lys	Phe	Thr	Leu	Gln
				95					100					105
Asp	Ser	Arg	Thr	Asn	Thr	Ala	Tyr	Val	Gly	Ser	Ala	Val	Met	Thr
				110					115					120
Asp	Glu	Thr	Ser	Val	Val	Ser	Ser	Pro	Pro	Pro	Tyr	Thr	Ala	Tyr
				125					130					135
Ala	Ala	Pro	Ala	Pro	Glu	Gln	Ala	Tyr	Gly	Tyr	Gly	Pro	Tyr	Gly
				140					145					150
Gly	Ala	Tyr	Pro	Pro	Gly	Thr	Gln	Val	Val	Tyr	Ala	Ala	Asn	Gly
				155					160					165
Gln	Ala	Tyr	Ala	Val	Pro	Tyr	Gln	Tyr	Pro	Tyr	Ala	Gly	Leu	Tyr
				170					175					180
Gly	Gln	Gln	Pro	Ala	Asn	Gln	Val	Ile	Ile	Arg	Glu	Arg	Tyr	Arg
				185					190					195
Asp	Asn	Asp	Ser	Asp	Leu	Ala	Leu	Gly	Met	Leu	Ala	Gly	Ala	Ala
				200					205					210
Thr	Gly	Met	Ala	Leu	Gly	Ser	Leu	Phe	Trp	Val	Phe			
				215					220					

<210> 15

<211> 924

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2685253CD1

<400> 15

Met	Ser	Arg	Ser	Arg	Gln	Pro	Pro	Leu	Val	Thr	Gly	Ile	Ser	Pro
1				5					10					15
Asn	Glu	Gly	Ile	Pro	Trp	Thr	Lys	Val	Thr	Ile	Arg	Gly	Glu	Asn
				20					25					30
Leu	Gly	Thr	Gly	Pro	Thr	Asp	Leu	Ile	Gly	Leu	Thr	Ile	Cys	Gly
				35					40					45
His	Asn	Cys	Leu	Leu	Thr	Ala	Glu	Trp	Met	Ser	Ala	Ser	Lys	Ile
				50					55					60
Val	Cys	Arg	Val	Gly	Gln	Ala	Lys	Asn	Asp	Lys	Gly	Asp	Ile	Ile
				65					70					75
Val	Thr	Thr	Lys	Ser	Gly	Gly	Arg	Gly	Thr	Ser	Thr	Val	Ser	Phe
				80					85					90
Lys	Leu	Leu	Lys	Pro	Glu	Lys	Ile	Gly	Ile	Leu	Asp	Gln	Ser	Ala
				95					100					105
Val	Trp	Val	Asp	Glu	Met	Asn	Tyr	Tyr	Asp	Met	Arg	Thr	Asp	Arg
				110					115					120
Asn	Lys	Gly	Ile	Pro	Pro	Leu	Ser	Leu	Arg	Pro	Ala	Asn	Pro	Leu
				125					130					135
Gly	Ile	Glu	Ile	Glu	Lys	Ser	Lys	Phe	Ser	Gln	Lys	Asp	Leu	Glu
				140					145					150
Met	Leu	Phe	His	Gly	Met	Ser	Ala	Asp	Phe	Thr	Ser	Glu	Asn	Phe
				155					160					165
Ser	Ala	Ala	Trp	Tyr	Leu	Ile	Glu	Asn	His	Ser	Asn	Thr	Ser	Phe
				170					175					180
Glu	Gln	Leu	Lys	Met	Ala	Val	Thr	Asn	Leu	Lys	Arg	Gln	Ala	Asn
				185					190					195
Lys	Lys	Ser	Glu	Gly	Ser	Leu	Ala	Tyr	Val	Lys	Gly	Gly	Leu	Ser
				200					205					210
Thr	Phe	Phe	Glu	Ala	Gln	Asp	Ala	Leu	Ser	Ala	Ile	His	Gln	Lys
				215					220					225
Leu	Glu	Ala	Asp	Gly	Thr	Glu	Lys	Val	Glu	Gly	Ser	Met	Thr	Gln
				230					235					240
Lys	Leu	Glu	Asn	Val	Leu	Asn	Arg	Ala	Ser	Asn	Thr	Ala	Asp	Thr
				245					250					255

Leu	Phe	Gln	Glu	Val	Leu	Gly	Arg	Lys	Asp	Lys	Ala	Asp	Ser	Thr
				260					265					270
Arg	Asn	Ala	Leu	Asn	Val	Leu	Gln	Arg	Phe	Lys	Phe	Leu	Phe	Asn
				275					280					285
Leu	Pro	Leu	Asn	Ile	Glu	Arg	Asn	Ile	Gln	Lys	Gly	Asp	Tyr	Asp
				290					295					300
Val	Val	Ile	Asn	Asp	Tyr	Glu	Lys	Ala	Lys	Ser	Leu	Phe	Gly	Lys
				305					310					315
Thr	Glu	Val	Gln	Val	Phe	Lys	Lys	Tyr	Tyr	Ala	Glu	Val	Glu	Thr
				320					325					330
Arg	Ile	Glu	Ala	Leu	Arg	Glu	Leu	Leu	Leu	Asp	Lys	Leu	Leu	Glu
				335					340					345
Thr	Pro	Ser	Thr	Leu	His	Asp	Gln	Lys	Arg	Tyr	Ile	Arg	Tyr	Leu
				350					355					360
Ser	Asp	Leu	His	Ala	Ser	Gly	Asp	Pro	Ala	Trp	Gln	Cys	Ile	Gly
				365					370					375
Ala	Gln	His	Lys	Trp	Ile	Leu	Gln	Leu	Met	His	Ser	Cys	Lys	Glu
				380					385					390
Gly	Tyr	Val	Lys	Asp	Leu	Lys	Gly	Asn	Pro	Gly	Leu	His	Ser	Pro
				395					400					405
Met	Leu	Asp	Leu	Asp	Asn	Asp	Thr	Arg	Pro	Ser	Val	Leu	Gly	His
				410					415					420
Leu	Ser	Gln	Thr	Ala	Ser	Leu	Lys	Arg	Gly	Ser	Ser	Phe	Gln	Ser
				425					430					435
Gly	Arg	Asp	Asp	Thr	Trp	Arg	Tyr	Lys	Thr	Pro	His	Arg	Val	Ala
				440					445					450
Phe	Val	Glu	Lys	Leu	Thr	Lys	Leu	Val	Leu	Ser	Gln	Leu	Pro	Asn
				455					460					465
Phe	Trp	Lys	Leu	Trp	Ile	Ser	Tyr	Val	Asn	Gly	Ser	Leu	Phe	Ser
				470					475					480
Glu	Thr	Ala	Glu	Lys	Ser	Gly	Gln	Ile	Glu	Arg	Ser	Lys	Asn	Val
				485					490					495
Arg	Gln	Arg	Gln	Asn	Asp	Phe	Lys	Lys	Met	Ile	Gln	Glu	Val	Met
				500					505					510
His	Ser	Leu	Val	Lys	Leu	Thr	Arg	Gly	Ala	Leu	Leu	Pro	Leu	Ser
				515					520					525
Ile	Arg	Asp	Gly	Glu	Ala	Lys	Gln	Tyr	Gly	Gly	Trp	Glu	Val	Lys
				530					535					540
Cys	Glu	Leu	Ser	Gly	Gln	Trp	Leu	Ala	His	Ala	Ile	Gln	Thr	Val
				545					550					555
Arg	Leu	Thr	His	Glu	Ser	Leu	Thr	Ala	Leu	Glu	Ile	Pro	Asn	Asp
				560					565					570
Leu	Leu	Gln	Thr	Ile	Gln	Asp	Leu	Ile	Leu	Asp	Leu	Arg	Val	Arg
				575					580					585
Cys	Val	Met	Ala	Thr	Leu	Gln	His	Thr	Ala	Glu	Glu	Ile	Lys	Arg
				590					595					600
Leu	Ala	Glu	Lys	Glu	Asp	Trp	Ile	Val	Asp	Asn	Glu	Gly	Leu	Thr
				605					610					615
Ser	Leu	Pro	Cys	Gln	Phe	Glu	Gln	Cys	Ile	Val	Cys	Ser	Leu	Gln
				620					625					630
Ser	Leu	Lys	Gly	Val	Leu	Glu	Cys	Lys	Pro	Gly	Glu	Ala	Ser	Val
				635					640					645
Phe	Gln	Gln	Pro	Lys	Thr	Gln	Glu	Glu	Val	Cys	Gln	Leu	Ser	Ile
				650					655					660
Asn	Ile	Met	Gln	Val	Phe	Ile	Tyr	Cys	Leu	Glu	Gln	Leu	Ser	Thr
				665					670					675
Lys	Pro	Asp	Ala	Asp	Ile	Asp	Thr	Thr	His	Leu	Ser	Val	Asp	Val
				680					685					690
Ser	Ser	Pro	Asp	Leu	Phe	Gly	Ser	Ile	His	Glu	Asp	Phe	Ser	Leu
				695					700					705
Thr	Ser	Glu	Gln	Arg	Leu	Leu	Ile	Val	Leu	Ser	Asn	Cys	Cys	Tyr
				710					715					720
Leu	Glu	Arg	His	Thr	Phe	Leu	Asn	Ile	Ala	Glu	His	Phe	Glu	Lys
				725					730					735
His	Asn	Phe	Gln	Gly	Ile	Glu	Lys	Ile	Thr	Gln	Val	Ser	Met	Ala
				740					745					750
Ser	Leu	Lys	Glu	Leu	Asp	Gln	Arg	Leu	Phe	Glu	Asn	Tyr	Ile	Glu

Leu Lys Ala Asp	755	Pro Ile Val Gly Ser	760	Leu Glu Pro Gly Ile	765
Ala Gly Tyr Phe	770	Asp Trp Lys Asp Cys	775	Leu Pro Pro Thr Gly	780
Arg Asn Tyr Leu	785	Lys Glu Ala Leu Val	790	Asn Ile Ile Ala Val	795
Ala Glu Val Phe	800	Thr Ile Ser Lys Glu	805	Leu Val Pro Arg Val	810
Ser Lys Val Ile	815	Glu Ala Val Ser Glu	820	Glu Leu Ser Arg Leu	825
Gln Cys Val Ser	830	Ser Phe Ser Lys Asn	835	Gly Ala Leu Gln Ala	840
Leu Glu Ile Cys	845	Ala Leu Arg Asp Thr	850	Val Ala Val Tyr Leu	855
Pro Glu Ser Lys	860	Ser Ser Phe Lys Gln	865	Ala Leu Glu Ala Leu	870
Gln Leu Ser Ser	875	Gly Ala Asp Lys Lys	880	Leu Leu Glu Glu Leu	885
Asn Lys Phe Lys	890	Ser Ser Met His Leu	895	Gln Leu Thr Cys Phe	900
Ala Ala Ser Ser	905	Met Met Lys Thr	910		915
	920				

<210> 16

<211> 435

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2762252CD1

<400> 16

Met Ala Pro Phe Gly	Arg Asn Leu Leu Lys	Thr Arg His Lys Asn
1 5	10	15
Arg Ser Pro Thr Lys	Asp Met Asp Ser Glu	Lys Glu Ile Val
20 25	30	35
Val Trp Val Cys Gln	Glu Glu Lys Leu Val	Cys Gly Leu Thr Lys
35 40	45	50
Arg Thr Thr Ser Ala	Asp Val Ile Gln Ala	Leu Leu Glu Glu His
50 55	60	65
Glu Ala Thr Phe Gly	Glu Lys Arg Phe Leu	Leu Gly Lys Pro Ser
65 70	75	80
Asp Tyr Cys Ile Ile	Glu Lys Trp Arg Gly	Ser Glu Arg Val Leu
80 85	90	95
Pro Pro Leu Thr Arg	Ile Leu Lys Leu Trp	Lys Ala Trp Gly Asp
95 100	105	110
Glu Gln Pro Asn Met	Gln Phe Val Leu Val	Lys Ala Asp Ala Phe
110 115	120	125
Leu Pro Val Pro Leu	Trp Arg Thr Ala Glu	Ala Lys Leu Val Gln
125 130	135	140
Asn Thr Glu Lys Leu	Trp Glu Leu Ser Pro	Ala Asn Tyr Met Lys
140 145	150	155
Thr Leu Pro Pro Asp	Lys Gln Lys Arg Ile	Val Arg Lys Thr Phe
155 160	165	170
Arg Lys Leu Ala Lys	Ile Lys Gln Asp Thr	Val Ser His Asp Arg
170 175	180	185
Asp Asn Met Glu Thr	Leu Val His Leu Ile	Ile Ser Gln Asp His
185 190	195	200
Thr Ile His Gln Gln	Val Lys Arg Met Lys	Glu Leu Asp Leu Glu
200 205	210	215
Ile Glu Lys Cys Glu	Ala Lys Phe His Leu	Asp Arg Val Glu Asn
215 220	225	230
Asp Gly Glu Asn Tyr	Val Gln Asp Ala Tyr	Leu Met Pro Ser Phe
230 235	240	245
Ser Glu Val Glu Gln	Asn Leu Asp Leu Gln	Tyr Glu Glu Asn Gln

Thr	Leu	Glu	Asp	245	Leu	Ser	Glu	Ser	Asp	250	Gly	Ile	Glu	Gln	Leu	Glu	255
				260						265							270
Glu	Arg	Leu	Lys	275	Tyr	Tyr	Arg	Ile	Leu	280	Ile	Asp	Lys	Leu	Ser	Ala	285
				290						295							300
Glu	Ile	Glu	Lys	305	Glu	Val	Lys	Ser	Val	310	Cys	Ile	Asp	Ile	Asn	Glu	315
				320						325							330
Asp	Ala	Glu	Gly	335	Glu	Ala	Ala	Ser	Glu	340	Leu	Glu	Ser	Ser	Asn	Leu	345
				350						355							360
Glu	Ser	Val	Lys	365	Cys	Asp	Leu	Glu	Lys	370	Ser	Met	Lys	Ala	Gly	Leu	375
				380						385							390
Lys	Ile	His	Ser	395	His	Leu	Ser	Gly	Ile	400	Gln	Lys	Glu	Ile	Lys	Tyr	405
				410						415							420
Ser	Asp	Ser	Leu	425	Leu	Gln	Met	Lys	Ala	430	Lys	Glu	Tyr	Glu	Leu	Leu	435
Ala	Lys	Glu	Phe		Asn	Ser	Leu	His	Ile		Ser	Asn	Lys	Asp	Gly	Cys	
Gln	Leu	Lys	Glu		Asn	Arg	Ala	Lys	Glu		Ser	Glu	Val	Pro	Ser	Ser	
Asn	Gly	Glu	Ile		Pro	Pro	Phe	Thr	Gln		Arg	Val	Phe	Ser	Asn	Tyr	
Thr	Asn	Asp	Thr		Asp	Ser	Asp	Thr	Gly		Ile	Ser	Ser	Asn	His	Ser	
Gln	Asp	Ser	Glu		Thr	Thr	Val	Gly	Asp		Val	Val	Leu	Leu	Ser	Thr	

<210> 17

<211> 321

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3452009CD1

<400> 17

Met	Glu	Phe	Leu	Leu	Gly	Asn	Pro	Phe	Ser	Thr	Pro	Val	Gly	Gln			
1				5					10					15			
Cys	Leu	Glu	Lys	Ala	Thr	Asp	Gly	Ser	Leu	Gln	Ser	Glu	Asp	Trp			
				20					25					30			
Thr	Leu	Asn	Met	Glu	Ile	Cys	Asp	Ile	Ile	Asn	Glu	Thr	Glu	Glu			
				35					40					45			
Gly	Pro	Lys	Asp	Ala	Ile	Arg	Ala	Leu	Lys	Lys	Arg	Leu	Asn	Gly			
				50					55					60			
Asn	Arg	Asn	Tyr	Arg	Glu	Val	Met	Leu	Ala	Leu	Thr	Val	Leu	Glu			
				65					70					75			
Thr	Cys	Val	Lys	Asn	Cys	Gly	His	Arg	Phe	His	Ile	Leu	Val	Ala			
				80					85					90			
Asn	Arg	Asp	Phe	Ile	Asp	Ser	Val	Leu	Val	Lys	Ile	Ile	Ser	Pro			
				95					100					105			
Lys	Asn	Asn	Pro	Pro	Thr	Ile	Val	Gln	Asp	Lys	Val	Leu	Ala	Leu			
				110					115					120			
Ile	Gln	Ala	Trp	Ala	Asp	Ala	Phe	Arg	Ser	Ser	Pro	Asp	Leu	Thr			
				125					130					135			
Gly	Val	Val	His	Ile	Tyr	Glu	Glu	Leu	Lys	Arg	Lys	Gly	Val	Glu			
				140					145					150			
Phe	Pro	Met	Ala	Asp	Leu	Asp	Ala	Leu	Ser	Pro	Ile	His	Thr	Pro			
				155					160					165			
Gln	Arg	Ser	Val	Pro	Glu	Val	Asp	Pro	Ala	Ala	Thr	Met	Pro	Arg			
				170					175					180			
Ser	Gln	Ser	Gln	Gln	Arg	Thr	Ser	Ala	Gly	Ser	Tyr	Ser	Ser	Pro			
				185					190					195			
Pro	Pro	Ala	Pro	Tyr	Ser	Ala	Pro	Gln	Ala	Pro	Ala	Leu	Ser	Val			
				200					205					210			
Thr	Gly	Pro	Ile	Thr	Ala	Asn	Ser	Glu	Gln	Ile	Ala	Arg	Leu	Arg			
				215					220					225			

Ser	Glu	Leu	Asp	Val	Val	Arg	Gly	Asn	Thr	Lys	Val	Met	Ser	Glu
				230					235					240
Met	Leu	Thr	Glu	Met	Val	Pro	Gly	Gln	Glu	Asp	Ser	Ser	Asp	Leu
				245					250					255
Glu	Leu	Leu	Gln	Glu	Leu	Asn	Arg	Thr	Cys	Arg	Ala	Met	Gln	Gln
				260					265					270
Arg	Ile	Val	Glu	Leu	Ile	Ser	Arg	Val	Ser	Asn	Glu	Glu	Val	Thr
				275					280					285
Glu	Glu	Leu	Leu	His	Val	Asn	Asp	Asp	Leu	Asn	Asn	Val	Phe	Leu
				290					295					300
Arg	Tyr	Glu	Arg	Trp	Glu	Pro	Asp	Phe	Phe	Phe	Phe	Phe	Phe	Pro
				305					310					315
Leu	Lys	Arg	Leu	Leu	Pro									
				320										

<210> 18

<211> 499

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4644780CD1

<400> 18

Met	Pro	Ala	Val	Ser	Gly	Pro	Gly	Pro	Leu	Phe	Cys	Leu	Leu	Leu
1				5					10					15
Leu	Leu	Leu	Asp	Pro	His	Ser	Pro	Glu	Thr	Gly	Cys	Pro	Pro	Leu
				20					25					30
Arg	Arg	Phe	Glu	Tyr	Lys	Leu	Ser	Phe	Lys	Gly	Pro	Arg	Leu	Ala
				35					40					45
Leu	Pro	Gly	Ala	Gly	Ile	Pro	Phe	Trp	Ser	His	His	Gly	Asp	Ala
				50					55					60
Ile	Leu	Gly	Leu	Glu	Glu	Val	Arg	Leu	Thr	Pro	Ser	Met	Arg	Asn
				65					70					75
Arg	Ser	Gly	Ala	Val	Trp	Ser	Arg	Ala	Ser	Val	Pro	Phe	Ser	Ala
				80					85					90
Trp	Glu	Val	Glu	Val	Gln	Met	Arg	Val	Thr	Gly	Leu	Gly	Arg	Arg
				95					100					105
Gly	Ala	Gln	Gly	Met	Ala	Val	Trp	Tyr	Thr	Arg	Gly	Arg	Gly	His
				110					115					120
Val	Gly	Ser	Val	Leu	Gly	Gly	Leu	Ala	Ser	Trp	Asp	Gly	Ile	Gly
				125					130					135
Ile	Phe	Phe	Asp	Ser	Pro	Ala	Glu	Asp	Thr	Gln	Asp	Ser	Pro	Ala
				140					145					150
Ile	Arg	Val	Leu	Ala	Ser	Asp	Gly	His	Ile	Pro	Ser	Glu	Gln	Pro
				155					160					165
Gly	Asp	Gly	Ala	Ser	Gln	Gly	Leu	Gly	Ser	Cys	His	Trp	Asp	Phe
				170					175					180
Arg	Asn	Arg	Pro	His	Pro	Phe	Arg	Ala	Arg	Ile	Thr	Tyr	Trp	Gly
				185					190					195
Gln	Arg	Leu	Arg	Met	Ser	Leu	Asn	Ser	Gly	Leu	Thr	Pro	Ser	Asp
				200					205					210
Pro	Asp	Asp	His	Asp	Val	Leu	Ser	Phe	Leu	Thr	Phe	Ser	Leu	Ser
				215					220					225
Glu	Pro	Ser	Pro	Glu	Val	Pro	Pro	Gln	Pro	Phe	Leu	Glu	Met	Gln
				230					235					240
Gln	Leu	Arg	Leu	Ala	Arg	Gln	Leu	Glu	Gly	Leu	Trp	Ala	Arg	Leu
				245					250					255
Gly	Leu	Gly	Thr	Arg	Glu	Asp	Val	Thr	Pro	Lys	Ser	Asp	Ser	Glu
				260					265					270
Ala	Gln	Gly	Glu	Gly	Glu	Arg	Leu	Phe	Asp	Leu	Glu	Glu	Thr	Leu
				275					280					285
Gly	Arg	His	Arg	Arg	Ile	Leu	Gln	Ala	Leu	Arg	Gly	Leu	Ser	Lys
				290					295					300
Gln	Leu	Ala	Gln	Ala	Glu	Arg	Gln	Trp	Lys	Lys	Gln	Leu	Gly	Pro
				305					310					315

```

Pro Gly Gln Ala Arg Pro Asp Gly Gly Trp Ala Leu Asp Ala Ser
320 325
Cys Gln Ile Pro Ser Thr Pro Gly Arg Gly Gly His Leu Ser Met
335 340
Ser Leu Asn Lys Asp Ser Ala Lys Val Gly Ala Leu Leu His Gly
350 355
Gln Trp Thr Leu Leu Gln Ala Leu Gln Glu Met Arg Asp Ala Ala
365 370
Val Arg Met Ala Ala Glu Ala Gln Val Ser Tyr Leu Pro Val Gly
380 385
Ile Glu His His Phe Leu Glu Leu Asp His Ile Leu Gly Leu Leu
395 400
Gln Glu Glu Leu Arg Gly Pro Ala Lys Ala Ala Ala Lys Ala Pro
410 415
Arg Pro Pro Gly Gln Pro Pro Arg Ala Ser Ser Cys Leu Gln Pro
425 430
Gly Ile Phe Leu Phe Tyr Leu Leu Ile Gln Thr Val Gly Phe Phe
440 445
Gly Tyr Val His Phe Ser Arg Gln Glu Leu Asn Lys Ser Leu Gln
455 460
Glu Cys Leu Ser Thr Gly Ser Leu Pro Leu Gly Pro Ala Pro His
470 475
Thr Pro Arg Ala Leu Gly Ile Leu Arg Arg Gln Pro Leu Pro Ala
485 490
Ser Met Pro Ala

```

<210> 19
 <211> 879
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4946103CD1

```

<400> 19
Met Thr Ala Ile Lys His Ala Leu Gln Arg Asp Ile Phe Thr Pro
1 5 10 15
Asn Asp Glu Arg Leu Leu Ser Ile Val Asn Val Cys Lys Ala Gly
20 25 30
Lys Lys Lys Lys Asn Cys Phe Leu Cys Ala Thr Val Thr Thr Glu
35 40 45
Arg Pro Val Gln Val Lys Val Val Lys Val Lys Lys Ser Asp Lys
50 55 60
Gly Asp Phe Tyr Lys Arg Gln Ile Ala Trp Ala Leu Arg Asp Leu
65 70 75
Ala Val Val Asp Ala Lys Asp Ala Ile Lys Glu Asn Pro Glu Phe
80 85 90
Asp Leu His Phe Glu Lys Ile Tyr Lys Trp Val Ala Ser Ser Thr
95 100 105
Ala Glu Lys Asn Ala Phe Ile Ser Cys Ile Trp Lys Leu Asn Gln
110 115 120
Arg Tyr Leu Arg Lys Lys Ile Asp Phe Val Asn Val Ser Ser Gln
125 130 135
Leu Leu Glu Glu Ser Val Pro Ser Gly Glu Asn Gln Ser Val Thr
140 145 150
Gly Gly Asp Glu Glu Val Val Asp Glu Tyr Gln Glu Leu Asn Ala
155 160 165
Arg Glu Glu Gln Asp Ile Glu Ile Met Met Glu Gly Cys Glu Tyr
170 175 180
Ala Ile Ser Asn Ala Glu Arg Phe Ala Glu Lys Leu Ser Arg Glu
185 190 195
Leu Gln Val Leu Asp Gly Ala Asn Ile Gln Ser Ile Met Ala Ser
200 205 210
Glu Lys Gln Val Asn Ile Leu Met Lys Leu Leu Asp Glu Ala Leu
215 220 225

```

Lys	Glu	Val	Asp	Gln	Ile	Glu	Leu	Lys	Leu	Ser	Ser	Tyr	Glu	Glu
				230					235					240
Met	Leu	Gln	Ser	Val	Lys	Glu	Gln	Met	Asp	Gln	Ile	Ser	Glu	Ser
				245					250					255
Asn	His	Leu	Ile	His	Leu	Ser	Asn	Thr	Asn	Asn	Val	Lys	Leu	Leu
				260					265					270
Ser	Glu	Ile	Glu	Phe	Leu	Val	Asn	His	Met	Asp	Leu	Ala	Lys	Gly
				275					280					285
His	Ile	Lys	Ala	Leu	Gln	Glu	Gly	Asp	Leu	Ala	Ser	Ser	Arg	Gly
				290					295					300
Ile	Glu	Ala	Cys	Thr	Asn	Ala	Ala	Asp	Ala	Leu	Leu	Gln	Cys	Met
				305					310					315
Asn	Val	Ala	Leu	Arg	Pro	Gly	His	Asp	Leu	Leu	Leu	Ala	Val	Lys
				320					325					330
Gln	Gln	Gln	Gln	Arg	Phe	Ser	Asp	Leu	Arg	Glu	Leu	Phe	Ala	Arg
				335					340					345
Arg	Leu	Ala	Ser	His	Leu	Asn	Asn	Val	Phe	Val	Gln	Gln	Gly	His
				350					355					360
Asp	Gln	Ser	Ser	Ser	Leu	Pro	Gln	His	Cys	Val	Ser	Thr	Gly	Phe
				365					370					375
Thr	Gln	Ser	Ser	Ser	Ile	Ser	Gln	Arg	Phe	Pro	Pro	Ile	Ala	Lys
				380					385					390
Leu	Met	Glu	Trp	Leu	Lys	Ser	Thr	Asp	Tyr	Gly	Lys	Tyr	Glu	Gly
				395					400					405
Leu	Thr	Lys	Asn	Tyr	Met	Asp	Tyr	Leu	Ser	Arg	Leu	Tyr	Glu	Arg
				410					415					420
Glu	Ile	Lys	Asp	Phe	Phe	Glu	Val	Ala	Lys	Ile	Lys	Met	Thr	Gly
				425					430					435
Thr	Thr	Lys	Glu	Ser	Lys	Lys	Phe	Gly	Leu	His	Gly	Ser	Ser	Gly
				440					445					450
Lys	Leu	Thr	Gly	Ser	Thr	Ser	Ser	Leu	Asn	Lys	Leu	Ser	Val	Gln
				455					460					465
Ser	Ser	Gly	Asn	Arg	Arg	Ser	Gln	Ser	Ser	Ser	Leu	Leu	Asp	Met
				470					475					480
Gly	Asn	Met	Ser	Ala	Ser	Asp	Leu	Asp	Val	Ala	Asp	Arg	Thr	Lys
				485					490					495
Phe	Asp	Lys	Ile	Phe	Glu	Gln	Val	Leu	Ser	Glu	Leu	Glu	Pro	Leu
				500					505					510
Cys	Leu	Ala	Glu	Gln	Asp	Phe	Ile	Ser	Lys	Phe	Phe	Lys	Leu	Gln
				515					520					525
Gln	His	Gln	Ser	Met	Pro	Gly	Thr	Met	Ala	Glu	Ala	Glu	Asp	Leu
				530					535					540
Asp	Gly	Gly	Thr	Leu	Ser	Arg	Gln	His	Asn	Cys	Gly	Thr	Pro	Leu
				545					550					555
Pro	Val	Ser	Ser	Glu	Lys	Asp	Met	Ile	Arg	Gln	Met	Met	Ile	Lys
				560					565					570
Ile	Phe	Arg	Cys	Ile	Glu	Pro	Glu	Leu	Asn	Asn	Leu	Ile	Ala	Leu
				575					580					585
Gly	Asp	Lys	Ile	Asp	Ser	Phe	Asn	Ser	Leu	Tyr	Met	Leu	Val	Lys
				590					595					600
Met	Ser	His	His	Val	Trp	Thr	Ala	Gln	Asn	Val	Asp	Pro	Ala	Ser
				605					610					615
Phe	Leu	Ser	Thr	Thr	Leu	Gly	Asn	Val	Leu	Val	Thr	Val	Lys	Arg
				620					625					630
Asn	Phe	Asp	Lys	Cys	Ile	Ser	Asn	Gln	Ile	Arg	Gln	Met	Glu	Glu
				635					640					645
Val	Lys	Ile	Ser	Lys	Lys	Ser	Lys	Val	Gly	Ile	Leu	Pro	Phe	Val
				650					655					660
Ala	Glu	Phe	Glu	Glu	Phe	Ala	Gly	Leu	Ala	Glu	Ser	Ile	Phe	Lys
				665					670					675
Asn	Ala	Glu	Arg	Arg	Gly	Asp	Leu	Asp	Lys	Ala	Tyr	Thr	Lys	Leu
				680					685					690
Ile	Arg	Gly	Val	Phe	Val	Asn	Val	Glu	Lys	Val	Ala	Asn	Glu	Ser
				695					700					705
Gln	Lys	Thr	Pro	Arg	Asp	Val	Val	Met	Met	Glu	Asn	Phe	His	His
				710					715					720
Ile	Phe	Ala	Thr	Leu	Ser	Arg	Leu	Lys	Ile	Ser	Cys	Leu	Glu	Ala

Glu Lys Lys Glu	725	Ala Lys Gln Lys Tyr	730	Thr Asp His Leu Gln	735
Tyr Val Ile Tyr	740	Ser Leu Gly Gln Pro	745	Leu Glu Lys Leu Asn	750
Phe Phe Glu Gly	755	Val Glu Ala Arg Val	760	Ala Gln Gly Ile Arg	765
Glu Glu Val Ser	770	Tyr Gln Leu Ala Phe	775	Asn Lys Gln Glu Leu	780
Lys Val Ile Lys	785	Glu Tyr Pro Gly Lys	790	Glu Val Lys Lys Gly	795
Asp Asn Leu Tyr	800	Lys Lys Val Asp Lys	805	His Leu Cys Glu Glu	810
Asn Leu Leu Gln	815	Val Val Trp His Ser	820	Met Gln Asp Glu Phe	825
Arg Gln Tyr Lys	830	His Phe Glu Gly Leu	835	Ile Ala Arg Cys Tyr	840
Gly Ser Gly Val	845	Thr Met Glu Phe Thr	850	Ile Gln Asp Ile Leu	855
Tyr Cys Ser Ser	860	Ile Ala Gln Ser His	865		870
	875				

<210> 20

<211> 298

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5562355CD1

<400> 20

Met Asp Asn Ala	Gly	Lys Glu Arg Glu	Ala	Val Gln Leu Met	Ala
1	5		10		15
Glu Ala Glu Lys	Arg	Val Lys Ala Ser	His	Ser Phe Leu Arg	Gly
	20		25		30
Leu Phe Gly Gly	Asn	Thr Arg Ile Glu	Glu	Ala Cys Glu Met	Tyr
	35		40		45
Thr Arg Ala Ala	Asn	Met Phe Lys Met	Ala	Lys Asn Trp Ser	Ala
	50		55		60
Ala Gly Asn Ala	Phe	Cys Gln Ala Ala	Lys	Leu His Met Gln	Leu
	65		70		75
Gln Ser Lys His	Asp	Ser Ala Thr Ser	Phe	Val Asp Ala Gly	Asn
	80		85		90
Ala Tyr Lys Lys	Ala	Asp Pro Gln Glu	Ala	Ile Asn Cys Leu	Asn
	95		100		105
Ala Ala Ile Asp	Ile	Tyr Thr Asp Met	Gly	Arg Phe Thr Ile	Ala
	110		115		120
Ala Lys His His	Ile	Thr Ile Ala Glu	Ile	Tyr Glu Thr Glu	Leu
	125		130		135
Val Asp Ile Glu	Lys	Ala Ile Ala His	Tyr	Glu Gln Ser Ala	Asp
	140		145		150
Tyr Tyr Lys Gly	Glu	Glu Ser Asn Ser	Ser	Ala Asn Lys Cys	Leu
	155		160		165
Leu Lys Val Ala	Ala	Tyr Ala Ala His	Leu	Glu Gln Tyr Gln	Asn
	170		175		180
Ala Ile Glu Ile	Tyr	Glu Gln Val Gly	Ala	Asn Thr Met Asp	Asn
	185		190		195
Pro Leu Thr Thr	Tyr	Ser Ala Lys Asp	Tyr	Phe Phe Lys Ala	Ala
	200		205		210
Leu Cys His Phe	Ile	Val Asp Glu Leu	Asn	Ala Lys Leu Ala	Leu
	215		220		225
Glu Gln Tyr Glu	Asp	Met Phe Pro Ala	Phe	Thr Asp Ser Arg	Glu
	230		235		240
Cys Lys Leu Leu	Lys	Lys Leu Leu Glu	Ala	His Glu Glu Gln	Asn
	245		250		255
Ser Glu Ala Tyr	Thr	Glu Ala Val Lys	Glu	Phe Asp Ser Ile	Ser

Arg	Leu	Asp	Gln	260	Trp	Leu	Thr	Thr	Met	265	Leu	Leu	Arg	Ile	Lys	270	Lys
				275						280							285
Ser	Ile	Gln	Gly	Asp	Gly	Glu	Gly	Asp	Gly	Asp	Leu	Lys					
				290					295								

<210> 21
 <211> 941
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5678824CD1

<400> 21

Met	Ala	Ala	Tyr	Leu	Gln	Trp	Arg	Arg	Phe	Val	Phe	Phe	Asp	Lys
1				5					10					15
Glu	Leu	Val	Lys	Glu	Pro	Leu	Ser	Asn	Asp	Gly	Ala	Ala	Pro	Gly
				20					25					30
Ala	Thr	Pro	Ala	Ser	Gly	Ser	Ala	Ala	Ser	Lys	Phe	Leu	Cys	Leu
				35					40					45
Pro	Pro	Gly	Ile	Thr	Val	Cys	Asp	Ser	Gly	Arg	Gly	Ser	Leu	Val
				50					55					60
Phe	Gly	Asp	Met	Glu	Gly	Gln	Ile	Trp	Phe	Leu	Pro	Arg	Ser	Leu
				65					70					75
Gln	Leu	Thr	Gly	Phe	Gln	Ala	Tyr	Lys	Leu	Arg	Val	Thr	His	Leu
				80					85					90
Tyr	Gln	Leu	Lys	Gln	His	Asn	Ile	Leu	Ala	Ser	Val	Gly	Glu	Asp
				95					100					105
Glu	Glu	Gly	Ile	Asn	Pro	Leu	Val	Lys	Ile	Trp	Asn	Leu	Glu	Lys
				110					115					120
Arg	Asp	Gly	Gly	Asn	Pro	Leu	Cys	Thr	Arg	Ile	Phe	Pro	Ala	Ile
				125					130					135
Pro	Gly	Thr	Glu	Pro	Thr	Val	Val	Ser	Cys	Leu	Thr	Val	His	Glu
				140					145					150
Asn	Leu	Asn	Phe	Met	Ala	Ile	Gly	Phe	Thr	Asp	Gly	Ser	Val	Thr
				155					160					165
Leu	Asn	Lys	Gly	Asp	Ile	Thr	Arg	Asp	Arg	His	Ser	Lys	Thr	Gln
				170					175					180
Ile	Leu	His	Lys	Gly	Asn	Tyr	Pro	Val	Thr	Gly	Leu	Ala	Phe	Arg
				185					190					195
Gln	Ala	Gly	Lys	Thr	Thr	His	Leu	Phe	Val	Val	Thr	Thr	Glu	Asn
				200					205					210
Val	Gln	Ser	Tyr	Ile	Val	Ser	Gly	Lys	Asp	Tyr	Pro	Arg	Val	Glu
				215					220					225
Leu	Asp	Thr	His	Gly	Cys	Gly	Leu	Arg	Cys	Ser	Ala	Leu	Ser	Asp
				230					235					240
Pro	Ser	Gln	Asp	Leu	Gln	Phe	Ile	Val	Ala	Gly	Asp	Glu	Cys	Val
				245					250					255
Tyr	Leu	Tyr	Gln	Pro	Asp	Glu	Arg	Gly	Pro	Cys	Phe	Ala	Phe	Glu
				260					265					270
Gly	His	Lys	Leu	Ile	Ala	His	Trp	Phe	Arg	Gly	Tyr	Leu	Ile	Ile
				275					280					285
Val	Ser	Arg	Asp	Arg	Lys	Val	Ser	Pro	Lys	Ser	Glu	Phe	Thr	Ser
				290					295					300
Arg	Asp	Ser	Gln	Ser	Ser	Asp	Lys	Gln	Ile	Leu	Asn	Ile	Tyr	Asp
				305					310					315
Leu	Cys	Asn	Lys	Phe	Ile	Ala	Tyr	Ser	Thr	Val	Phe	Glu	Asp	Val
				320					325					330
Val	Asp	Val	Leu	Ala	Glu	Trp	Gly	Ser	Leu	Tyr	Val	Leu	Thr	Arg
				335					340					345
Asp	Gly	Arg	Val	His	Ala	Leu	Gln	Glu	Lys	Asp	Thr	Gln	Thr	Lys
				350					355					360
Leu	Glu	Met	Leu	Phe	Lys	Lys	Asn	Leu	Phe	Glu	Met	Ala	Ile	Asn
				365					370					375
Leu	Ala	Lys	Ser	Gln	His	Leu	Asp	Ser	Asp	Gly	Leu	Ala	Gln	Ile

	380		385		390
Phe Met Gln Tyr	Gly Asp His Leu Tyr	Ser Lys Gly Asn His	Asp		
	395		400		405
Gly Ala Val Gln	Gln Tyr Ile Arg Thr	Ile Gly Lys Leu Glu	Pro		
	410		415		420
Ser Tyr Val Ile	Arg Lys Phe Leu Asp	Ala Gln Arg Ile His	Asn		
	425		430		435
Leu Thr Ala Tyr	Leu Gln Thr Leu His	Arg Gln Ser Leu Ala	Asn		
	440		445		450
Ala Asp His Thr	Thr Leu Leu Leu Asn	Cys Tyr Thr Lys Leu	Lys		
	455		460		465
Asp Ser Ser Lys	Leu Glu Glu Phe Ile	Lys Lys Lys Ser Glu	Ser		
	470		475		480
Glu Val His Phe	Asp Val Glu Thr Ala	Ile Lys Val Leu Arg	Gln		
	485		490		495
Ala Gly Tyr Tyr	Ser His Ala Leu Tyr	Leu Ala Glu Asn His	Ala		
	500		505		510
His His Glu Trp	Tyr Leu Lys Ile Gln	Leu Glu Asp Ile Lys	Asn		
	515		520		525
Tyr Gln Glu Ala	Leu Arg Tyr Ile Gly	Lys Leu Pro Phe Glu	Gln		
	530		535		540
Ala Glu Ser Asn	Met Lys Arg Tyr Gly	Lys Ile Leu Met His	His		
	545		550		555
Ile Pro Glu Gln	Thr Thr Gln Leu Leu	Lys Gly Leu Cys Thr	Asp		
	560		565		570
Tyr Arg Pro Ser	Leu Glu Gly Arg Ser	Asp Arg Glu Ala Pro	Gly		
	575		580		585
Cys Arg Ala Asn	Ser Glu Glu Phe Ile	Pro Ile Phe Ala Asn	Asn		
	590		595		600
Pro Arg Glu Leu	Lys Ala Phe Leu Glu	His Met Ser Glu Val	Gln		
	605		610		615
Pro Asp Ser Pro	Gln Gly Ile Tyr Asp	Thr Leu Leu Glu Leu	Arg		
	620		625		630
Leu Gln Asn Trp	Ala His Glu Lys Asp	Pro Gln Val Lys Glu	Lys		
	635		640		645
Leu His Ala Glu	Ala Ile Ser Leu Leu	Lys Ser Gly Arg Phe	Cys		
	650		655		660
Asp Val Phe Asp	Lys Ala Leu Val Leu	Cys Gln Met His Asp	Phe		
	665		670		675
Gln Asp Gly Val	Leu Tyr Leu Tyr Glu	Gln Gly Lys Leu Phe	Gln		
	680		685		690
Gln Ile Met His	Tyr His Met Gln His	Glu Gln Tyr Arg Gln	Val		
	695		700		705
Ile Ser Val Cys	Glu Arg His Gly Glu	Gln Asp Pro Ser Leu	Trp		
	710		715		720
Glu Gln Ala Leu	Ser Tyr Phe Ala Arg	Lys Glu Glu Asp Cys	Lys		
	725		730		735
Glu Tyr Val Ala	Ala Val Leu Lys His	Ile Glu Asn Lys Asn	Leu		
	740		745		750
Met Pro Pro Leu	Leu Val Val Gln Thr	Leu Ala His Asn Ser	Thr		
	755		760		765
Ala Thr Leu Ser	Val Ile Arg Asp Tyr	Leu Val Gln Lys Leu	Gln		
	770		775		780
Lys Gln Ser Gln	Gln Ile Ala Gln Asp	Glu Leu Arg Val Arg	Arg		
	785		790		795
Tyr Arg Glu Glu	Thr Thr Arg Ile Arg	Gln Glu Ile Gln Glu	Leu		
	800		805		810
Lys Ala Ser Pro	Lys Ile Phe Gln Lys	Thr Lys Cys Ser Ile	Cys		
	815		820		825
Asn Ser Ala Leu	Glu Leu Pro Ser Val	His Phe Leu Cys Gly	His		
	830		835		840
Ser Phe His Gln	His Cys Phe Glu Ser	Tyr Ser Glu Ser Asp	Ala		
	845		850		855
Asp Cys Pro Thr	Cys Leu Pro Glu Asn	Arg Lys Val Met Asp	Met		
	860		865		870
Ile Arg Ala Gln	Glu Gln Lys Arg Asp	Leu His Asp Gln Phe	Gln		
	875		880		885

His	Gln	Leu	Arg	Cys	Ser	Asn	Asp	Ser	Phe	Ser	Val	Ile	Ala	Asp	
				890					895					900	
Tyr	Phe	Gly	Arg	Gly	Val	Phe	Asn	Lys	Leu	Thr	Leu	Leu	Thr	Asp	
				905					910					915	
Pro	Pro	Thr	Ala	Arg	Leu	Thr	Ser	Ser	Leu	Glu	Ala	Gly	Leu	Gln	
				920					925					930	
Arg	Asp	Leu	Leu	Met	His	Ser	Arg	Arg	Gly	Thr					
				935					940						

<210> 22

<211> 336

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5870962CD1

<400> 22

Met	Ser	Lys	Ser	Val	Pro	Ala	Phe	Leu	Gln	Asp	Glu	Val	Ser	Gly	
1				5					10					15	
Ser	Val	Met	Ser	Val	Tyr	Ser	Gly	Asp	Phe	Gly	Asn	Leu	Glu	Val	
				20					25					30	
Lys	Gly	Asn	Ile	Gln	Phe	Ala	Ile	Glu	Tyr	Val	Glu	Ser	Leu	Lys	
				35					40					45	
Glu	Leu	His	Val	Phe	Val	Ala	Gln	Cys	Lys	Asp	Leu	Ala	Ala	Ala	
				50					55					60	
Asp	Val	Lys	Lys	Gln	Arg	Ser	Asp	Pro	Tyr	Val	Lys	Ala	Tyr	Leu	
				65					70					75	
Leu	Pro	Asp	Lys	Gly	Lys	Met	Gly	Lys	Lys	Lys	Thr	Leu	Val	Val	
				80					85					90	
Lys	Lys	Thr	Leu	Asn	Pro	Val	Tyr	Asn	Glu	Ile	Leu	Arg	Tyr	Lys	
				95					100					105	
Ile	Glu	Lys	Gln	Ile	Leu	Lys	Thr	Gln	Lys	Leu	Asn	Leu	Ser	Ile	
				110					115					120	
Trp	His	Arg	Asp	Thr	Phe	Lys	Arg	Asn	Ser	Phe	Leu	Gly	Glu	Val	
				125					130					135	
Glu	Leu	Asp	Leu	Glu	Thr	Trp	Asp	Trp	Asp	Asn	Lys	Gln	Asn	Lys	
				140					145					150	
Gln	Leu	Arg	Trp	Tyr	Pro	Leu	Lys	Arg	Lys	Thr	Ala	Pro	Val	Ala	
				155					160					165	
Leu	Glu	Ala	Glu	Asn	Arg	Gly	Glu	Met	Lys	Leu	Ala	Leu	Gln	Tyr	
				170					175					180	
Val	Pro	Glu	Pro	Val	Pro	Gly	Lys	Lys	Leu	Pro	Thr	Thr	Gly	Glu	
				185					190					195	
Val	His	Ile	Trp	Val	Lys	Glu	Cys	Leu	Asp	Leu	Pro	Leu	Leu	Arg	
				200					205					210	
Gly	Ser	His	Leu	Asn	Ser	Phe	Val	Lys	Cys	Thr	Ile	Leu	Pro	Asp	
				215					220					225	
Thr	Ser	Arg	Lys	Ser	Arg	Gln	Lys	Thr	Arg	Ala	Val	Gly	Lys	Thr	
				230					235					240	
Thr	Asn	Pro	Ile	Phe	Asn	His	Thr	Met	Val	Tyr	Asp	Gly	Phe	Arg	
				245					250					255	
Pro	Glu	Asp	Leu	Met	Glu	Ala	Cys	Val	Glu	Leu	Thr	Val	Trp	Asp	
				260					265					270	
His	Tyr	Lys	Leu	Thr	Asn	Gln	Phe	Leu	Gly	Gly	Leu	Arg	Ile	Gly	
				275					280					285	
Phe	Gly	Thr	Gly	Lys	Ser	Tyr	Gly	Thr	Glu	Val	Asp	Trp	Met	Asp	
				290					295					300	
Ser	Thr	Ser	Glu	Glu	Val	Ala	Leu	Trp	Glu	Lys	Met	Val	Asn	Ser	
				305					310					315	
Pro	Asn	Thr	Trp	Ile	Glu	Ala	Thr	Leu	Pro	Leu	Arg	Met	Leu	Leu	
				320					325					330	
Ile	Ala	Lys	Ile	Ser	Lys										
				335											

<210> 23

<211> 163
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2818605CD1

<400> 23
 Met Asp Asp Lys Glu Pro Lys Arg Trp Pro Thr Leu Arg Asp Arg
 1 5 10 15
 Leu Cys Ser Asp Gly Phe Leu Phe Pro Gln Tyr Pro Ile Lys Pro
 20 25 30
 Tyr His Leu Lys Gly Ile His Arg Ala Val Phe Tyr Arg Asp Leu
 35 40 45
 Glu Glu Leu Lys Phe Val Leu Leu Thr Arg Tyr Asp Ile Asn Lys
 50 55 60
 Arg Asp Arg Lys Glu Arg Thr Ala Leu His Leu Ala Cys Ala Thr
 65 70 75
 Gly Gln Pro Glu Met Val His Leu Leu Val Ser Arg Arg Cys Glu
 80 85 90
 Leu Asn Leu Cys Asp Arg Glu Asp Arg Thr Pro Leu Ile Lys Ala
 95 100 105
 Val Gln Leu Arg Gln Glu Ala Cys Ala Thr Leu Leu Leu Gln Asn
 110 115 120
 Gly Ala Asp Pro Asn Ile Thr Asp Val Phe Gly Arg Thr Ala Leu
 125 130 135
 His Tyr Ala Val Tyr Asn Glu Asp Thr Ser Met Ile Glu Lys Leu
 140 145 150
 Leu Ser His Gly Thr Asn Ile Glu Glu Cys Ser Lys Val
 155 160

<210> 24
 <211> 1983
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 381039CB1

<400> 24
 gtcaggggag ccgggcgtgc ggagggcggt gatgagggcg gaagggcgga gtggttgaag 60
 ggggtgattgc tgactagcgg ggagggagtg ggcagcgatg gtgatggcgg cgaagaaggg 120
 cccgggcccgg ggccggcgggg tcagcggggg caagggcgag gccggagcgg cctcggaggt 180
 gtggtgcccgt cgggtgcggg agctgggtgg ctgcagccag gccgggaacc gccactgctt 240
 cgagtgcgcc cagcgcgggg tcacctacgt gcatatcacc gtgggcagct tcgtgtgcac 300
 cacctgctcc ggccctctga gagggctgaa ccccccctcat cgtgtcaagt caatctccat 360
 gacaactttc actgagcctg aagtagtatt cctgcaatcc cgtggaaatg aggtttgccg 420
 gaagatttgg ttgggtctgt ttgatgctcg gacatcttta gtaccagatt ccagggatcc 480
 tcagaaagtg aaggagtttc tccaggaaaa atatgagaag aagagatggt atgtccccc 540
 agaccaagtc aaggggcccc cttataccaa aggcagtgcc tccaccctg tgcagggtc 600
 catcccagaa gggaagcccc ttgggacact tctgggtgat cctgcaccgt ctctctcagt 660
 tgctgcctcc acctcgagcc agcccgtcag tcagtctcac gctcggacat cccaggcccc 720
 gagcactcag ccacctcccc actcctctgt caaaaaagcc agtactgacc tgctggctga 780
 catcgggtgga gaccctttt ctgcacccca gatggcacca gcttttgctg cattccctgc 840
 ctttgggggc cagacacctt cccaaggagg ctttgccaac tttgatgcct ttagcagtgg 900
 cccagctctt tctgtgtttg gaagcctccc tccagctggt caagcctcgt tccaggcccc 960
 gccaaactct gcaggagca gccaggggac tccatttggt gccactcccc tggcaccgcc 1020
 cagtcagcca aacagcctcg cagacgtggg cagcttctct ggaccggggg tgcccgtctg 1080
 aggtgttctt agcagcctct tcgggatggc tggccaggte cccccgctcc agtctgtcac 1140
 gatggggggc ggccggcgga gcagcacagg gctggccttt ggagccttca ctaacccttt 1200
 cacagctccc gccgcccagt ccccgctgcc ttccaccaac ecgttccagc ccaatggctt 1260
 ggcgccaggg cccggttttg ggatgagcag tgctgggcct ggcttcccc aggcagtgcc 1320
 acccactggg gcctttgcca gctccttccc agcaccgctg ttccccccgc agaccctgct 1380
 tgttcagcag cagaatggct cttccttogg ggacttagga tcagccaagt tggggcagag 1440
 gccactgagc cagccagctg ggatctccac caacccttc atgactggac cctcatcaag 1500

```

cccattcgcc tccaaacctc caaccaccaa ccccttcttg tagcactgtg tttttggggg 1560
gcctcttccc tgccctctgg ggccctcttg ctccctagag ctctgggtgac cacttgccctg 1620
tgggcatttc tatgggcctt ggggatgggt gaggtgctaa tgctttgctt ggggcctaca 1680
ggtgaaaggt ggctgccctc agattccaca aagcctctct cccctccctc gcatcacccc 1740
caccacaggca ggaagcccag gaggagtgcg ggcagggcct gacctggagg agtgatgggt 1800
gagggcgagg gatttttttc acatgatcag tccctgtgtg aacagctctt ccctgggcct 1860
agctcgccag cgctgtgctc ctcatggacg ggagcgcag gggaaagtag gggaaagatg 1920
aggcacagtg ttgatggcgc agtgaccaga cagttctgag cccagaagag ccatgtgtgc 1980
caa

```

<210> 25

<211> 1559

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 383249CB1

<400> 25

```

cccagcgaga gatctgcagc taggetggct gcacttgctc caggggtcag gggatcggag 60
ggggattgaa gaatgcgcca ttaaaaggaa agatcaagga gtaaaccaga agaagaagaa 120
aaagaggact tcaaagctgg gaaggatgag ttcttgagc aacgtctgtg ggtccaggca 180
ggcacaggct gcagctgagg gtggttacca gcgctatgga gtccggctct acctgcacca 240
gttttatgag gactgtacag cctcaatttg ggagtatgag gatgatttcc agatccaaag 300
atcacctaac aggttgagct cagtattctg gaaggttggg ctcatctcag gtacagtttt 360
tgtgatccct ggattgactg ttctggcagt gggctttctt gtgcccccca aaatcgaagc 420
atttggcgaa gccgattttg tgggtgtcga cacacatgct gtccagttta acagtgtctc 480
ggacatgtac aagctggcag gagctgttct cttctgcatt ggaggcacgt ccatggcagg 540
gtgcctgctg atgtcgggtg ttgtaaagag ctactccaaa gaagaaaaat tcctccagca 600
gaagttaaaa gaacgaatcg cagacatcaa agcccacacc cagccgggta caaaagctcc 660
agggccaggg gaaacaaaga ttccagtcac ttgtccagg gttcaaaatg tccagcctct 720
actggcaacc tgaaaccttt cccaccccag ttcttctgt ctgatttatg cccgtgggta 780
aaaagagcag gccagttttc gagataaaga agatttggcg ttgactgcc tagggctgtg 840
ttcagctgtg ggcaatataa tgggtggact cacacttget cagttcaggc agctctgctg 900
gagggcggtg ccatgcctaa gcctgtgcac atgcatccag ctgcttaaga tgggtgcttc 960
cttgtaccgg gccaccagaa aacccttga actcctctt catagatcgt gactttgtgt 1020
tattttccgt gttgtttgaa agtgccgaag agtttagacc agctcaccaa tggctaattg 1080
ggtccatttt attttctaat attggtttat ctaatgtttt ctgcgtgggtg ctgtcttcca 1140
ttcctgtctc actatgtgta agattttgtc atatgtgttc atagaaacat ggaattctct 1200
gattttgagc cgaacatgag tttttagact gtatcttgac atgaggttcc tgacattgga 1260
tacaaagtta tcagcattca caaatctttt tgttttctcc atttttttcc cctttgagat 1320
gaagtgtgtg tcttttgcct aggttggagt acagcgggtg aatcatagcg cactgacgcc 1380
ttgaactcct gggctcaagc gatcctccca cctcagcctc ccaagtagct gggactacag 1440
gcacctgccca ccaccccag ttaatatatt ttttaatttt ttttagagggc aggatcttgc 1500
tgtccaattt tttttaaaaa gctaaaattg ccaactttag aatcttgacc aaaaaaaaa 1559

```

<210> 26

<211> 1643

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 618769CB1

<400> 26

```

gaaatcatgc ccctcgtaga gcagcaggtc caagcagggc tgctggctat ttttccaaaa 60
agtgaggcag ttttaaaaaa agggcgagaa ctagaattat agaataatgg cacattttgt 120
gtattttgaa aactaacggc ttgcatgggt cacaacccat ttcttatgcc tgtgttttcc 180
ttggcagcaa aatttctgtg gttcctctca ctccacctcc tgtcagcgag agccagtga 240
gccgcagtc ttggcaggag gtcagtgca cagatattct gaaacctctc aatcaagagg 300
atcccaaatg ctctactaac cctattttga agcaacagaa tctcccatcc agtccggcac 360
ccagtaccat attctctgga ggttttagac acggaagttt aattagcatt gacagcacct 420
gtacagagat gggcaatttt gacaatgcta atgtcactgg agaaatagaa tttgccattc 480
attattgctt caaaacccat tctttagaaa tatgcatcaa ggctgtgaag aaccttgcc 540
atggagaaga aaagaagaaa aagtgcaatc cgtatgtgaa gacctacctg ttgcccagca 600

```

```

gatcctccca gggaaagcgc aagactggag tccaaaggaa caccgtggac ccgaccttcc 660
aggagacctt gaagtatcag gtggcccctg cccagctggg gaccggcag ctgcaggctc 720
cgggtgtggca tctgggcacg ctggcccggg gagggtttct tggagaagtg atcattcctc 780
tgccacagtg ggactttgaa gacagcacia cacagtcctt ccgctggcat ccgctccggg 840
ccaaggcgga gaaatacgaa gacagcgctc ctcagagtaa tggagagctc acagtcgggg 900
ctaagctggg tctcccttca cggcccagaa aactccaaga ggctcaagaa gggacagatc 960
agccatcact tcatgggtcaa ctttgtttgg tagtgctagg agccaagaat ttacctgtgc 1020
ggccagatgg caccttgaac tcatttggtt agggctgtct cactctgcca gaccaacaaa 1080
aactgagact gaagtgcgca gtccctgagg agcaggcttg cccccagtg aaacactcat 1140
ttgtcttcag tggcgtaacc ccagctcagc tgaggcagtc gagcttggag ttaactgtct 1200
gggatcaggc cctcttttga atgaatgacc gcttgcttgg aggaaccaga cttgggtcaa 1260
agggagacac agctgttggc ggggatgcat gctcactatc gaagctccag tggcagaaag 1320
tcctttccag cccaatcta tggacagaca tgactcttgt cctgactga catgaaggcc 1380
tcaaggttcc aggttgcagc aggcgtgagg cactgtgcgt ctgcagaggg gctacgaacc 1440
aggtgcaggg tcccagctgg agaccccttt gaccttgagc agtctccatc tgcggccctg 1500
tcccatggct taaccgccta ttggtatctg tgtatattta cgttaaacac aattatgtta 1560
cctaagcctc tgggtgggtta tctcctcttt gagatgtaga aaatggccag attttaataa 1620
acgttggttac ccaaaaaaaaa aaa 1643

```

<210> 27

<211> 1613

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1234837CB1

<400> 27

```

cccgggccag gggcgggcgc cgccatgggt aacctgttgc gccgcaagaa gcagagccgc 60
gtcacggagc aggacaaggc catcctgcaa ctgaagcagc agcgggacaa gctgaggcag 120
taccagaaga ggatcgccca gcagctggag cgcgagcgcg ccctggcccg gcagctgctg 180
cgggacggca ggaaggaacg ggccaagctg ctgctcaaga agaagcgata ccaggagcag 240
ctcctggaca ggacggagaa ccagatcagc agcctggagg ccattggttca gaggattgag 300
ttcaccacaga tcgaaatgaa agtgatggag gggctgcagt ttggaaatga gtgtctgaac 360
aagatgcacc aggtgatgtc cattgaagag gtggagagga tcttgacga gacgaggag 420
gccgtggagt accagcgcca aatagacagc ctctggcag gaagcttcac tcaggaggat 480
gaagacgcca tcttgaggga gctgagcgca atcactcagg aacaaataga gctgccagag 540
gttccctccg agccccttcc tgagaagatc ccagaaaacg tccctgtcaa ggccaggccc 600
aggcaggcgg agctgggtggc agcttcgtaa cgtggcctcg tcttgtggga ctacgggga 660
tgccccaggg actgtggccc acagagagtt tgggtcacgg ccagcccctg accgggttcc 720
ctggagccca gtgcgcacgg tgctgagcag agctgcagcc acgcaggcgc attgcaggag 780
gactccagag cgtctcctgg agaccttgag cctgaacgca ctcaggcgcc actggcctgc 840
tctcagtcgg gattaactct cgaccgagcc cagcttctgc cggttgtggg ctcccccggt 900
ggccgaggcc caggcccacg gcctctgggt ctgttccct gcagtcccag ccccgctgg 960
ctcgcgctcg tctgtaggga agacacctcc agaccttggg gtcccgcgc tctctcttgc 1020
tctctcgtgc tccattagc tgggtcaggc ttccgttaag gggctccctc cttggcctgg 1080
cttcccggcg cacctcagct tccctgctgg tggggggatc cccaggagac cagcatgtgc 1140
tgaacctctc tgtgcctctg cctccgcacc ctagacaccc acctccagtt tgaaggtggc 1200
gggcccaggg ctttcttgtc gaattgacga ctccgagagc cctgactccc gccttgccac 1260
tcacggctct gtccactagg gctcagcct gctgagaaag gacctccgat gcttgggaga 1320
cgctgctccg gcaggtgcag ccccgggaag ttgtccatgg gggctcccgc ggttggggct 1380
catggaactg cgagaccggg gacctcctg ccctgcgggt ccccgagcca ccagcagcca 1440
ggactggagg ctgtggggca tggcgtgact tctcgtgcac agggctggtt tgggttatgag 1500
acgatctcgc tgggaccgcc cctgcccgtg gaaagccaca aggacaaagg gacggccac 1560
ctcgaccccc agacaggcgg gcccggtatta aagttggccc tgcacgcca aaa 1613

```

<210> 28

<211> 2810

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1607223CB1

<400> 28

tacagggttgg	gcatttgetgg	gcactgtatg	taggtcgccct	gaccctatct	cccatgggag	60
tgatgaccag	tcccagaggg	ctgtgtgact	ctgaatcttc	caacctcaca	ggccattccc	120
tcccttctga	aagcccaccc	atgggcttct	cgccctgtgc	caaggcggtg	gagggaggcc	180
acggcctctg	atgcacctgg	gcgtcccttc	ccatccccc	tccctgcaca	ggtgctggaa	240
acatgcatga	agagctgcgg	caagcgggtc	cacgacgaag	tgggcaagtt	ccgctttctc	300
aacgagctca	tcaaggctgt	gtctcccaag	tatctgggct	ctcggacatc	ggagaagggtg	360
aagaacaaga	tcttggagct	cctctacagc	tggacagtgg	gcctgcccga	ggaggtgaaa	420
atcgagaggg	cctaccagat	gctaaagaag	caggggattg	taaagtcoga	ccccaaagctt	480
ccagatgaca	ctacctttcc	ccttctctct	ccacggccga	agaatgtgat	ctttgaagat	540
gaggagaaat	ccaagatgct	ggcccgccctg	ctgaagagct	cccatcccga	agacctccgc	600
gcagccaata	agctcatcaa	agagatgggtg	caggaggacc	agaagcggat	ggagaagatc	660
togaagaggg	tgaatgccat	cgaggagggtg	aacaacaatg	tgaactgct	cacggagatg	720
gtgatgagcc	acagccaggg	cggcgacga	gctggcagca	gcgaggacct	catgaaggaa	780
ctgtaccagc	gctgtgagcg	gatgcggccc	acgctcttcc	gactggcgag	tgacacagag	840
gacaatgatg	aggccttagc	ggagatcctg	caggccaatg	acaacctcac	ccaggtgatc	900
aacctgtata	agcagctggg	gcgggggtgag	gaggtcaacg	gtgatgccac	agccggctcc	960
atccctggga	gcacctcggc	cctgtctggat	ctctcaggcc	tggatctccc	gcctgcgggc	1020
accacctacc	cagctatgcc	caccgcctct	ggcgagcagg	ccagccctga	gcagccaggt	1080
gcctcagttt	ccctgcttga	cgacgagctc	atgtctctgg	gcctcagtg	ccccacaccc	1140
ccttcaggcc	caagcctgga	tggtagccga	tggacagct	tccagtcgtc	ggatgccact	1200
gagcccccag	cccctgctct	ggcccaggcc	cccagtatgg	aaagccgacc	cccagcgag	1260
acatccctgc	cagcaagcag	cggctctggac	gacctagacc	tccctggggaa	gacctctctg	1320
cagcagctgc	tgcccccgga	atcccagcaa	gtgcgggtgg	agaagcagca	gacacacccc	1380
cggctcacac	tccgggacct	gcagaataag	agcagcagct	gcagctcccc	cagctccagc	1440
gccaccagcc	ttctccacac	cgtgtcccca	gagcccccca	ggcctccgca	gcagcccgt	1500
ccaaccgagc	tctcactggc	cagcatcact	gtgccccctg	agtccatcaa	accagcaac	1560
atcctgcccg	tgactgtgta	tgaccagcac	ggcttccgca	tctcttcca	ttttgcccgg	1620
gacctcagtc	cgagcgctc	cgagctgtcg	gtggtgtgg	tttccatgct	gagcacccgc	1680
ccccagccca	tccgcaacat	cgtgttccag	tcagctgtcc	ccaaggttat	gaagggtgaag	1740
ctgcagccac	cctcgggcac	ggagctgcca	gcttttaacc	ccatcgtcca	cccctcagca	1800
atcaccaggg	tctgtctgct	tgccaacccc	cagaaggaga	aggttcgcc	ccgctacaag	1860
ctcaccctta	ccatgggtga	ccagacctac	aacgagatgg	gggatgtgga	ccagttcccc	1920
ccacctgaaa	cctggggtag	cctctagaac	agaggggctg	gggagaggaa	ggggcagagg	1980
gaccggtcac	tgtccagcct	ggagggaggc	attggtggcc	aaggacaccc	tttgttgccc	2040
atggccattc	accccaggg	ctggtgcttc	tccccacacc	cctgtaggcc	tcaagtgact	2100
cttccccctc	ctgctccggc	cccgcctctg	ctgagccaaa	cccagtagga	ggctgggct	2160
gggtttgtgc	cgtctggggtc	tccatcaccc	ggacctggag	agggaggggc	tgtgtagcct	2220
tgggaagaact	tgggtcatgg	ggagggaagc	cagctgttgg	ggaaggggcca	ggacctcagg	2280
cccagcccca	accccagctg	gggtgggggtc	ttccccacct	gtctcttatg	ccttatggga	2340
aggcccagcc	ataactcggg	ggccatgctg	gagctgggga	ccagcttagg	cctcctccat	2400
aggaacccag	tgactggggg	gtgacgccta	cacccccagc	tatttgact	ctggtgtgtg	2460
gtttgactgc	gctttcttc	cggattggcc	ctgtgtgac	agcctcaggg	ggcagggtg	2520
ggggaacctc	acctggcccg	tactcctggg	gggtttccct	tgccattggg	ccccctagg	2580
gactgtgggg	gctcaagggt	aatgccagag	gcccattggc	ccagcgaggg	gctgtggggc	2640
acctagagtt	ctcgggtgtg	ctccttcatt	cattggcctc	tgctggggcc	tctatgggt	2700
gtcttacgtc	tgtccatcca	tctgtccgtg	gtcagaagtg	gggtcagttg	gtgagtgaga	2760
gcaggagtat	ttatgaaaat	aaaacgtcgt	tttctctgga	aaaaaaaaaa		2810

<210> 29

<211> 2321

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1621554CB1

<400> 29

tcccaagtga	gtcagagggg	gatcccgact	ccagtcgagg	gccttggcca	gcggagccgc	60
ggtattcgga	agcgggaatc	ccactcagag	cccgggcctg	taggggagg	gcgtcccg	120
cacccggtg	tggggcgctc	cccgctcgtg	acgggggcac	cggcgactca	cccggaagga	180
gaagccgtga	tctggctata	tggtagggcg	cggcggtgtg	cgtgtgggg	agctgggtg	240
gtttctcagat	gtttccttcc	aatggggctt	tgggtgtagga	tgtcggagaa	ccaagaacag	300
gaggaggtga	ttacagtgcg	tgttcaggac	ccccgagtgc	agaatgaggg	ctcctggaac	360
tcttatgtgg	attataagat	attcctccat	accaacagca	aagcctttac	tgccaagact	420
tctgtgtg	ggcgccgcta	ccgtgagttc	gtgtggctga	gaaagcagct	acagagaaat	480
gctgggttgg	tgccgtgtcc	tgaacttcc	gggaagtcaa	ccttcttcgg	cacctcagat	540

gagtttcattg	agaagcgacg	acaagggtctg	cagcacttec	ttgaaaaggt	cctgcagagt	600
gtggtttctcc	tgtagacacg	ccagttgcac	ctattcctgc	aaagccagct	ctcgggtgcct	660
gagatagaag	cctgtgtcca	gggccgaagt	accatgactg	tgtctgatgc	cattcttcga	720
tatgctatgt	caaactgtgg	ctgggcccag	gaagagaggc	agagctcttc	tcacctggct	780
aaaggagacc	agcctaagag	ttgctgcttt	cttccaagat	cgggtaggag	gagctctccc	840
tcaccgcctc	ccagtgaaga	aaaggaccat	ttagaagtgt	gggctccagt	tgttgactct	900
gaggttcctt	ccttggaag	tcccactctc	ccacccctct	cctcaccatt	atgctgtgat	960
tttggaagac	ccaaagaggg	aacctccact	cttcagtctg	tgaggaggggc	tgtgggagga	1020
gatcatgctg	tgccttttga	ccctgggtcag	ttagaaacag	ttttggaaaa	gtgagctctc	1080
ggtttctgctc	tgagatggtc	agagaagatg	cgggccagga	gacttactca	ggtgggactg	1140
ggcacaggggc	aggtatgtgg	gaggctgggc	tgcttagtgt	cttctagtca	cctctgcttg	1200
ggctgattga	cagaggtcag	tcattacagc	cccttatgcc	tcttccatgg	gaacaaatac	1260
tgtgcagatg	tttgtaagtt	aaacataaga	cacaggggct	gttgcttttg	aacagaaccc	1320
tatattactc	tcctgggac	tgagttttctg	caggtcattt	gtatgtagga	ccaggagtat	1380
ctcctcaggt	gaccagtttt	ggggaccctg	atgtggcaaa	ttctaagctg	ccatattgaa	1440
catcatccca	ctgggagtg	ttatgttgta	tcccactctt	ggctggcttc	agtttttgct	1500
gtagccctga	agcactttgt	ttgtgggagg	ctggcctctt	gcctaactcc	ttgcatggac	1560
aggggatga	atatttactt	tcccactctc	ttgcttttct	tttactgat	accactgaat	1620
ggaactgggtg	ctgtgactcc	tgctgctggg	gatttatgtc	ccgagacctt	agcctggctg	1680
agtggagcct	gagacctgca	caacagctca	tggtcatgca	tgagagagaa	gtggctggcc	1740
acagccagag	ggaacagtaa	cagcccagg	gcctttatatt	tgggaaaggc	tgtccggggc	1800
tgttactgtc	tcttctgggt	ataaagcaga	catgtggcca	tcttttccgc	aggggttagg	1860
tgggtcctct	tcttttttga	atccttttct	tctccttttg	tagcagctcc	ctggctccag	1920
ggcttccgcc	accagcgtct	ctgctgtgtt	gogcagtgca	gtggggtgca	agggctttgt	1980
ttctgcctgc	ctgaaagaga	gggctctggg	gatggagatg	agaaacaaca	cgctctcctt	2040
cagacaatga	ggcattctgt	cctcctgctg	ccattcttca	tctccactga	gagccagagc	2100
tggtaggagc	cgagtgcac	aggcattctg	ctctgtctca	ctcttaggtt	tgtgtgtgtg	2160
atccttcccc	tccctgtcgc	ccactcctcc	ctcctctggc	tatcctaccc	tgtctgtggg	2220
ctcttttact	accagcctat	gctgtgggac	tgcatggca	tttagttcag	agtggagggg	2280
ctttggcctg	aaataaaatg	caagtattta	aaaaaaaaa	a		2321

<210> 30

<211> 3581

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1751553CB1

<400> 30

gcgccgcgcg	ggcgccgag	cgggcggttg	gacgtggaga	cccgagagag	gcgccgggac	60
ccggcgccgg	aaaggcgcg	cgtggccttg	ctcaggtgog	cttctccac	ctggcagctc	120
gctcagggct	gtggggggcg	cctgtgaggg	cgccgcgcgt	tgcggcgtgg	gagattaata	180
ttaagatttg	aagtttgtgt	cttttctgtg	atattggaaa	ttgaatgtaa	tggcaacaga	240
atttataaag	agttgtctgt	gaggatgttt	ctatgggtga	acagaaaaac	acaacttttc	300
tgtggaaaga	gattttaaag	cagcagctcc	aaatagtcaa	aatgctacta	tctctgtacc	360
tccattgact	tctgtttctg	taaagcctca	gcttggtgtg	actgaggatt	atttgccttc	420
caaattacca	tctgatggca	aagaagtacc	atgtgtgggtg	cccaagttaa	agttatctta	480
cattcaaccc	aggacacaag	aaactccttc	acatctggaa	gaacttgaag	gatctgccag	540
agcatctttt	ggagatcgaa	aggtagaact	ttccagtcca	tcccagcacg	gacctagcta	600
tgatgtgtat	aaaccattct	atatgtatca	gcacatttca	cctgatttga	gtcgacgctt	660
tcttccccgt	tcagaagtga	cgagactgta	tggatcggtt	tgtgatttaa	ggacgaacaa	720
acttccccgt	tcccctgggc	taagcaaatc	tatgtttgat	cttacaact	catctcagcg	780
attcatccag	agacatgatt	cattgtccag	tgtaccagct	agttcttctt	caaggaaaaa	840
ttctcagggg	agtaacagaa	gcttgatata	aattactcta	tcaggagatg	aaagggactt	900
tgggagagct	aatgtgaaat	tgttttataa	ttcttcagta	gaacagatct	ggatcacagt	960
tttacagtgc	agagatttaa	gttggccctc	tagttatgga	gacactccta	ctgtttctat	1020
aaaaggaata	cttacattgc	ccaaaccagt	gcatttcaaa	tcttcagcca	aggaaggttc	1080
caacgctatt	gaatttatgg	aaacgtttgt	atttgcattt	aaacttcaaa	atctacaac	1140
tgttaagactt	gtatttaaga	ttcaaaccac	gactccaggg	aagaaaacca	ttggagaatg	1200
ctcaatgtca	ctcagaaccc	ttagcacaca	ggaaatggat	tactcttttg	atataacacc	1260
accttcaaaa	atttctgttt	gccatgcaga	acttgaattg	gggacttggt	ttcaagcagt	1320
aaatagcaga	attcagttac	aaattcttga	ggcacggtac	cttccaagct	catcaacacc	1380
tctgactttg	agttttttcg	tgaaggtggg	aatgttttagc	tcgggagagt	tgatttataa	1440
gaaaaagaca	gccttactga	aggcctccaa	tgggaagatc	aagtggggag	agacttatgat	1500
ttttccactt	atacagagtg	aaaaagaaat	tgtttttctc	attaagcttt	acagtcgaag	1560

```

ctctgtaaga agaaaacact ttgtggggcca gatttggata agtgaagaca gtaataacat 1620
tgaagcagtg aaccagtgga aagagacagt aataaatcca gaaaagggtg ttatcagggtg 1680
gcacaaatta aatccatctt gaagacttca cacattaatt tgggtgaagaa cttgacattc 1740
ttttagaaga cttatgattt caatttgcta ccaatgagaa gaggcacaaatc aacaaatttg 1800
tcaatttatg ggggctataa ttatggtata taatgtatct gatagaaaat ttgataagaa 1860
aatgtaatga attttatcag atatccaaag taaaggaaat gtttttaaac tgcaacaaga 1920
gacacagaca gtaaaatcaa agtattatta ggatgactaa ataaattata aagtctgtga 1980
gaatatcaac catagatagt tctttctata ttatgttttt gcttttgtat ttaagcctt 2040
acttagatat tcaaaacctg gtatatcaag tctctgttag tactattggc atttagaaga 2100
ctttaccatt atttcagtc taggcattat tgattaggtc ttggctccac tgtttacctc 2160
ttgctatgta ttttctcccc gtaaaaatga attgaaccat ttcaactatt ttctatattt 2220
ggagaaagtt tgtgcctgtt gttttataat ttttttacct ataagacatc acattatccc 2280
tttgtaagct acttatctcc aaaaaacttc agaaatagaa aactacattt tggcaggaa 2340
aattgaaaac accagaaggt tgaagtttaa ttggaaaccc agaataatac tactttgctg 2400
ttttcttccc tcaaatattt tactatttgt tttatttgga gttaaaataa gagtatcatc 2460
catatggtcc atcctaattc acagaattaa atgagcttaa atagaaaatt cagtatttta 2520
tgataatcac ttcgttttta gtttttaaaa tttagattat tctataattt accgtgtttg 2580
agtattttct catttttttc ataaccatac ctgattatac tgtgtaacaa atattttcta 2640
ttgcagtttt ctttccagta cttattagaa ctcagtattt ggaaataatt tcagcttaat 2700
tgaccataag aactgtggcc aaaaagaaca gttttttgga gaggcagatg acattatacc 2760
tgatttttaga aaatctcact ttatttttgc taataagtag actaagtgtc ctgtgttctc 2820
agtcttccct ttttttctgc ccccatctct actttgtccc aggcagtcag agaaagatgg 2880
tgatatttta ggccaggagt atacctgtct ataacctaa catgccttct ttattccagc 2940
tctatgttcc tgtgtatata attaacattt tcccaataa acacttaatt ctcttttccc 3000
taggtgccat ctctcaagc tacaaaatgt ccacatctta tatcccttt gcttctactg 3060
ctctgatttt gtggtaccag tactctctgc cactgaacat tttgaaatat ttttgtttta 3120
gatttgcaaa aaatgacata taggtcagta ctcacatgga tttttaagat aaatcacctg 3180
tgtgataata ttttgaaatc gagacgaata caacttttaa aaattgtttt taaaataaga 3240
cttttttttt tagagcagtt gttaggttaac agaaaaattg agaggaagat agagatttcc 3300
tttctccctt gacaaagccc tcaacagcct cccaggctat cagtatcctg caccacaatg 3360
gtacatttgt tacaatcaat gaacctactc tgaacatca ttatcatcca aagttcatgg 3420
ttacatttag agtccctct tgggtgtata catgctagag gacaaatata tgatgatag 3480
tatgcatac tataatatag tataagtctc ctgcctaaa catcctctgc aaataaaact 3540
attttaatga aacctaaga agaaaatgta tagaaatagc t 3581

```

<210> 31

<211> 1014

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1832403CB1

<400> 31

```

cccacgcgtc cgcgcggcgc cctcgcggca ggtttcgggc ttcaggacaa ttcgtgatgg 60
cgggggctgg ttcgcgcgct gtatcggggg ggtggcgggg cccacaggcc 120
gcgacctttt cgcggaaggc ctgctggagt tctcgcgacc cgctgtgcag cagctcgact 180
ctcacgtaca cgcgcgcaga gagagccagg tagagctccg ggaacaaatt gacaacctag 240
ccacagaact gtgcgcgata aatgaggatc agaagggtgg cctggatctt gacccctatg 300
ttaagaagct acttaatgcc cggcgacgcg ttgtcttggg taacaacatt ctacagaatg 360
ctcaggaacg actgagacgg ctaaacacac gtgttgccaa ggaaacagcc cgcaggagag 420
caatgctgga ttcgggaatt taccctctg gctccccagg caaataacag atgagcctat 480
ggactcagta gcacaagtac tgttccccag ctgccttgtt tcaacagaca tgcaaagatc 540
ctaggagaca gtcccatag accttcagac attaaaaagg gagcgtaca gtttgtttga 600
agcacttcgt cttaccttct taattatgac tcttcttccc cctacacaca gccagaatga 660
ggttcccaaa ggacttacat taattatgac tcttcttccc tttcacaat gagctgaggc 720
ctctactttt ttttttaagc tgcatacttg aggcttacct tcttcaggac tagttaacca 780
gaggggcttc ctttgtatgt tacatgcctg gttacatggg cctggacagc atgtcctcta 840
cctgtgactt ctcatcttcc tgtttacact ggggatttgg agggggcagg caaagtcaaa 900
gtgaatgacc tctgtccacc cactttttta ttgactggc ttgaatacag tagcagtgtt 960
gatagaatca ttttattcaa taaatactta aaatgataaa aaaaaaaaaa aagg 1014

```

<210> 32

<211> 4195

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1971747CB1

<400> 32

```

gcaagactag gcaacctcca gccagtcocct gggctcggggc gatcctccca gaggtggcac 60
aatggagcga tctccaggag aggggccccag cccagcccc atggaccagc cctctgtccc 120
ctccgacccc actgaccagc cccccgctgc tcacgcaaag ccagaccagc gttctggggg 180
ccaacctgct ggccctggcg cggcggtgga ggccctggcg gtgctgactt cattcggggg 240
gcggttgctg gtgctgatac ctgtgtatct ggccggggca gtgggactca gcgtgggttt 300
cgtgtctctc ggccctgccc tctacctggg ctggcgccgg gtccgcgacg agaaaagaac 360
gagccttcga gcagcagggc agctactgga cgacgaggag cagctcactg cgaaaactct 420
ctatatgagt catcgagagc tacctgcctg ggtcagcttc ccagacgtgg aaaaggctga 480
atggctcaat aagattgtgg cccaggtctg gcccttctct ggccagtata tggagaagct 540
tctggctgaa actgtggctc cggctgttag gggatctaac ccccatctgc aaacatttac 600
atttacacga gtggaactgg gtgaaaagcc attgcgcatac attggagtca aggttcaccc 660
aggtcagaga aaagagcaga tctgtctgga cttgaacata agctatgtag gtgatgtgca 720
gattgatgtg gaagtgaaga aatatttttg caaagcagga gtcaagggca tgcagctaca 780
tggcggtttt cggttgatac tggagccact cattggggac cttcccttcg tgggggctgt 840
gtcaatgttc ttcacccgac gcccgacct gctcactctc tgacaccatg atcatggact ccattgctgc 960
gctggatata ccaggactta gttactggtt gccccttggt cctgaccttc aagatgtggc 1020
tcagtgtcgt tcccctctgc ccaggggcat tattcgaatt cacctgctgg ctgctcgagg 1080
gctgagttcc aaggacaaat atgtgaaggg cctgattgag ggcaagtcag acccatatgc 1140
acttgtgcgt ttgggtaccc agacattctg cagctgtgtc attgatgaag aactcaaccc 1200
acagtgggga gagacttatg aggtgatggt acacgaggtc ccagggcagg agattgaagt 1260
ggaggtgttc gacaaggatc cagataaaga tgactttctg ggcagaatga agtggatgt 1320
aggggaaggtg ttacaggcta gcgttctgga tgattggttc cctctacaag gtgggcaagg 1380
ccaagttcac ttgaggctag aatggctgtc acttttgtca gatgcagaga aactggagca 1440
ggttctacag tgggaattggg gagtctctctc tcgaccagat ccccgcgcag ctgccatctt 1500
agttgtctac ctggatcggg cccaggatct tctctgaag aaggggaaca aggaacccaa 1560
ccctatggta caactgtcaa ttcaggatgt gactcaggag agcaaggctg tctacagtac 1620
caactgcccc gtgtgggagg aagcgttccg gttcttctca caagacctc aaagccagga 1680
gctcgatgtg caagtgaagg atgattccag ggccctgact ttaggagcac tgacgtgcc 1740
tctggccccg ctgctgactg ccccgaaact catcctggac cagtgggtcc agctcagcag 1800
ctctggtcca aactccagac tctatatgaa actagtcact aggatcctgt acttggttc 1860
atcagaaata tgcctcccca cggtgccctg ttgtcctggt gcttgggacg tggcagtgta 1920
gaatccccag agaggcagca gtgtggatgc cccacctcga cctgtgcaca cgactcctga 1980
tagccagttt gggactgagc atgtgcttcg gatccatgta ttagaggccc aggacctgat 2040
tgccaaagac cgtttcttgg ggggactggt gaagggcaag tcagacctc atgtcaact 2100
aaagtgggca ggacgaagct tccggagcca tgtttctcgg gaagatctca atccccgtg 2160
gaatgaggtt tttgaggtga tcgtcacatc agttccagge caagagctag aggttgaagt 2220
ctttgacaag gacttggaac aggatgattt tctgggcagg tgtaaagtgc gtctcaccac 2280
agtcttaaac agtggtctcc ttgatgagtg gctgacctg gaggatgtcc catctggccg 2340
cctgcacttg cgctggagc gtctcaccoc cgtcccact gctgctgagt tagaggaggt 2400
gctcagagtg aatagtttga tccagactca gaagagtgc gagctggctg cggccctgct 2460
atccatctat atggagcggg ccgctgcga aaaggcaca agcacctcag 2520
cccttatgct actctcactg tgggagatag ttctcataaa accaagacta tttcgcaaac 2580
ttcagccccc gtctgggatg agagtgcctc cttctcatc aggaaaccac aactgagag 2640
cctagagttg caggttcggg gtgagggcac tggcgtgctg ggctcattat cctgcccc 2700
ctcagagctc ctctggctg accagctctg cttggaccgc tgggtttacac tcagcagtg 2760
tcaggggcag gtgctactga gaggcacgct agggatcctg gtgtcccagc actcgggag 2820
ggaagctcat agccacagct acagccacag ctctcatcgt ctgagtgaag aaccagagct 2880
ctcgggggga cccctcaca tcacctctc agccccagag ctccggcagc gcctaacaca 2940
tggttacagt ccccttgagg ctccagccgg gccctctggg cagggtgaaac tgactctgtg 3000
gtactacagt gaagaacgaa agctgggtcag catgttctat ggttgccggg ccttcgaca 3060
gaatggacgt gatcctctg atccctatgt gtcactgttg ctactgccag acaagaaccg 3120
aggcaccaag aggaggacct cacagaagaa gaggacctg agtctgaat ttaatgaacg 3180
gttttgagtgg gaactcccc tggatgaggg ccagagacga aagctggatg tctctgtcaa 3240
gtctaattcc tcttctatgt caagagagcg tgagctgctg ggggaaggtg agctggacct 3300
agctgagaca gacctttccc aggggtgtag ccggtggtat gacctgatgg acaacaagga 3360
caagggcagc tctaggagc tggcgagtcc cagcctgact gctctgtctt cctgccttcg 3420
tctcgctcca tcaccgctc aatgtgatga gcctaaagct aggggtccaag ggcagagcct 3480
gtgcccctca gccctttcac ctaacaggcc catattcggg cctttgcctg accaaagaga 3540
agaaccgtat gttcccttta ctgcaaggcc tttatccttc tgggccccctg gggcggggac 3600
ctgagctggc tgttctctgc tttgctctga cattgtctc ccttctctcc aactcctcag 3660
ggccttctgt atctgtgcct ggccagtggc agcactagca gtggtattag cttatgccaa 3720

```

atacagcttt	ggaaggatct	ttttttcttt	aactagatgg	tcaccttctt	ccctaccaca	3780
catgggtggg	aaggtggaca	ggetaacctc	tccagctgtg	agcctcttag	actactgcat	3840
gtagcaaatg	ttcagcagct	caggccccc	tgtccagttc	tgccccact	gtcctcaacc	3900
ctgtcctgaa	aattctactg	ctttgatggc	tggtggccagt	ctcttgtcac	tttggaaact	3960
gaggacgcgt	ggattctact	caagcctcca	agtagtggca	tatcagtctt	ggagctccta	4020
gctgggtgata	cggagagggc	tttggaggac	ttggggacagc	agggccaatt	tttttgccca	4080
agtgcctagg	ctgctaactc	actgactaga	acttaatctg	gtactttaca	gttttgcacc	4140
aactctgcc	agccactgga	tcttacatta	aacatcatac	tcaaaaaaaa	aaaaa	4195

<210> 33

<211> 3343

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2285348CB1

<400> 33

gaaaaagaca	aagatgatct	ggggcctgac	agattctcaa	cactcacaga	tgatcccagc	60
cctagactca	gtgcacaagc	tcaggtggct	gaggatattc	tgacaaaata	caggaatgcc	120
attaaacgga	ccagccccag	tgatggagca	atggcaaaact	atgaaagtac	agaggttatg	180
gggtgatggg	aaagtgcaca	tgattctccc	cgtgacgaag	cactgcagaa	catctcggct	240
gatgatctcc	cagactctgc	aagccaagca	gccccccgc	aggattcagc	tttctcttac	300
agagatgcaa	aaaagaaact	gaggcttgct	ctttgctctg	cggactctgt	tgcttccca	360
gtgctgaccc	attcaacaag	gaatggttta	ccagaccaca	cagaccacaga	agacaatgaa	420
attgtatgct	tcttaaaagt	tcaaatagct	gaagcaatta	atttacaaga	taagaatcta	480
atggctcaac	ttcaagaaac	aatgcgctgt	gtgtgcccgt	ttgataatag	gacttgtagg	540
aaactgctgg	cttcgattgc	tgaggactac	agaaaaagag	ccccatata	tgcttatctc	600
actcgttgct	gacaaggact	acagaccaca	caggctcacc	tggaagggt	attgcaaaga	660
gtttttgctg	acaaagaagt	ggccaatcga	tactttacca	ctgtctgtgt	gagattactg	720
cttgagagca	aagaaaaaga	gatcagggaa	ttcattcaag	acttttcagaa	actcaccgca	780
gctgacgata	aaactgctca	ggtagaagat	tttctgcagt	ttcttttatgg	tgcaatggcc	840
caggatgtca	tatggcaaaa	cgcgagtga	gaacagcttc	aagatgcaca	gctggccatt	900
gagcgaagcg	tgatgaaccg	gattttcaag	ctcgcttct	accctaata	agatggggac	960
atacttcgcg	accaggttct	tcatgaacat	atccagagat	tgtctaaagt	agtgactgca	1020
aatcacagag	ctcttcagat	accagaggtt	tatcttcgag	aagcaccatg	gccatctgca	1080
caatcagaaa	tcaggacaat	aagtgccttat	aaaaccccc	gggacaaaagt	gcagtgcata	1140
ctgagaatgt	gctctacgat	tatgaacctc	ctgagcctgg	ccaatgagga	ctctgtccct	1200
ggagcggatg	actttgttcc	tgtgttggtg	tttgtgttga	taaaggcaaa	tccaccctgt	1260
ttgctgtcta	ctgtgcagta	tatcagtagc	ttttatgcta	gctgtctgtc	tgagaggag	1320
tcctattggc	ggatgcagtt	cacagcagca	gtagaattca	ttaaaaccat	cgatgccga	1380
aagtgaccaa	gaccaaggcc	caccaaggca	cgagactggt	aatcagacaa	acagatctct	1440
gagaagggtgc	atcagctgct	ttgaaggctg	aagattgttt	tgtatgatac	tgacagcat	1500
caggcatttt	aaagcagatc	tttactaaac	agggttaatga	gctaacaagc	aggttctctc	1560
gtctttgggc	tctttctctt	ctgagttgca	tattctattt	tcttgtcccc	aagtagagac	1620
tagtactaca	aaaagggacc	acatttttca	agtattttta	agtataaaaa	acaaaaaaa	1680
aatctcttag	gaaatgtcta	gacctccatt	cttggattcc	ctttctttcc	ttttatttta	1740
aaaaagaaca	gtacccctct	tttaagatgc	tgtcttacat	taatgagcat	ctaattgaaa	1800
gaagggtatga	gttgactgca	ggattagaat	agtggtgcgt	tagtggcatt	atctataaat	1860
acactcacct	aaattgaaag	ctaagaagga	aatgtaaata	taatataat	ttatatttga	1920
tgtaatatgg	acatctgcag	attctaataa	acaaggacta	ttgctgatag	taggctgtga	1980
catactgtct	tgtgaaatgg	tttcttgac	aaaatttaag	ctgagcttaa	aagcaaaaaa	2040
caaaaagtac	acagaaatat	ttattaaaat	gtaatacagt	ttattgaact	ttctaggtat	2100
ggagtttgat	ggacagggt	gcctttaatg	agtggtgaag	tcactaagtc	acttagacat	2160
ctcacctggg	aagtttgtga	gcctgcatta	ggagatagac	tgattaccat	acatgacata	2220
aaaagggaaca	gtggatagct	catactttat	gtgggttctt	ctcctccgaa	ataatatact	2280
gcagaaatcc	cagacagagc	tccttacaaa	cctttaattg	taatataatt	ttgatgatta	2340
ttcacattga	atgcacagac	caagaattca	gtgaatgtca	ttttttaaaa	aactaatttg	2400
tattgtctgc	tctagtata	caagttttac	tagtgataaa	ctattttta	caaccatact	2460
attcttatgg	aaaaaaatat	ctattttggc	aggtttctgt	gccttttatt	ccctctctct	2520
aaaaaaagtc	tgtgttttca	tagtttgggt	tgcattgtat	atcaataatt	aatcaggaat	2580
gggttttggt	gcctgaaaaa	ttggccatgg	aggcacacca	aagcttcaag	cacaagtctt	2640
gtacatgggc	catcactgtc	tggtttcact	tcgtgtgttt	cctaaacaca	tttagctgct	2700
tttttaacaa	actcagcccc	atacttgagt	cccttggtgt	tgaggagcatt	tccaggcatc	2760
ttttaaggga	actgtgacaa	acagcctcgg	gcagatgaac	acggaggctc	tctgttgtct	2820
gtctctgaga	tctttgtgtc	tggaatgcc	taaagatttt	atTTTTTTTT	ctttgttttt	2880

```

atatttatattt attttatattt tttgagacag agtctcaccg tgttgcccag gctggagtgc 2940
aatgggtgcga tcttggtctca ctgcaacctc cacctcccag ttcaagtgat tcccctgcct 3000
cagcctcgcg agtagctagg actacaggcg catgtcacca agcccggcta atttttgtat 3060
tttttagtaga aacgggggttt caccatgttg gccaggatga tcttcaatct cctgacctcg 3120
tgatccaccc gccttggcct cccaaagtgc gggattacaa gcgtgaacca cctgcccag 3180
ccagaaacta gatttttcttt atgcctccac cccttcttat tcatttactt tacaatacca 3240
aaataacaaa ttgcatagga gtgtgggatg tggtttctgc cttctagaga gagatcagaa 3300
ttaaactagt actatgaaat gtccttttga atgttaggtc aag 3343

```

<210> 34

<211> 2456

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2374186CB1

<400> 34

```

tgagtcagca aactccgchg cccgcaagcc cggctcggcc cggccctgct ctgttctgcc 60
cggaggagcc gccattgat cgtgtcctgt gctgaagatg tttccggaac aacagaaaga 120
ggaatttgta agtgtctggg ttcgagatcc taggattcag aaggaggact tctggcattc 180
ttacattgac tatgagatat gtattcatac taatagcatg tgttttacia tgaaaacatc 240
ctgtgtacga agaagatata gagaattcgt gtggctgagg cagagactcc aaagtaatgc 300
gttgtctggt caactgccag aacttccatc taaaaacctg tttttcaaca tgaacaatcg 360
ccagcacgtg gatcagcgtc gccagggtct ggaagatttc ctcagaaaag tcttacagaa 420
tgcacttttg ctttcagata gcagccttca cctcttctta cagagccatc tgaattcaga 480
agacattgag gcgtgtgttt ctgggcagac taagtactct gtggaagaag caattcacia 540
gtttgcctta atgaatagac gtttccctga agaagatgaa gaaggaaaaa aagaaaatga 600
tatagattat gattcagaaa gttcacatcc tgggcttgga cacagtagtg atgacagcag 660
ttcacatgga tgtaaagtaa atacagctcc gcaggaatcc tgaaaaaataa ttctaattgt 720
actatcttag gaatagcaaa ttatgtccag tcatagagaa gaaagcttca taataatata 780
ttcttaccta aagctcactg tcatgatgtt aggtatttaa attcttaaag atgttgggtt 840
gtttattagt ggtattttta tgttgtctta ttttaggtaa gcttctgtgt aaagctaaaa 900
atcctgtgaa tacaatacta tcttttacag gcagacatta ttggtaaaaa agatcttgcc 960
ctccaatgaa atgacttaca tgttttaaaa aaccgagttg gttttattga atttaaaaag 1020
ataggttaact aagtagcatt taaaatcaag atagagcatt ccttcttgta tcagtggggc 1080
agtgttacca taaacacggt gtatatgttg ttaaacccta tgaagagtaa cagtgtagac 1140
cagactgcct ctctcagata tgtgcctgat attttgtgga tacctcccct gcactggcaa 1200
aacactatgc ttttgggtgt tagactgaaa tattttaaga gtatttaacc tttccagat 1260
tctgtttcac gcttagatgg aaatgtatct tatgaataga gacataatga aataattgtt 1320
acatcttaga aaaaacatag atagtgttag taatattact tataactgta atatatagat 1380
tcagaaatac attttcatta tccaaaatca gcttcaacia atggtttctg gagacaaata 1440
atttgttttc attatcattg tataatcagg ttaatgattt attttttgac taaatgtgca 1500
atttcttata actagataac tttcagtatc agtgggtggt acttattact taaatcagag 1560
gaaggatttt ataaagatta ataaatttaa ttttaccat aaatattccc ataatttga 1620
aaaggatgtc gacttgctaa tttcagaaat aattattcat ttttaaaaag ccccttttaa 1680
agcatctact tgaagattgg tataattttc ataaaatgtc ttttttttta gtgtcccaa 1740
gatattcttag ataaactatt ttgaagttca gatttcagat gaggcaacat tttctgtgaa 1800
taattaccca agtttcatcc atgttgaatg gtacaaaata tttctgtgaa actaacagga 1860
agatattttc agataactag gataacttgt tgctttgtta ccagcctaa ttgaagagt 1920
gcagaggcta ctacaaaaag caaccttttc attttcacta agagttaaaa agctattgta 1980
ttattaaaaa gtcttttacia tgcttgtttc aaagaaccaa cagaaaaaaa agctaagaaa 2040
actgagaact aacattaaaa aaattaaaat tagaataaga atgatttctt taatttgtcc 2100
tttttttctt tggcttaaaa cattattaaa tttttgtaaa tattttgatt taatgtgtct 2160
tagatctcca ttatttttaac acaggaaaag aaaagattta gtaatttctt accatgcta 2220
tatgtaaaag tcatgccatc caggcattta agagcgatcc tcatcccttc agcaatatgt 2280
atttgagttc acactatttc tgttttacag cagttttgaa aaacacatac tatgccacca 2340
attgtcatat tattttttaga tgatgtaaca tagccatcaa aattaatatt atgtaatgcc 2400
taatacttag tatgtaaatg tcacgagatc atttttacat taaacgtgaa aaaaaa 2456

```

<210> 35

<211> 2004

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature
<223> Incyte ID No: 2476232CB1

<400> 35

cgcgagacagc	ggtggaggcgc	gatttctctgg	gccccgccct	ctggcgctac	catggcgctt	60
ggcaagagtc	accgggatcc	ctacgcgacc	tccgtgggcc	acctcataga	aaaggctaca	120
tttgctggag	ttcagactga	agattggggc	cagttcatgc	acatctgtga	cataattaac	180
actaccagg	atgggccaaa	agatgcagtg	aaagctttga	agaaaaggat	ttccaaaaac	240
tacaatcata	aagaaatcca	acttaccttg	tcacttattg	acatgtgtgt	gcagaactgt	300
ggtccaagtt	tccagtctct	gattgtgaag	aagggaattg	ttaaagagaa	tttagttaag	360
ctactgaatc	ccagatacaa	cttgccatta	gacattcaga	atagaatctt	gaatttcatt	420
aagacttggt	cacagggctt	cccaggaggt	gtggatgtaa	gcgaagtcaa	agaagtatac	480
ctcgacctgg	ttaagaaagg	cgttcagttt	cctccctcag	aagcagaggc	tgaaacagca	540
agacaagaga	ctgctcaaat	ctcatcaaat	cctccaacat	ctgtccctac	tgaccagct	600
ctttcttctg	taattgctcc	aaagaactcg	actgttacat	tggtcccaga	acagattgga	660
aaactgcaca	gtgaattgga	tatggtgaaa	atgaatgtgc	gagtgatgtc	cgccatattg	720
atggagaata	ctcctgggtc	tgaaaaccat	gaagacatag	agcttctgca	gaaactctat	780
aaaacaggtc	gggagatgca	ggagaggatc	atggacctgc	ttgtggtggt	ggagaacgaa	840
gatgtaactg	ttgagctaat	tcagggtgaat	gaggatttga	ataatgctat	ccttggtat	900
gagaggttta	ctagaaacca	acaaaggatt	ttggagcaaa	ataagaacca	gaagggaagc	960
accaatacta	ccagtgaagc	ttctgcccc	tctcaagatc	tctcgcacct	aagtcccagt	1020
ccccggatgc	ctagggccac	tctgggagaa	ctcaacacca	tgaataatca	actttcaggc	1080
ttaaatttca	gccttccaa	ttctgatgta	acaaacaact	taaaaccag	tcttcatcca	1140
cagatgaact	tgctagcctt	ggagaatata	gagatacccc	cgtttgcccc	aaggaccagc	1200
caaaacctca	cctcaagcca	cgcatatgat	aattttctgg	aacattcaaa	ttcagtgttt	1260
ctacagccag	ttagtctaca	aaccattgca	gcagcaccat	caaaccagag	tctgccacct	1320
ttgccagca	atcatccagc	gatgacaaaa	agtgatctcc	agccacctaa	ttactacgag	1380
gtaatggagt	ttgatccctt	agctcctgct	gtcactacag	aagctattta	tgaagaaatt	1440
gatgctcacc	agcacaaagg	agctcaaaat	gatggtgact	gagaagaaag	tggtatgatca	1500
gctcactacc	acatcaaagg	tgccaactct	ctaaaacgta	gactctgtgc	agctttgaag	1560
cctggaagac	aatacctacc	aacatgtcaa	agccatggtg	gcacatttct	gctataatga	1620
agattaaata	gaataacagt	tccaggataa	cactgatctc	tgacaacagc	atgagatttc	1680
aacagaactt	gtttggaaca	aatactcact	taaaacttca	gcagaagaaa	aattacttag	1740
tccttaggcc	aaccaattta	actgcagtgt	catgtttcac	aggccttctt	acatttagaa	1800
atcgtcacac	agctgtgata	agagtagatt	attttactat	gaaataattc	tgaatagatg	1860
aaagcataaa	atgtgagaaa	ctgaatgtat	tattcaggaa	gaatactgag	tgcccttcatt	1920
taactaaagt	tgaatgtaaa	agtcattttg	cacttcttta	taatcctctg	gtttagaatt	1980
atgaaaaata	gtggcaggct	cgta				2004

<210> 36

<211> 4019

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2503986CB1

<400> 36

ccctggcgac	agtgtcttgc	cggggagtag	tagccgggct	ggtaactgga	gtttgagatt	60
aggagacttt	cagacccttg	tgcaaaaga	ggtgcggtcc	ccgaggcagg	atgaagttaa	120
aggaagttaga	tcgtacagcc	atgcaggcat	ggagccctgc	ccagaatcac	ccattttacc	180
tagcaacagg	aacatctgct	cagcaattgg	atgcaacatt	tagtacaaat	gcttcccttg	240
agatatttga	attagacctc	tctgatccat	ccttggtat	gaaatcttgt	gccacattct	300
cctcttctca	caggtaccac	aagttgattt	ggggggccta	taaaatggat	tccaaaggag	360
atgtctcttg	agttctgatt	gcaggtgggt	aaaatggaaa	tattattctc	tatgatcctt	420
ctaaaattat	agctggagac	aaggaagttg	tgattgcccc	gaatgacaag	catactggcc	480
cagtgaagac	cttggtatgt	aacattttcc	agactaatct	ggtagcttct	ggtgctaatt	540
aatctgaaat	ctacatatgg	gatctaaata	attttgcaac	cccaatgaca	ccaggagcca	600
aaacacagcc	gccagaagat	atcagctgca	ttgcatggaa	cagacaagtt	cagcatattt	660
tagcatcagc	cagtcccagt	ggccggggca	ctgtatggga	tcttaggaaa	aatgagccaa	720
tcatacaagt	cagtgaacct	agtaacagaa	tgcatgttct	tggttggtgc	tgcatcctg	780
atgttgctac	tcagatggtc	cttgccctcg	aggatgaccg	gttaccagtg	atccagatgt	840
gggatcttct	atttgcttcc	tctccacttc	gtgtcctgga	aaaccatgcc	agggggattt	900
tggcaattgc	ttggagcatg	gcagatcctg	aattgttact	gagctgtgga	aaagatgcta	960
agattctctg	ctccaatcca	aacacaggag	aggtgttata	tgaacttccc	accaacacac	1020
agtgggtgctt	cgatattcag	tggtgtcccc	gaaatcctgc	tgtcttatca	gctgcttcgt	1080

ttgatggg	cgt	tatcagtg	tattctatca	tgggaggaag	cacagatgg	taaagacaga	1140
aacaagttga	caagcttttca	tcattctttt	ccacagcaga	ggaatcttga	tcccttttggc	acaggacagc	1200
cccttcctcc	gttacaaaatt	ccacagcaga	ccacagcaga	ctgctcagca	tagtatagtg	ctgcctctga	1260
agaagccgcc	caagtggatt	cgaaggcctg	ttggtgcttc	tttttcattt	ggaggcaaac	1320	
tggttacgtt	tgagaatgtc	agaatgcctt	ctcatcagg	agctgagcag	cagcagcagc	1380	
agcaccatgt	gttcattagt	caggttgtta	cagaaaagga	gttcctcagc	cgatcagacc	1440	
aacttcagca	ggctgtgcag	tcacaaggat	ttatcaatta	ttgccaaaaa	aaaatttgat	1500	
cttctcagac	tgaatttgag	aaaaatgtgt	ggctcttttt	gaaggtaaac	tttgaggatg	1560	
attctcgtgg	aaaatacctt	gaacttctag	gatacagaaa	agaagatcta	ggaaagaaga	1620	
ttgcttttggc	cttgaacaaa	gtggatggag	ccaatgtggc	tcttaaagac	tctgaccaag	1680	
tagcacagag	tgatggggag	gagagccctg	ctgctgaaga	gcagctcttg	ggagagcaca	1740	
ttaaagagga	aaaagaagaa	tctgaatttc	taccctcctc	tggaggaaca	tttaatatct	1800	
ctgtcagttg	ggacattgat	ggtttaatta	ctcaggcttt	gctgacgggc	aattttgaga	1860	
gtgctgttga	cctttgttta	catgataacc	gcatggccga	tgccattata	ttggccatag	1920	
caggtggaca	agaactcctg	gctcgaacc	agaaaaata	ccttcgcaaaa	tcccaaagca	1980	
aaattaccag	gctcatcact	gcagtgggtga	tgaagaactg	gaaagagatt	gttgagtctt	2040	
gtgatcttaa	aaattggaga	gaggcttttag	ctgcagtatt	gacttatgca	aagccggatg	2100	
aattttcag	cctttgtgat	ccttttgggaa	ccaggcttga	aaatgaagga	gatagcctcc	2160	
tgcagactca	agcatgtctc	tgctatatatt	gtgcagggaa	tgtagagaaa	ttagtgtgat	2220	
gttgactaa	agctcaagat	ggaagccacc	cctttgtcact	tcaggatctg	attgagaaag	2280	
ttgtcactct	gcgaaaagct	gtgcaactca	ctcaagccat	ggacactagt	actgtaggag	2340	
ttctcttggc	tgcaaatgtg	agtcagtatg	ccaatttgg	ggcagctcag	ggcagtattg	2400	
ctgcagcctt	ggcttttctt	cctgacaaca	ccaaccagcc	aaatatcatg	cagcttcgtg	2460	
acagactttg	tagagcacaa	ggagagcctg	tagcaggaca	tgaatcacct	aaaattccgt	2520	
acgagaaaca	gcagctcccc	aagggcaggc	ctggaccagt	tgctggccac	caccagatgc	2580	
caagagttca	aactcaacaa	tattatcccc	atggagaaaa	tcctccacct	ccgggtttca	2640	
taatgcattg	aaatgttaat	ccaaatgtct	ctggtcagct	tcccacatct	ccaggtcata	2700	
tgacacacca	ggtaccacct	tatccacagc	cacagcctta	tcaaccagcc	cagccgtatc	2760	
ccttcggaac	aggggggtca	gcaatgtatc	gacctcagca	gectgttgct	cctcctactt	2820	
caaacgctta	ccctaaccac	ccttacatat	cctctgcttc	ttcctatact	gggcagtctc	2880	
agctgtacgc	agcacagcac	caggectctt	cacctaccte	cagccctgct	acttctttcc	2940	
ctctctcccc	ttcctctgga	gcataccttc	agcatggcgg	accaggagct	ccaccatcat	3000	
cttcagctta	tgactgcct	cctggaacaa	caggtacact	gcctgctgcc	agttagctgc	3060	
ctgcgtccca	aagaacaggt	cctcagaatg	gttggaatga	ccctccagct	ttgaacagag	3120	
tacccaaaaa	gaagaagatg	cctgaaaact	tcattgcctcc	tgttcccatc	acatcaccaa	3180	
tcattgaaccc	gttgggtgac	ccccagtcac	aaatgctgca	gcaacagcct	tcagctccag	3240	
taccactgtc	aagccagtct	tcattcccac	agccacatct	tccagggtggc	cagcccttcc	3300	
atggcggtaca	gcaacctctt	gggtcaaacag	gcatgccacc	atctttttca	aagcccaata	3360	
ttgaaggtgc	cccaggggct	cctattggaa	ataccttcca	gcatgtgcag	tctttgcca	3420	
caaaaaaaat	taccaagaaa	cctattccag	atgagcacct	cattctaaag	accacatttg	3480	
aggatcttat	tcagcgtctg	ctttcttcag	caacagaccc	tcaaaccaag	aggaagctag	3540	
atgagccag	caaacgtttg	gagtttctgt	atgataaact	taggggaacag	acactttcac	3600	
caacaatcac	cagtgggtta	cacaacattg	caaggagcat	tgaaactcga	aactactcag	3660	
aaggattgac	catgcatacc	cacatagtta	gcaccagcaa	cttcagttag	acctctgctt	3720	
tcattgccagt	tctcaaagtt	gttctcacc	aggccaataa	gctgggtgtc	taaaaggaca	3780	
gcttctcttc	cactcaatat	tgccattttt	ccaaagaac	atgttaaaaa	aaaaaattat	3840	
aagacattga	ctagtctcca	ttagcatgtt	cagtgatcaa	ccagtcaaga	gcatttacac	3900	
tattttctgct	gatatactca	ccttagaact	gctcagaacc	ctggtgcttt	atttttggtt	3960	
ttaatctttt	gttgccagtg	atgattttcc	tattctgcaa	atagtgcac	tccgcgact	4019	

<210> 37

<211> 1793

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2596566CB1

<400> 37

cgacgcgg	cgtgcggcag	ttgcaggcga	gcaggcgagg	aatcgccgtg	gctcttgg	60
gtttccag	ctggttcgca	ggtgaagaga	tggcgttt	gaagagtggc	tggttgctgc	120
gacagagtac	tattttgaag	cgctggaaga	agaactgg	tgatctgtgg	tcggatgg	180
acctgatcta	ttatgatgac	cagactcggc	agaatatcga	ggataagg	cacatgccaa	240
tggactgcat	caacatccgc	acggggcagg	aatgtcggga	tactcagccc	ccggatggaa	300
agtcaaaaga	ctgcattgtc	cagattgttt	gtcagatgg	gaaaacaatt	agtcttgg	360
cagaaagcac	agatgattgc	ttggcctgga	aatttacact	ccaagattct	aggacaaaca	420

cagcgtatgt	gggctctgca	gtcatgaccg	atgagacatc	cgtgggtttcc	tcacctccac	480
catacacggc	ctatgctgca	ccggccccctg	agcaggctta	tggctatggg	ccatacgggtg	540
gtgcgtaccc	gccaggaaact	caagttgtct	acgctgcgaa	tgggcaggcg	tatgccgtgc	600
cctaccagta	cccatatgca	ggacttttatg	gacagcagcc	tgctaaccac	gtcatcattc	660
gagagcgcta	tcgagacaac	gacagcgacc	tggcactggg	catgctggca	ggagcagcca	720
cgggcatggc	cttaggggtct	ctatttttggg	tcttctaggg	gcctcaagggt	cttgatgtgc	780
atagcttctg	ataaccctgt	gtgcaataat	atgatttgca	gggcattttct	gtttgtgaca	840
aaagttttta	ataatagttt	taatcattcc	tttgaaagta	gtgatgtcat	aattgtacta	900
atccacataa	gtaccacaga	gaagggtttg	aactgtgcta	ttttgttcaa	atgttgactc	960
tccgggggca	ctgggtcatt	ccaagactgt	tcttggtgca	ctctcagaat	accttatttg	1020
agcataccctg	ttttgaaagg	cattttcttt	ttagagttag	gtgtagtgtc	taaggggttaa	1080
tttattttca	tgttatgcca	gtaatatagt	gttgatgccc	tattgagtga	ttgtggcaag	1140
aaaagctaca	gcttctttgc	gtttaacttt	ttcaaaccac	agaccagaac	tggttgcatg	1200
ttacttttagg	agttgtgggt	tggtaagctc	ccagggtactt	cccagggtca	tggtgtgaga	1260
gcccccgccc	tgccctctgg	ggctccacag	gccccctggca	aggccgatgg	ctcaggatga	1320
tggggcacag	cccgcctttg	aacaatcatg	cttcagaaat	ctgcctgacc	ctagctgctg	1380
ctgctgctca	ctttattctt	gtatggcttt	ggtaggcata	cttggagaac	atatcccaac	1440
ttagggaattg	atttaagcct	gagagttaga	gggctttaat	cctttaaaac	ttggagaagc	1500
tggctggggcg	cggtggctca	cgctgtaat	cccagcactt	tgagagaccg	aggcggggcg	1560
atcacgaggt	caggagatcg	agaccatcct	ggctaacacg	gtgaaacccc	atctctacta	1620
aaaatacaaaa	aaattagctg	ggcgtgggtg	caggcgcttg	tgggtcccagc	tactcgggag	1680
gctgaggcag	gagaatagtg	tgaaccagg	aggcggagct	tgcagtgagc	caagatagtg	1740
ccactgcact	tcagcctggg	tgacagagt	agactctgtc	tcaaaaaaaa	aaa	1793

<210> 38

<211> 4427

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2685253CB1

<400> 38

cttcggcgag	acctaccggg	ccctgtctgt	gccggccgtg	agctgcggca	ggacggctgg	60
agattacttc	tctaacagga	tcaccagttt	gtccacaaaa	gcacaatgtc	tcgatcacga	120
caaccccccc	ttgtgacggg	catctctcca	aatgaaggga	taccatggac	gaaggtcaca	180
atcaggggag	aaaatctggg	gactggcccc	accgacctca	taggcttgac	catttggtga	240
cataattgcc	tcctgacggc	agaatggatg	tctgcaagta	aaatagtatg	tcgagtggga	300
caagccaaaa	atgacaaaag	agacattatt	gtcaccacta	agtcagggtg	cagaggaacc	360
tcaacagtct	ctttcaagct	actcaaacct	gagaaaaatg	gcatttttga	tcagtctgct	420
gtgtgggttg	atgaaatgaa	ttattatgat	atgcgcactg	acaggaacaa	aggaattccg	480
ccctgtctct	tacgtctctg	taaccgcgtt	ggcattgaga	ttgaaaaaag	ttaattttcg	540
cagaaggact	tagaaatgct	attccatgga	atgagtgtct	atttttacaag	tgagaatttc	600
tcagcagcct	ggtatcttat	agagaatcac	tcaaacacca	gttttgagca	gctcaaaatg	660
gcagtcacca	acctaaagag	acaggctaac	aagaagagtg	agggcagcct	ggcctatgtg	720
aaaggcggtc	tcgtacatt	cttcgaagca	caggatgcc	tctcagccat	ccatcaaaaa	780
ctagaagcag	atggaacgga	aaaagtagaa	ggatccatga	cgcagaaact	ggagaatgtt	840
ctgaacagag	caagtaatac	tgcagacaca	ttgtttcaag	aagtattagg	tcggaaagac	900
aaggcagatt	ccactagaaa	tgcactcaat	gtgcttcagc	gatttaagtt	tcttttcaac	960
cttctcttaa	atatgaaag	gaatattcaa	aagggtgatt	atgatgtggt	tattaatgat	1020
tatgaaaagg	ccaagtcact	ttttgggaaa	acggaggtgc	aagttttcaa	gaaatattat	1080
gctgaagttag	aaacaaggat	tgaagcttta	agagaattac	ttctggataa	attgcttgag	1140
acaccatcaa	ctttacatga	ccaaaaacgt	tacataaggt	acctgtctga	ccttcatgcg	1200
tctggtgacc	ctgcttgcca	atgcattgga	gcccacaca	agtggatcct	tcagctcatg	1260
cacagttgca	aagagggtca	cgtgaaagat	ctgaaaggta	acccaggcct	gcacagtccc	1320
atgttggatc	ttgataatga	tacacgtccc	tcagtgttgg	gccatctcag	tcagacagcg	1380
tcctgaaga	ggggcagcag	ctttcagctc	ggtcgagacg	acacgtggag	atacaaaact	1440
ccccacaggg	tggcctttgt	tgaaaaattg	acaaaactcg	tcttgagcca	gctgcctaac	1500
ttctggaaac	tctggatctc	ctacgttaat	ggaagcctct	tcagtggagac	tgctgagaag	1560
tcaggccaga	ttgaaagatc	aaagaatgta	aggcaaagac	aaaatgattt	taagaaaaatg	1620
attcaggaag	tcagtcactc	cctgggtgaag	cttaccgcgc	gagccctgct	tcccctcagc	1680
atccgggatg	gggaagccaa	gcagtacgga	ggctggggag	tgaagtgcga	gctctccgga	1740
cagtggctcg	ctcacgccat	ccagactgta	agacttactc	atgaatcggt	gactgccctt	1800
gaaattccta	atgacctgtt	acagactatc	caggatctca	tcttggatct	ccgagtacgt	1860
tgcgtaatgg	ccacgttgca	gcacacggcg	gaagaaataa	agagattagc	tgaaaaagaa	1920
gactggattg	ttgacaatga	aggactgact	tctctacat	gtcagtttga	acagtgcac	1980

gtgtgtttctc	tgcagtcact	gaaggggggtt	ctggagtgca	agccggggaga	ggccagtgte	2040
ttccaacaac	ctaaaacaca	ggaggagggtt	tgccagctaa	gcatcaatat	aatgcagggt	2100
tttatatact	gtctggaaca	gttgagcacc	aagcctgatg	cagatataga	tactacacat	2160
ctctctgttg	atgttttctt	ccctgacttg	tttggaagta	tccatgaaga	cttcagcttg	2220
acctcagaac	agcgcccttt	gatagtccta	agtaattgct	gctatctaga	acgtcacacc	2280
ttcctaaata	tgcgagaaca	ttttgaaaag	cacaacttcc	agggaataga	aaaaatcaca	2340
caggttagca	tggccctcatt	gaaagaacta	gatcaaagac	tctttgaaaa	ttacatcgag	2400
ttgaaagcag	atcccatcgt	tggctcctta	gaacctggaa	tttatgcagg	atattttgat	2460
tgggaaggact	gcctgcctcc	aacagggtgt	agaaactatt	taaaagaagc	actgggtgaat	2520
ataattgccc	tgcattgcaga	ggtgttcacc	atttccaaag	aactgggtccc	tcgggtacta	2580
tccaagggtga	tagaagcagt	ttctgaagag	ctcagtcgac	tgatgcagtg	tgtttcatcc	2640
ttcagcaaaa	atggagcttt	acaggcgaga	cttgaaatct	gtgctttgag	ggacactgtg	2700
gctgtttacc	tgacacccga	aagcaagtca	agttttaagc	aggctttgga	agccctgccc	2760
cagctttcca	gtggagcaga	taaaaagtta	ctggaagagc	tcctgaacaa	gttcaagagt	2820
agcatgcact	tgcagctcac	ctgtttccaa	gcagcttctt	caaccatgat	gaaaacataa	2880
atatctgcca	cataaaagaa	gtccaggaaa	ataacacgta	ataagactgt	tcactctcta	2940
agtaccccta	aggtatttgg	tgtattaaac	attgggtttg	ccatttttct	cttttttctt	3000
cctctgactt	cgaaaattgt	tggatacat	ttcaacccaa	atgacctcat	ttgaaatgcc	3060
caggaagtat	ttgtttttgcc	attcttacta	gatcagatcc	tgtatgactt	taaaatacat	3120
tttaaaatat	attttgcate	tcagcagaag	gttgagtgtc	gcaggaagag	ctgcttctgc	3180
agggagtttt	ctggaatggc	gtggcacata	ttgaagcatt	tctcagtgte	ccgttcacgg	3240
aagcgcaggc	agctcccacg	acacacggag	aaactgactg	atacgagatt	tggaaagcta	3300
tgtagacatc	tttgagctc	ttactgtcct	aaactgtaca	gctgtgctta	aaacccttat	3360
ttcatataaa	tggccttaag	ttttcttaatt	caagcgggtt	tttgaaaaaa	tttatggtct	3420
ccattaaaaat	acataattaca	actggggtag	attattttgt	gtccaggtgt	ctgtgattta	3480
actttgcgtt	ttgctatctg	attttttatt	ttcacagggc	taagcatgag	ctttcattct	3540
cactcactct	taatttgcg	tgcgtcacta	cacatgcacg	tgttgcagtc	cctgaggccc	3600
tgtgtgttat	ctgtgatgga	gtgtgaatgt	gtaacgggca	ctgtgttaca	ctctcaggtg	3660
ttggcggggc	ggtcgcagac	ttcagggtcc	cctaacggaa	aggccaggct	ccgcgtggac	3720
ggccaactcc	ctgcccgcct	cttcagcagg	tgactgtctc	tgccacttct	tacctgctga	3780
aggatcttgc	tcagtagctg	gaacaatgct	gctgtcacac	agtctcttct	ctgaaacttc	3840
aggatgctcc	ttggtcacca	ggcaatggga	gctgtagacc	agccgcattg	acttgcccca	3900
cattcactgc	tgactggctt	cactggaata	ggttcaggtc	accgggactt	ctctcagagt	3960
cagcagccca	cccaactccc	tcataccgtc	gcatctgaaa	attttcagag	aggaattctc	4020
tttgtagtcg	gcttgtcagg	gttttcaagt	ttttctatgg	ttatttttaa	atctcttttt	4080
ttaaatgcta	aaattaatgt	cctcattaat	gcaagtaatt	ttagagcaca	gtgtagccca	4140
gtacagtttc	cattattttaa	aataaataaa	tgtgttgggt	catttgggtg	atggggggac	4200
atcatggtaa	ctggaaatga	aagtttaatc	tttcacaaat	gtgcttgata	aagctgttaa	4260
gaattgtctt	tcctaagtat	gttaactgtt	gtatctaact	caactgtgag	ctaagtttaa	4320
aaactgcaat	catgaatgtt	tgtgatattt	taaatatcta	ctacagcaca	ttcagccaat	4380
aaagaatctg	cctttaatgt	aagttgtaat	ttacttgaaa	aaaaaaa		4427

<210> 39

<211> 1583

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2762252CB1

<400> 39

gagctgaaag	ccaggaacat	ccgaggagaa	gagaaagctt	ccagccctcc	ttccttcacc	60
ctggaaatcc	agacaccccc	acccccaccc	tcagatcact	ttaagataat	ttctttatct	120
gtttgcccga	cagaccatgg	ctcccttttg	aagaaacttg	ctaaagactc	ggcataaaaa	180
cagatctcca	actaaagaca	tggattcaga	agagaaggaa	attgtgggtt	gggtttgcca	240
agaagagaag	cttgtctgtg	ggctgactaa	acgcaccacc	tctgtctgat	tcattccaggc	300
tttgcttgag	gaacatgagg	ctacgttttg	agagaaacga	tttcttctgg	ggaagcccag	360
tgattactgc	atcatagaga	agtggagagg	ctccgagagg	gttcttctcc	cactaactag	420
aatcctgaag	ctttggaaa	cgtggggaga	tgagcagccc	aatatgcaat	ttgttttggg	480
taaagcagat	gcttttcttc	cagttccttt	ctggcggaca	gctgaagcca	aattagtgc	540
aaacacagaa	aaattgtggg	agctcagccc	agcaaactac	atgaagactt	taccaccaga	600
taaacaacaaa	agaatagtca	ggaaaacttt	ccggaaactg	gctaaaatta	agcaggacac	660
agttttctcat	gatcgagata	atatggagag	attagttcat	ctgatcattt	cccaggacca	720
tactattcat	cagcaagtca	agagaatgaa	agagctggat	ctggaaattg	aaaagtgtga	780
agctaagttc	catcttgatc	gagtagaaaa	tgatggagaa	aactatgttc	aggatgcata	840
tttaatgccc	agtttcagtg	aagttgagca	aaatctagac	ttgcagtatg	aggaaaacca	900

gactctggag	gacctgagcg	aaagtgatgg	aattgaacag	ctggaagaac	gactgaaata	960
ttaccgaata	ctcattgata	agctctctgc	tgaaatagaa	aaagaggtaa	aaagtgtttg	1020
cattgatata	aatgaagatg	cgggaagggg	agctgcaagt	gaactggaaa	gctctaattt	1080
agagagtgtt	aagtgtgatt	tggaagaaag	catgaaagct	ggtttgaaaa	ttcactctca	1140
tttgagtggc	atccagaaag	agattaaata	cagtgaactca	ttgcttcaga	tgaaagcaaa	1200
agaatatgaa	ctcctggcca	aggaattcaa	ttcacttcac	attagcaaca	aagatgggtg	1260
ccagttaaag	gaaaacagag	cgaaggaatc	tgaggttccc	agtagcaatg	gggagattcc	1320
tccctttact	caaagagtat	ttagcaatta	cacaaatgac	acagactcgg	acactgggat	1380
cagttctaac	cacagtcagg	actccgaaac	aacagtagga	gatgtggtgc	tggtgtcaac	1440
atagttocaa	tggtctcttt	ctgacctgct	ttcatgtttt	aatgtttgtt	taatttaata	1500
ggaaacctca	ttttaaatat	aacactcaaa	aaaaatgtaa	atcatattgt	agtattcaat	1560
agttataaaa	aactcgagaa	atg				1583

<210> 40
 <211> 1416
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3452009CB1

<400> 40						
gtccggcg	ttacagcgga	ggcctaggtg	gcagacaggg	ggcccgggcc	gctgcgtgtt	60
gtccacccaa	gatggagttc	ctcctgggga	accggttcag	cacaccagtg	gggcagtgcc	120
tggaaaaggc	aacagatggc	tccctgcaaa	gtgaggattg	gacgttgaat	atggagatct	180
gtgacatcat	caatgagacg	gaggaagggc	caaaggatgc	cattcgagcc	ctgaagaagc	240
ggctcaacgg	gaaccggaa	tacagagagg	tgatgtctgg	attaacagtg	ctggagacat	300
gtgtgaagaa	ctgtggccac	cgttccaca	tccttgtggc	caaccgagat	ttcatcgaca	360
gtgttcttgt	caaaattata	tctcccaaga	acaacctcc	caccattgta	caggacaaag	420
tgcttgctct	gatccaggca	tggtctgatg	ccttttogaag	cagtcctgat	ctcaccggcg	480
ttgtgcacat	atataggagg	ctgaagagga	aaggggttga	atttcccatg	gcagacttgg	540
acgtctgtgc	tcccatacac	acaccacagc	ggagtgtccc	tgaagtggat	ccagctgcga	600
ccatgcccag	gtcccaatca	cagcagagga	caagtgtctg	ttcctattcc	tcgccgcctc	660
ctgtcccta	ctccgcaccg	caggcccccag	ctctgagtg	gactggcccc	atcacagcca	720
attcagaaca	gattgccagg	ctgcggagtg	aactggacgt	cgttcgagga	aacacaaaag	780
tcattgtctga	gatgttaaca	gaaatggctc	ctggacagga	ggattcatct	gatctggagt	840
tgctgcagga	gctcaacagg	acctgtcggg	ccatgcagca	gcgcacgtg	gagctcatct	900
ccgcgtgtc	caatgaggag	gtcaccgagg	agctgtctga	tgtgaacgat	gacctcaaca	960
acgtcttctc	tgcatacgag	aggtggggag	cggatttctt	tttttttttt	tttcccttga	1020
aaagattgtt	gccataacag	agggatttct	ttcataaaca	aaggaaagag	agggagaggg	1080
agagagatta	gagttggaaa	atactcatga	gagatgctaa	tttgttccc	gctgcagga	1140
aaaatgcttc	ttgtctctc	gggaactgag	cagcagaagc	agcaggaagt	tgacgtgcag	1200
aattaggctc	tgccctttgt	aagcctcagg	ctgctctgcc	tttcccact	gccccgccac	1260
ccccagcccc	agggaggggt	agcacttggg	ggcctctggg	ccccagagct	tctgcacctg	1320
ctcctttgat	gatcctttga	atgtgaaaag	gattaaaggg	aaaaaaaaatc	tctagtaaga	1380
aaaaggattt	cctctttccc	atctgtgaaa	aaaaaa			1416

<210> 41
 <211> 1662
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4644780CB1

<400> 41						
cgatgccggc	ggtcagtggt	ccaggctcct	tattctgcct	tctcctcctg	ctcctggacc	60
cccacagccc	tgagacgggg	tgctctcttc	tacgcagggt	tgagtacaag	ctcagcttca	120
aaggcccaag	gctggcattg	cctggggctg	gaataccctt	ctggagccat	catggagacg	180
ccatcctggg	cctggaggaa	gtgcggctga	gcacatccat	gaggaaccgg	agtggcgccg	240
tgtggagcag	ggcctctgtc	cccttctctg	cctgggaagt	agaggtgcag	atgaggggtga	300
cgggactggg	gcgcggggga	gcccagggca	tgccctgtgt	gtacaccggg	ggcagggggc	360
atgtaggctc	tgctcttggg	gggctggctt	cgtgggaagg	catcgggatc	ttctttgact	420
ctccggcaga	ggatactcag	gacagtcctg	ccatccgtgt	gctggccagc	gacgggcaca	480
tcccctctga	gcagcctggg	gatggagcta	gccaaagggt	gggtcctgtg	cattgggact	540

tccggaaccg	gccacacccc	ttcagagcac	ggatcaccta	ctgggggcag	aggctgcgca	600
tgctccttgaa	cagtggcctc	actcccagtg	atccagatga	tcattgatgtc	ctgtccttcc	660
tgaccttcag	cctgagttag	cccagcccag	agggttcccc	tcagcccttc	ctggagatgc	720
agcagctccg	cctggcgagg	cagctggaag	ggctgtgggc	aaggctgggc	ttgggcacca	780
gggaggatgt	aactccaaaa	tcagactctg	aagctcaagg	agaaggggaa	aggctctttg	840
acctggagga	gacgctgggc	agacaccgcc	ggatcctgca	ggctctgcgg	ggctctctcca	900
agcagctggc	ccaggetgag	agacaatgga	agaagcagct	ggggccccc	ggccaagcca	960
ggcctgacgg	aggctggggc	ctggatgctt	cctgccagat	tccatccacc	ccagggaggg	1020
gtggccacct	ctccatgtca	ctcaataagg	actctgccaa	ggctcgggtgc	ctgctccatg	1080
gacagtggac	tctgtctccag	gccctgcaag	agatgaggga	tgacagctgtc	cgcatggctg	1140
cagaagccca	ggtctcctac	ctgcctgtgg	gcattgagca	tcattttctta	gagctggacc	1200
acatctctggg	cctcctgcag	gaggagcttc	ggggcccgcc	gaaggcagca	gccaaaggcc	1260
cccgcctacc	tggcgagccc	ccaaggccct	cctcgtgcct	gcagcctggc	atcttctctgt	1320
tctacctcct	cattcagact	gtaggcttct	tgggtacgt	gcacttcagc	aggcaggagc	1380
tgaacaagag	ccttcaggag	tgtctgtcca	caggcagcct	tcctctgggt	cctgcaccac	1440
acacccccag	ggccctgggg	attctgagga	ggcagcctct	ccctgccagc	atgcctgcct	1500
gacccacctc	agagcctgct	ttgcatcact	gggaagcagg	cagtgtcttg	gggtgggggt	1560
tggtcagtat	cctctccgtc	tgggtgcccc	gtccccacgc	acacctgagc	tttcggcatg	1620
ctccacctc	gttaaagggtg	atttcctctc	caaaaaaaaa	aa		1662

<210> 42

<211> 3387

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4946103CB1

<400> 42

ccggtctccg	ttttggaaga	cccgcctcgg	cacagccagg	ctcagtcagg	cettgcggta	60
agccttcggc	cgcggtcgcc	cggtagtccc	ggcgggcgcg	gacagacgag	ctgacaggca	120
ccagggtcta	aggcggtccc	tcagtcgggc	tgctgtctcc	acgcctgggg	tcgggcaccg	180
ctcctttctga	ccttcctttc	ccggtttgtc	ccgctgagaa	aagatgacag	caatcaagca	240
tgcatataca	agagacattt	ttacacccaa	tgatgaacgc	ctgctgagca	ttgtgaatgt	300
ctgcaaagca	ggaaaaaaga	aaaagaactg	ttttttatgt	gccacagtga	caactgaacg	360
ccctgtgcag	gttaagggtg	tcaaagtcaa	gaaatccgat	aaggagagatt	tctacaaaaag	420
gcagattgca	tgggcccctc	gagatcttgc	tgtggtagat	gccaaaagatg	ctatcaaaga	480
aaatcctgaa	tttgatttac	actttgaaaa	aatatataaa	tgggttgcca	gcagcactgc	540
tgaaaagaat	gcattttattt	catgcatttg	gaaattgaat	cagcgatatc	tccggaagaa	600
aattgatttt	gtcaatgtta	gtcacagct	tttggaagaa	tctgttccaa	gtggagaaaa	660
tcagagtgtg	acaggaggtg	atgaagaagt	agttagtgaa	taccaagagt	taaatgcaag	720
agaagaacag	gatatcgaaa	taatgatgga	aggctgtgaa	tatgcaatct	cgaatgcgga	780
acgcttttga	gaaaaattgt	ccagagagct	gcagggtgcta	gatgggggcta	acatccagtc	840
aatcatggca	tctgaaaaac	aagtcaacat	cctgatgaaa	ttgctagatg	aggctctaaa	900
ggaggtagat	cagattgaat	tgaactgag	cagttatgag	gaaatgtctc	aaagtgtaaa	960
agaacaaatg	catagatctc	ctgaagcaga	ccacctaat	catcttagta	acactaataa	1020
tgtaaaactc	ctatctgaga	tagagtctct	tgtgaaccac	atggacttgg	ccaaagggtca	1080
tataaaggcc	cttcaggaag	gagatcttgc	ttcttccaga	ggcattgagg	cctgcaccaa	1140
tgtgtctgat	gcccttctgc	agtgcattga	tgtagctctt	cgaccaggcc	atgacttgct	1200
tctggcagtc	aaacagcaac	agcagcgatt	cagtgatttg	cgagagcttt	ttgccgggag	1260
actggccagt	cacctcaaca	atgtttttgt	tcaacagggc	cacgatcaga	gttcgtctct	1320
cccccagcac	tgtgtctcaa	ccgggtttac	ccaatcatca	tccatttcac	agagatttcc	1380
tccgattgcc	aagctgatgg	agtggctaaa	gagtacagat	tatggaaaat	atgaaggact	1440
aacaaagaat	tacatggatt	atztatcccg	actatatgaa	agagaaatca	aagatttctt	1500
tgaagttgca	aagatcaaga	tgactggcac	aactaaagaa	agcaagaagt	ttggtcttca	1560
tggaaagttcg	gggaaattaa	ctggatctac	ttctagtcta	aataagctca	gtgttcagag	1620
ttcagggaat	cgcagatctc	agtcattctc	cctgtttggat	atgggaaaca	tgtctgcctc	1680
tgatctcgat	gttgccgaca	ggaccaaaat	tgataagatc	tttgaacagg	tactaagtga	1740
actggagccc	ctatgtcttg	cagaacagga	cttcataagt	aaatttttca	aactacagca	1800
acatcaaagt	atgcctggaa	ctatggctga	agcagaggac	ctggatggag	gaacattatc	1860
acggcaacat	aattgtggga	caccactgcc	tgtttcatct	gagaaagata	tgatccgcca	1920
aatgatgatt	aaaatatttc	gctgcattga	gccagagctg	aacaacctaa	ttgcattagg	1980
agacaaaatt	gatagcttta	actctcttta	tatgttagtc	aaaatgagtc	atcatgtgtg	2040
gactgcacaa	aatgtggacc	ctgcttcttt	cctaagtact	acattgggaa	atgttttggg	2100
gactgtcaaa	aggaactttg	acaaatgcat	tagtaaccaa	ataaggcaaa	tggaaagaagt	2160
aaagatctca	aaaaagagta	aagttggaaat	tcttccattt	gttgctgaat	ttgaagaatt	2220

tgctggactt	gcagaatcaa	tcttcaaaaa	tgctgagcgt	cgtggagacc	tgataaaagc	2280
atacaccaaa	cttatcagag	gagtatttgt	taatgtggag	aaagtagcaa	atgaaagcca	2340
gaagaccccc	agggatgtgg	ttatgatgga	aaactttcac	catatttttg	caactccttc	2400
tcgattgaaa	atctcatgtc	tagaagcaga	aaaaaaagaa	gccaaacaaa	aatacacaga	2460
tcaccttcag	tcttatgtca	tttactcttt	aggacaacct	cttgaaaaac	taaatcattt	2520
ctttgaaggt	gttgaagctc	gcgtggcaca	gggcataagg	gaggaggaag	taagttacca	2580
acttgcattt	aacaaacaag	aacttcgtaa	agtcattaag	gagtaccctg	gaaaggaagt	2640
aaaaaaaggt	ctagataacc	tctacaagaa	agttgataaa	catttatgtg	aagaagagaa	2700
cttacttcag	gtgggtgtgg	actccatgca	agatgaattt	atacgccagt	ataagcactt	2760
tgaagggttg	atagctcgct	gttatcctgg	atctgggtgt	acaatggaat	tcactattca	2820
ggacattctg	gattattgtt	ccagcattgc	acagtcccac	taaaccttgt	gaaagaagaa	2880
aagataactg	aatgaagcat	ttgagtataa	cagacactat	acaaaaatac	caagcaactg	2940
ttttgagaac	ccagacttaa	aattttatgt	attattaaat	gttagataaa	tgggtagtac	3000
catactacaa	atattttaa	gcaaaattac	caacctatat	agcagtttta	tttgccctat	3060
aggttgcata	ctaacttaag	cattcatgtc	accataaaat	gccttttagc	tttctcaatg	3120
actggatggg	aaattttcct	ttattgccta	gctgcttggt	tttgagtggg	tgctctatga	3180
gcaatgcatt	tggagttctt	cagctttcac	tacttctctg	ttgcttgcta	atcatgtaac	3240
tactaaaata	ctgtacaaaa	ttgttttttc	acactaacaa	atgtgtatat	ggagaagagg	3300
gctcatgtga	tgatcatttg	tgaacttaga	tttttgagga	ttatgtgact	agtaataaat	3360
gtgaaataaa	ttttcaaaaa	aaaaaaa				3387

<210> 43

<211> 1043

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5562355CB1

<400> 43

gcgggcgaca	tggaacaacgc	ggggaaggag	cgtgaggcag	tacagctgat	ggcggaggcc	60
gagaagcgag	tcaaggcctc	ccactccttc	ctccgagggc	tgtttgaggg	aaacacaaga	120
atagaagagg	cttggtgaaat	gtataccaga	gctgcaaata	tgttcaagat	ggctaaaaat	180
tggagtgcctg	caggaaacgc	attttgtcag	gcagccaagc	tcacatgca	gcttcagagc	240
aaacatgact	ctgctaccag	ctttgtggat	gctggaaatg	cttacaacaa	ggcagatccc	300
caagaggcta	tcaactgctt	aaatgcagcc	atcgacattt	acacagacat	gggaagggtt	360
acaattgcag	ccaagcacca	cattactatt	gcagagatct	atgagactga	actgtagac	420
attgagaagg	ctattgcaca	ttatgaacaa	tctgctgatt	attacaaagg	agaagaatcc	480
aacagctcag	caaacaagtg	tctgctgaag	gtggcagcat	atgctgccc	tcttgagcag	540
taccagaacg	ccattgagat	ctatgagcag	gttggggcaa	acacaatgga	taatcctttg	600
acgacataca	gtgcaaagg	ttacttcttc	aaagctgccc	tctgccactt	catagtagac	660
gagttgaatg	ccaagcttgc	tcttgagcaa	tatgaggaca	tgtttccagc	atttactgat	720
tcaagagaat	gtaaattatt	gaaaaaactc	ctagaagctc	atgaagaaca	gaacagtga	780
gcttacactg	aagcagtga	ggaatttgac	tcaatatctc	gcttggatca	gtggctgacc	840
accatgttgc	ttcgcacaa	aaagtccatc	caaggggatg	gagaaggaga	tggagacct	900
aaatgaaatg	ttttgtctt	tgtggcatgc	agctaactcc	tctttagttt	tgtcttaggg	960
tcaagtgate	tttatgggat	gcctatttaa	tggcttaatt	ttgttgcata	tgagccagac	1020
ggcctgtgta	ttgtttaagc	tgc				1043

<210> 44

<211> 3207

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5678824CB1

<400> 44

ccggagggtgg	gagccctggg	ccaaaatggc	ggcctacctg	cagtggcggc	gcttcgtttt	60
cttcgacaag	gagctggtga	aggagccgct	gagcaatgat	ggggccgctc	ccggggccac	120
acctgcttct	ggatccgctg	cttccaagtt	cctttgcctc	cctcctggca	tcactgtctg	180
cgactcaggc	cgaggagacc	tggtcttttg	agatatggaa	ggccagatct	ggttcttgcc	240
acgttcccta	cagcttacag	gcttccaagc	ctacaaacta	cggttgacac	acctgtacca	300
actgaagcag	cacaatatc	tggcatctgt	tggagaagat	gaagagggca	tcaaccctt	360
ggttaagatc	tggaaacctg	agaagagaga	tgttgccaat	ccactctgca	ctcgaatctt	420

```

ccctgtctatt ccaggaacag agccaactgt tgtatcttgt ttgactgtcc atgaaaatct 480
caacttttatg gccatttggt tcacagatgg cagtgtttaca ttgaacaaag gagacatcac 540
ccgggaccgg catagcaaga cccagatttt gcacaagggc aactatcctg taactggatt 600
ggccttttcgc caagcaggaa agaccactca cttgttttgt gtgacaacag agaacgtcca 660
gtcctatata gtttctggaa aagactaccc tcgcgtggag ttggacaccc atgggtgtgg 720
cctgcgtgc tcagccctaa gtgacccttc tcaggacctg cagttcattg tggccgggga 780
tgagtgtgtc tacttgtacc agcctgatga acgtgggccc tgcttcgct ttgagggcca 840
taagtcatt gccactggt ttagaggcta ccttatcatt gtctcccggt accggaaggt 900
ttctcccaag tcagagtta ccagcaggga ttcacagagc tccgacaagc agattctaaa 960
catctatgac ctgtgcaaca agttcatagc ctatagcacc gtctttgagg atgtagtggg 1020
tgtgcttgct gagtggggct ccctgtactg gctgacgagg gatggggggg tccacgcact 1080
gcaggagaag gacacacaga ccaaactgga gatgctgttt aagaagaacc tatttgagat 1140
ggcgattaac cttgccaaaga gccagcatct ggacagtgat gggctggccc agattttcat 1200
gcagtatgga gaccatctct acagcaaggg caaccacgat ggggctgtcc agcaatatat 1260
ccgaaccatt ggaaagtgtg agccatccta tgtgatccgc aagtttctgg atgccacg 1320
cattcacaac ctgactgcct acctgcagac cctgcaccga caatccctgg ccaatggcca 1380
ccataccacc ctgctcctca actgctatac caagctcaag gacagctcga agctggagg 1440
gttcatcaag aaaaagagt agagtgaagt ccactttgat gtggagacag ccatcaaggt 1500
cctccggcag gctggctact actcccatgc cctgtatctg gcggagaacc atgcacatca 1560
tgagtgttac ctgaagatcc agctagaaga catthaagaat tatcaggaag ccttcgata 1620
catcggaag ctgccttttg agcaggcaga gagcaacatg aagcgctacg gcaagatcct 1680
catgcaccac ataccagag agacaactca gttgctgaag ggactttgta ctgattatcg 1740
gccagcctc gaaggccgca gcatagggga ggccccaggc tgcagggcca actctgagga 1800
gttcatcccc atctttgcca ataaccgcg agagctgaaa gccttcctag agcacatgag 1860
tgaagtgcag ccagactcac cccaggggat ctacgacaca ctccttgagc tgcgactgca 1920
gaactggggc cagcagaagg atccacaggt caaagagaag cttcacgcag aggccatttc 1980
cctgctgaag agtggctcgt tctgcaagc ccttgacaag gccctgggtc tgtgcagat 2040
gcagcacttc caggatggtg tctttacct ttatgagcag ggaagctgt tccagcagat 2100
catgcactac cacatgcagc acgagcagta ccggcaggtc atcagcgtgt gtgagcgcca 2160
tggggagcag gaccctcct tgtgggagca ggccctcagc tacttcgctc gcaaggagga 2220
ggactgcaag gagtgtgtg cagctgtcct caagcatatc gagaacaaga acctcatgcc 2280
acctcttcta gttgtgcaga cctgggcca caactccaca gccacactct ccgtcatcag 2340
ggactacctg gtccaaaaac tacagaaaca gagccagcag attgcacagg atgagctgcg 2400
ggtgcggcgg taccgagagg agaccacccg tatccgccag gagatccaag agctcaaggc 2460
cagtccaaag attttccaaa agaccaagtg cagcatctgt aacagtgcct tggagttgcc 2520
ctcagtcac ttctgtgtg gccactcct ccaccaacac tgctttgaga gttactcgga 2580
aagtgtgct cactgcccc cctgcctccc tgaaaaccgg aaggtcatgg atatgatccg 2640
ggcccaggaa cagaaacgag atctccatga tcaattccag catcagctca ggtgctccaa 2700
tgacagcttt tctgtgattg ctgactactt tggcagaggt gttttcaaca aattgactct 2760
gctgaccgac cctcccacag ccagactgac ctccagcctg gaggtgggc tgcaacgca 2820
cctactatg cactccagga ggggacttta agcagcctgg aggaagatgt gggcaacagt 2880
ggaggaccga gagaacagac acaatgggac ctgggggggc gttacacaga aggctggctg 2940
acatgcccag ggctccactc tcatctaatg tcacagccct cagaactaaa gcggactttc 3000
tttccctgcc ttcttattta gtcagcttgc catccctcct cttcactagc agtgtagatc 3060
attccagatc agtgggggag ggcacctcag caacctctga gtgtggacaa tagctgcttt 3120
cttctctatc caagagcacc aggtgtgtct tgggtccttg ctctcagagt ctataaataa 3180
aagaatataa tgattaaaaa aaaaaaa 3207

```

<210> 45

<211> 2356

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5870962CB1

<400> 45

```

ctgaactaac cctctaacce ttgggagctc gtgtgcagca gtgatgtgag ctgcatccgg 60
gctgaatagg gagctcccgc tatggcactg gctcaaggag ctaactctaa gagaaagtat 120
tttggggaag acatcttggt tgcaccctc attgcatctt gactaattgc ttccaactcc 180
gggggcttcc agggacaatc tcattgttct tttctctcca agggtcagcc atgttaacca 240
tttgtaatt tcttctgaac agatcaagca aaagtagata atcagccaga agaattagtg 300
cgtagtgtc aagatgatga gaaaccagat cagaagccag ttacaaatga atgcgtacca 360
agaatttcca cagtgcctac acaacctgat aatccatttt ctcacctga caaactcaaa 420
aggatggaca agtctgttcc agcatttctc caagatgagg tgagtggcag tgtgatgagt 480
gtttatagtg gagactttgg caatctggaa gttaaaggaa atattcagtt tgcaattgaa 540

```

tatgtggagt	cactgaagga	gttgcattgtt	tttgtggccc	agtgttaagga	cttagcagca	600
gcggatgttaa	aaaaacagcg	ttcagaccca	tatgtaaagg	cctatttgct	accagacaaa	660
ggcaaaatgg	gcaagaagaa	aacactcgta	gtgaagaaaa	ccttgaatcc	tgtgtataac	720
gaaatactgc	ggtataaaat	tgaaaaacaa	atcttaaaga	cacagaaatt	gaacctgtcc	780
atgtggcatc	gggatacatt	taagcgcaat	agtttcttag	gggaggtgga	acttgatttg	840
gaaacatggg	actgggataa	caaacagaat	aaacaattga	gatggtagcc	tctgaagcgg	900
aagacagcac	cagttgccct	tgaagcagaa	aacagagggt	aatgaaact	agctctccag	960
tatgtcccag	agccagtcct	tggtaaaaag	cttcctacaa	ctggagaagt	gcacatctgg	1020
gtgaaggaat	gccttgatct	accactgcta	agggggaagtc	atctaaattc	ttttgttaaa	1080
tgtaccatcc	ttccagatac	aagtaggaaa	agtcgccaga	agacaagage	tgtagggaaa	1140
accaccaacc	ctatcttcaa	ccacactatg	gtgtatgatg	ggttcaggcc	tgaagatctg	1200
atggaagcct	gtgtagagct	tactgtctgg	gaccattaca	aattaaccaa	ccaatttttg	1260
ggaggtcttc	gtattggctt	tggaaacaggt	aaaagttatg	ggactgaagt	ggactggatg	1320
gactctactt	cagaggaagt	tgtctctctg	gagaagatgg	taaactcccc	caatacttgg	1380
attgaagcaa	cactgcctct	cagaatgctt	ttgattgcca	agatttccaa	atgagcccaa	1440
attccactgg	ctcctccact	gaaaactact	aaaccgggtg	aatctgatct	tgaaaatctg	1500
agtaggtgga	caaatatctt	cactttctat	ctattgcacc	taaggaatac	tacacagcat	1560
gtaaaagtca	atctgcatgt	gcttctttga	ttacaaggcc	caagggattt	aaatataaca	1620
aaatgtgtaa	tttgtgactc	taatatataa	taagatatatt	gaacaagcta	ggaaaattga	1680
atttctgctg	ctgcttcaaa	gaaaaagctg	ccccagagca	ttaaactatg	ggtattgtta	1740
agaagcaaaa	tgttcttgtt	tgccatcatg	tgtttcacac	cacaattctg	tgccacagtt	1800
aagagggtct	ggtacccttg	caggaccttt	gtagggtgtg	ggaaaaagtc	gcagaaagat	1860
actcaaatg	gagcagggaa	tggagacaga	catcagtgat	gataaaaaaa	aaaaatggac	1920
cttaagaaac	tatttactct	gtaatctcta	ataaaatatg	gaattccata	ttagggcaat	1980
gagactgaaa	ctactggtgt	ttttctgcct	tgagaaaaca	aacagttaaa	acaagcctca	2040
aatgtatttt	agtgccacc	actggccata	ggtacaattc	agttgttggc	ttgttttgac	2100
tttaattctaa	aagctgtctc	aagcctgtat	ttttatgagt	ttattttttt	aaaacctgc	2160
atatatatga	ttgtttttct	tataacttta	ctatatgaaa	gcagcataag	agtagtcaca	2220
aacatgtttt	gcaacaaagt	tttaattaga	atgtaagttg	ctcagttata	ctgttcttct	2280
tatgtatgta	aaattttctg	attttgtaaa	aacccttaga	ataaattatc	atttgattta	2340
aattgtatta	gaaaaat					2356

<210> 46

<211> 1034

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2818605CB1

<400> 46

gcttttgctgt	tgttgcctct	eggaggcggc	gatccccgaa	ggcgagctga	aatacggctg	60
caggctacaa	tttgcagccg	accattatgg	atgacaagga	gccgaagagg	tggcccaccc	120
tcaggggaccg	cttgtgctcg	gatggcttct	tatttcccca	ataccctatt	aaaccgtatc	180
atctgaaggg	gatccacaga	gctgtcttct	atcgtgatct	ggaggaactg	aagttcgttc	240
tgtcacgcg	ttatgacatc	aataagagag	acaggaagga	aaggaccgcc	ctacatttgg	300
cctgtgccac	tggccaaccg	gaaatggtac	atctcctggg	gtccagaaga	tgtgagctta	360
acctctgcga	ccgtgaagac	aggacacctc	tgatcaaggc	tgtacaactg	aggcaggagg	420
cttgtgcaac	tcttctgctg	caaaatggcg	ccgatccaaa	tattacggat	gtcttttgaa	480
ggactgctct	gcactacgct	gtgtataatg	aagatatact	catgatagaa	aaacttcttt	540
cacatggtac	aaatattgaa	gaatgcagca	aggtataggt	caaccaatgt	tattttcaaa	600
ctatctgaaa	tgcattttatt	ttaacattga	cacatgtaag	ggtcaatttt	tcataatttg	660
aagctcaaac	attccttgaa	tgaaaatatt	ttgaaatgcc	ttaactgtct	aagattttac	720
tttaaatatt	ggaactttta	aagaagcatt	ataggggaaca	gccttttttc	atgcacttat	780
ggtaaaatac	tataaaaaa	aatgaattac	aataaattta	taattcatga	caactgaatt	840
tgggaaaggt	aatagtttaag	tgtttttcca	ctaaattact	ttttttctaa	tcagtgtgaa	900
gtgacacagg	aaagtaaaat	tgtcccttat	aaataggctt	tatttttaaat	gtcaaaagaa	960
attaaagaat	ttcacaataa	atgtacatgt	tgttgcctgt	gacaagtgtt	gtatgtgaag	1020
gtgatttcat	ctga					1034

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number
WO 01/46256 A3

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/47, 16/18, C12Q 1/68, A01K 67/00, A61K
38/17, G01N 33/68

(21) International Application Number: PCT/US00/34919

(22) International Filing Date:
21 December 2000 (21.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/172,968 21 December 1999 (21.12.1999) US
60/172,066 23 December 1999 (23.12.1999) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier applications:
US 60/172,968 (CIP)
Filed on 21 December 1999 (21.12.1999)
US 60/172,066 (CIP)
Filed on 23 December 1999 (23.12.1999)

(71) Applicant (for all designated States except US): **INCYTE
GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo
Alto, CA 94304 (US).

(72) Inventors: and

(75) Inventors/Applicants (for US only): **TANG, Y., Tom**
[CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US).
YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale,
CA 94087 (US). **BANDMAN, Olga** [US/US]; 366 Anna
Avenue, Mountain View, CA 94043 (US). **HILLMAN,
Jennifer, L.** [US/US]; 230 Monrow Drive #17, Mountain
View, CA 94040 (US). **BAUGHN, Mariah, R.** [US/US];
14244 Santiago Road, San Leandro, CA 94577 (US). **LU,**

Dyung, Aina, M. [US/US]; 233 Coy Drive, San Jose, CA
95123 (US). **AZIMZAI, Yalda** [US/US]; 5518 Boulder
Canyon Drive, Castro Valley, CA 94552 (US). **YANG,
Junming** [CN/US]; 7125 Bark Lane, San Jose, CA 95129
(US). **BURFORD, Neil** [US/US]; 105 Wildwood Circle,
Durham, CT 06522 (US). **AU-YOUNG, Janice** [US/US];
233 Golden Eagle Lane, Brisbane, CA 94005 (US).
REDDY, Roopa [IN/US]; 1233 W. McKinley Avenue #3,
Sunnyvale, CA 94086 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics,
Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
27 December 2001

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: VESICLE TRAFFICKING PROTEINS

(57) Abstract: The invention provides human vesicle trafficking proteins (VETRP) and polynucleotides which identify and encode VETRP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing disorders associated with expression of VETRP.

WO 01/46256 A3

INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/US 00/34919

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C07K16/18 C12Q1/68 A01K67/00
 A61K38/17 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal, WPI Data, BIOSIS, EMBL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 46744 A (FIORE PIER PAOLO DI ;ISTITUTO EUROP DI ONCOLOGIA S (IT); PELICCI P) 22 October 1998 (1998-10-22)</p> <p>page 2 -page 4; claims 1-6,10,27,28,32,33; figures 3,7 page 10 page 13 page 19 -page 20 page 22 & DATABASE EMBL 6 January 1999 (1999-01-06) SALCINI A.E. ET AL.: "human Rab-R protein" Database accession no. AF015042 the whole document</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	<p>1,3, 6-16,18, 19,22, 25-28</p>

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

25 April 2001

Date of mailing of the international search report

20.07.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

van Heusden, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/34919

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SALCINI A.E. ET AL.: "Binding specificity and in vivo targets of the EH domain. a novel protein-protein interaction module" GENES AND DEVELOPMENT, vol. 11, 1997, pages 2239-2249, XP002073498 page 2245 ---	1,3, 6-16,18, 19,22, 25-28
X	DATABASE EMBL,22 June 1999 (1999-06-22) HILLIER ET AL.: "WashU-NCI human EST project" Database accession no. AI880211 XP002165492 the whole document ---	3,12
A	SCHIMMOLLER F. ET AL.: "Rab GTPases, directors of vesicle docking" J. BIOL. CHEM., vol. 273, no. 35, 28 August 1998 (1998-08-28), pages 22161-22164, XP002165490 the whole document ---	1-19,22, 25-28
P,A	WATERS M.G. AND HUGHSON F.M.: "Membrane Tethering and fusion in the secretory and endocytic pathways" TRAFFIC, vol. 1, August 2000 (2000-08), pages 588-597, XP002165491 the whole document -----	1-19,22, 25-28

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/34919

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 18 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the composition.
2. ☒ Claims Nos.: 20,21,23,24
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-19,22,25-28 (partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-19, 22, 25-28 (partially)

An isolated polypeptide comprising an amino acid consisting of SEQ ID NO:1, an isolated polynucleotide encoding said polypeptide, an isolated polynucleotide comprising a sequence consisting of SEQ ID NO:24, said polynucleotide in a cell or in an organism and methods of recombinant production of said polypeptide. Methods of detection of said polynucleotide, pharmaceutical composition comprising said polypeptide, method of treating a VETRP-related disease by administering said composition. Methods of screening compounds for binding to or for effectiveness as agonists or antagonists of said polypeptide. Method of screening a compound for altering expression of said polynucleotide. Method of assessing toxicity of a test compound on said polypeptide.

2. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 2 and 25, respectively.

3. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 3 and 26, respectively.

4. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 4 and 27, respectively.

5. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 5 and 28, respectively.

6. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 6 and 29, respectively.

7. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 7 and 30, respectively.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

8. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 8 and 31, respectively.

9. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 9 and 32, respectively.

10. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 10 and 33, respectively.

11. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 11 and 34, respectively.

12. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 12 and 35, respectively.

13. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 13 and 36, respectively.

14. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 14 and 37, respectively.

15. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 15 and 38, respectively.

16. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 16 and 39, respectively.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

17. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 17 and 40, respectively.

18. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 18 and 41, respectively.

19. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 19 and 42, respectively.

20. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 20 and 43, respectively.

21. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 21 and 44, respectively.

22. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 22 and 45, respectively.

23. Claims: 1-19, 22, 25-28 (partially)

As in invention 1 but relating to SEQ ID NO 23 and 46, respectively.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 20,21,23,24

Present claims 20-21 and 23-24 relate to a compound defined by reference to a desirable characteristic or property, namely having agonist/antagonist activity on the VETRP polypeptide of claim 1. The claims cover all compounds having this characteristic or property. The application provides no support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search impossible.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

Information on patent family members

PCi, JS 00/34919

Form PCT/ISA/210 (patent family annex) (July 1992)

THIS PAGE BLANK (USPTO)